

Classifying Non-Periodic Sequences by Permutation Transducers

Hans Zantema^{1,2} and Wieb Bosma^{2,3}

¹ Department of Computer Science, TU Eindhoven, P.O. Box 513,
5600 MB Eindhoven, the Netherlands, email: h.zantema@tue.nl

² Radboud University Nijmegen, P.O. Box 9010,
6500 GL Nijmegen, the Netherlands, email: w.bosma@math.ru.nl

³ Centrum voor Wiskunde en Informatica, Amsterdam

Abstract. Transducers order infinite sequences into natural classes, but permutation transducers provide a finer classification, respecting certain changes to finite segments. We investigate this hierarchy for non-periodic sequences over $\{0, 1\}$ in which the groups of 0s and 1s grow according to simple functions like polynomials. In this hierarchy we find infinite strictly ascending chains of sequences, all being equivalent with respect to ordinary transducers.

1 Introduction

Equivalence under transducers organizes infinite sequences into a hierarchy with interesting properties, as ongoing research is revealing, see for example [3, 6] and the conference paper [5] at DLT 2016. In this setting the main definition is that for two sequences σ, τ we have that $\sigma \geq \tau$ if and only if there exists a transducer T that produces τ when consuming σ . Here a transducer is a deterministic automaton producing output strings on every transition. Two sequences σ, τ are called equivalent, notation $\sigma \sim \tau$ if both $\sigma \geq \tau$ and $\tau \geq \sigma$. A straightforward construction shows that $\sigma \sim u\sigma$ for any sequence σ and any finite string u , so prepending or removing a finite initial word remains inside the class. The pre-order \geq gives rise to an order on the equivalence classes of \sim ; the bottom element in this order consists of the class of ultimately periodic sequences.

In the current paper we investigate a more fine-grained hierarchy on sequences based on an alternative pre-order \geq_p . Here prepending or removing initial segments may change the class, but other basic properties are kept, like $\sigma \geq_p h(\sigma)$ for any morphism h . The idea is that we add the requirement that transducers should be *permutation transducers*. This means that not only for every state and symbol there is exactly one outgoing arrow (as is required by determinism), but also exactly one incoming arrow: it will thus be a permutation automaton (see [1, 7, 8]) with output, just like a finite state transducer is a DFA with output. Our original motivation for permutation transducers was to be able to compare and classify two-sided sequences as was elaborated in [2]. There we

already made some first investigations on ordering (one-sided) sequences by permutation transducers, raising several issues that we worked out in the current paper.

So we define $\sigma \geq_p \tau$ if and only if a permutation transducer P exists such that $P(\sigma) = \tau$, and $\sigma \sim_p \tau$ if and only if both $\sigma \geq_p \tau$ and $\tau \geq_p \sigma$. In [2] we already showed that $0^\omega \not\sim_p 10^\omega$, a clear illustration that initial segments matter in this context. Again the pre-order \geq_p on sequences gives rise to an order on \sim_p -equivalence classes; here the bottom element is the class of all periodic sequences. In [2] we showed that the ultimately periodic sequences that are not periodic form an atomic class. Here the focus is on sequences that are not ultimately periodic. In particular, we look at sequences of the shape

$$\langle f \rangle = 10^{f(0)}10^{f(1)}10^{f(2)} \dots \quad \text{and} \quad \llbracket f \rrbracket = 0^{f(0)}1^{f(1)}0^{f(2)} \dots,$$

for various functions $f : \mathbb{N} \rightarrow \mathbb{N}$, in particular polynomials. Based on ordinary transducers one has $\langle f \rangle \sim \llbracket f \rrbracket$ if $f(n) > 0$ for all $n \in \mathbb{N}$, and for all linear f, g it holds $\langle f \rangle \sim \langle g \rangle$. A main result of [4] states that the class containing the sequences of the shape $\langle f \rangle$ for f linear is *atomic*, that is, if $\langle f \rangle \geq \sigma$ for f linear, then either $\sigma \geq \langle f \rangle$ or σ is ultimately periodic. In [3] it was shown that a similar result holds for quadratic functions, while in [6, 5] it was shown that for higher degree it does not hold. Here we are interested in considering \geq_p and \sim_p instead.


In [2] we already showed that $\langle f \rangle \sim_p \langle g \rangle$ for f, g linear. Here we show that the corresponding class is *not* atomic: we show that for ascending f we have $\langle f \rangle \geq_p \llbracket f \rrbracket$ but not the other way around, and we even show that the class containing $\langle f \rangle$ for linear f is an upper bound of infinitely many distinct classes, in particular

$$\llbracket n \rrbracket <_p \llbracket n+2 \rrbracket <_p \llbracket n+4 \rrbracket <_p \llbracket n+8 \rrbracket <_p \llbracket n+16 \rrbracket <_p \dots \leq_p \langle n \rangle.$$

We write $\sigma <_p \tau$ if $\tau \geq_p \sigma$ but not $\sigma \geq_p \tau$, and use $\llbracket f(n) \rrbracket$ as shorthand for $\llbracket n \mapsto f(n) \rrbracket$, and similarly for $\langle \cdot \rangle$. While all $\langle f \rangle$ for f quadratic are equivalent under ordinary transduction, we show that this does not hold for \sim_p ; in particular, we obtain the infinite ascending chain

$$\langle (n+1)^2 \rangle <_p \langle n^2 \rangle <_p \langle (n-1)^2 \rangle <_p \langle (n-2)^2 \rangle <_p \langle (n-4)^2 \rangle <_p \langle (n-8)^2 \rangle <_p \dots$$

Typically, for proving $\sigma <_p \tau$ the easier part is giving an explicit permutation transducer P satisfying $P(\tau) = \sigma$. The hard part is showing that a permutation transducer for the other way around does not exist. For instance, the easy part of showing $\llbracket n \rrbracket <_p \llbracket n+2 \rrbracket$ can be done using the following permutation transducer, proving $\llbracket n+2 \rrbracket \geq_p \llbracket n \rrbracket$. In presenting a transducer by a picture an arrow labeled by $a|u$ means that an input symbol a is consumed and the string u is produced as output. The initial state is indicated by an incoming arrow not



The initial state is indicated by an incoming arrow not

starting in a state. When consuming $\llbracket n+2 \rrbracket = 0^2 1^3 0^4 1^5 \dots$ by this permutation transducer indeed $\llbracket n \rrbracket = 10^2 1^3 0^4 \dots$ is produced. Here the two 1-arrows between the two top states are never used, and may also produce anything else.

An even simpler ordinary transducer doing the same job, but which is not a permutation transducer, is easily found, as $\llbracket n \rrbracket$ can also be obtained from $\llbracket n+2 \rrbracket$ by simply putting a single symbol 1 in front.

The remaining proof obligation, that no permutation transducer exists transforming $\llbracket n \rrbracket$ to $\llbracket n+2 \rrbracket$, is much harder. For doing this we investigate the pattern of any sequence that can be obtained by applying a permutation transducer to $\llbracket n \rrbracket$, and then prove that $\llbracket n+2 \rrbracket$ does not satisfy this pattern. For all other claims in this paper containing ' $<_p$ ', we give similar arguments all being instances of the following three cases. The first case investigates the creation of isolated 1s, leading to $\llbracket f \rrbracket <_p \langle f \rangle$ for ascending f . The second one investigates transducts of $\langle f \rangle$ and $\llbracket f \rrbracket$ for those f (such as $f(n) = n!$) for which for every m there exists N such that $f(n) \equiv 0 \pmod m$ whenever $n > N$. The third one investigates transducts of $\langle f \rangle$ and $\llbracket f \rrbracket$ for f , such as polynomials, for which $n \mapsto f(n) \pmod m$ is periodic for every m .

We consider four basic ways to transform functions $f : \mathbb{N} \rightarrow \mathbb{N}$: transforming $f(n)$ to $f(n) + k$ and to $f(n + k)$ for any $k \geq 1$, and to $kf(n)$ and to $f(kn)$ for any $k > 1$. For all of them we investigate how $\langle f(n) \rangle$ and $\llbracket f(n) \rrbracket$ relate to their transformed variants, both with respect to ordinary transducers and permutation transducers.

The paper is organized as follows. We start by preliminaries in Section 2. In Section 3 we classify permutation transducts of particular sequences σ , in order to be able to prove $\sigma \not\geq_p \tau$ for certain τ . In Section 4 we investigate the effect of transforming f in the above given four ways. In Section 5 we investigate $\llbracket f \rrbracket$ and $\langle f \rangle$ for linear functions f ; in particular, we give an infinite strictly ascending chain of them. In Section 6 we investigate polynomials of higher degree; in particular, we give an infinite strictly ascending chain of sequences $\langle f \rangle$ for quadratic polynomials f .

2 Preliminaries

In the following we assume $\Sigma = \{0, 1\}$.

Definition 1. A finite state transducer $T = (Q, q_0, \delta, \lambda)$ consists of a finite set Q , $q_0 \in Q$, $\delta : Q \times \Sigma \rightarrow Q$, $\lambda : Q \times \Sigma \rightarrow \Sigma^*$. For $\sigma : \mathbb{N} \rightarrow \Sigma$ we define $T(\sigma) = \lambda(q_0, \sigma(0))\lambda(q_1, \sigma(1))\lambda(q_2, \sigma(2)) \dots$ for q_i defined by $q_{i+1} = \delta(q_i, \sigma(i))$ for $i \geq 0$.

A permutation transducer over Σ is a finite state transducer $T = (Q, q_0, \delta, \lambda)$ with the additional requirement that for every $a \in \Sigma$ the function $q \mapsto \delta(q, a)$ is a bijection from Q to Q .

For $\sigma, \tau : \mathbb{N} \rightarrow \Sigma$ we define \geq_p , \sim_p and $>_p$ by

$$\begin{aligned} \sigma \geq_p \tau &\iff \exists \text{ permutation transducer } T : \tau = T(\sigma), \\ \sigma \sim_p \tau &\iff \sigma \geq_p \tau \wedge \tau \geq_p \sigma, \quad \sigma >_p \tau \iff \sigma \geq_p \tau \wedge \neg(\tau \geq_p \sigma). \end{aligned}$$

In drawing pictures for transducers we write an arrow from p to q labeled by $a|u$ if $\delta(p, a) = q$ and $\lambda(p, a) = u$. We use $\geq, \sim, >$ for the similar relations on sequences based on ordinary finite state transducers, that is, without the additional bijectivity requirement. These were studied extensively in [4, 3, 6, 5]. To see the effect of the additional requirement of permutation transducers, throughout the paper in presenting properties of $\geq_p, \sim_p, >_p$ we often present the corresponding properties of $\geq, \sim, >$.

For a homomorphism $h : \Sigma \rightarrow \Sigma^+$ the transducer $T_h = (\{q_0\}, q_0, \delta, \lambda)$ defined by $\delta(q_0, a) = q_0$ and $\lambda(q_0, a) = h(a)$ for all $a \in \Sigma$ is a permutation transducer satisfying $T_h(\sigma) = h(\sigma)$ for all σ , proving that $\sigma \geq_p h(\sigma)$. In particular, for choosing h to be the identity we obtain that \geq_p is reflexive. A straightforward construction given in [2] shows that \geq_p is transitive. Hence \geq_p is a pre-order, yielding a partial order on equivalence classes with respect to the equivalence relation \sim_p . By defining $h(a) = 0$ for all $a \in \Sigma$ we obtain $T_h(\sigma) = 0^\omega$ for every σ . Hence the equivalence class of 0^ω is the bottom element in this order; it consists of all (purely) periodic sequences as was shown in [2].

A *partial permutation transducer* $T = (Q, q_0, \delta, \lambda)$ consists of a finite set Q and initial state $q_0 \in Q$, together with a partial function $\delta : Q \times \Sigma \rightarrow Q$ such that for every $q \in Q, a \in \Sigma$ there is at most one $q' \in Q$ such that $\delta(q', a) = q$, and $\lambda : Q \times \Sigma \rightarrow \Sigma^*$ is a partial function that is defined on the same pairs that δ is defined for. Thus, in a permutation transducer for every symbol $a \in \Sigma$ there is exactly one incoming and exactly one outgoing a -arrow for every state $q \in Q$, but in a partial permutation transducer 'exactly one' is weakened to 'at most one'. As observed in [2], just like every partial permutation of a set can be extended to a permutation, every partial permutation transducer can be extended to a permutation transducer. Sometimes we will present a permutation transducer by only giving a partial permutation transducer and leaving the extension implicit.

From the introduction recall the definitions

$$\langle f \rangle = \sigma_f = 10^{f(0)}10^{f(1)}10^{f(2)} \dots \quad \text{and} \quad \llbracket f \rrbracket = 0^{f(0)}1^{f(1)}0^{f(2)} \dots$$

for any $f : \mathbb{N} \rightarrow \mathbb{N}$. For the latter it is natural to require $f(n) > 0$ for all $n > 0$, to avoid collapsing groups; we will say that f is *positive* if it satisfies this property. Note that every sequence σ that is not eventually constant has a natural representation $\llbracket f \rrbracket$ for some (positive) function f . The same is true for $\langle f \rangle$ if $\sigma(0) = 1$, with f usually not positive.

Writing $\langle f(n) \rangle$ for $\langle f \rangle$, we obtain $\langle n \rangle = 11010010001 \dots$, and $\langle n+1 \rangle = 101001000 \dots$, so $\langle n+1 \rangle = \text{tail}(\langle n \rangle)$. Using similar shorthand notation, $\llbracket n \rrbracket = 1^1 0^2 1^3 0^4 1^5 \dots$, and $\llbracket n+1 \rrbracket = 0^1 1^2 0^3 1^4 0^5 \dots$, so $T_h(\llbracket n \rrbracket) = \llbracket n+1 \rrbracket$ and $T_h(\llbracket n+1 \rrbracket) = \llbracket n \rrbracket$ for $h(0) = 1, h(1) = 0$, proving $\llbracket n \rrbracket \sim_p \llbracket n+1 \rrbracket$.

We continue with a fruitful lemma.


Lemma 1. *Let P be a permutation transducer over a finite alphabet Σ . Then there exists an integer $N > 0$ such that for every state q and every $u \in \Sigma^+$ it holds that $\delta(q, u^N) = q$.*

Proof. Let n be the number of states. Then $q \mapsto \delta(q, u)$ is a permutation on the n states. Choose N to be the least common multiple of all k with $k \leq n$. Then $q \mapsto \delta(q, u^N) = (q \mapsto \delta(q, u))^N$ is the identity. \square

Proposition 1. *For every positive function $f : \mathbb{N} \rightarrow \mathbb{N}$ holds*

- $\langle f \rangle \sim \llbracket f \rrbracket$, and
- $\langle f \rangle \geq_p \llbracket f \rrbracket$.

Proof. This is proved by the following two transducers.



The left one is a permutation transducer replacing sequences of consecutive 0's that are demarcated by a single 1, alternately by the same number of 0's or 1's; hence transforming $\langle f \rangle$ to $\llbracket f \rrbracket$, showing $\langle f \rangle \geq_p \llbracket f \rrbracket$ and thus also $\langle f \rangle \geq \llbracket f \rrbracket$.

The right one is an ordinary transducer (but not a permutation transducer) transforming $\llbracket f \rrbracket$ to $\langle f \rangle$, showing that $\llbracket f \rrbracket \geq \langle f \rangle$. Together with the just observed $\langle f \rangle \geq \llbracket f \rrbracket$ this proves $\langle f \rangle \sim \llbracket f \rrbracket$. \square

Now we show that $\llbracket f \rrbracket \geq_p \langle f \rangle$ does not generally hold: for certain f no permutation transducer P exists transforming $\llbracket f \rrbracket$ to $\langle f \rangle$. The key idea is that the isolated 1s in $\langle f \rangle$ can not be created when the input only contains big groups of 0s and 1s as in $\llbracket f \rrbracket$. In fact we prove the following stronger result.

Theorem 1. *Let $f, g : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} g(n) = \infty$. Then no permutation transducer P such that $P(\llbracket f \rrbracket) = \langle g \rangle$.*

Proof. Assume such a $P = (Q, q_0, \delta, \lambda)$ exists. Use Lemma 1 to choose p such that $\delta(q, 0^p) = \delta(q, 1^p) = q$ for all states q . Write $u(q, 0) = \lambda(q, 0^p)$ and $u(q, 1) = \lambda(q, 1^p)$ for all states q . Since $\lim_{n \rightarrow \infty} f(n) = \infty$ a number N exists such that $f(n) > 2p$ for all $n \geq N$. Hence beyond a finite initial part, the sequence $\llbracket f \rrbracket$ is composed of strings 0^k and 1^k for $k > 2p$. For each such string the permutation transducer P produces a prefix of $u(q, i)^\omega$ that starts by $u(q, i)^2$ for some state q and $i \in \{0, 1\}$. Beyond a finite initial part, the resulting output $\langle g \rangle$ is the concatenation of such prefixes. Assume that one of the occurring strings $u(q, i)^2$ contains a symbol 1. Then it contains at least two symbols 1 at distance at most m , where m is the maximal size of all $u(q, i)$. Since $\lim_{n \rightarrow \infty} g(n) = \infty$, the total number of occurrences of $u(q, i)^2$ in $\langle g \rangle$ that contain a symbol 1, is finite. This contradicts the fact that $\langle g \rangle$ contains infinitely many 1s. \square

For proving more claims of the type $\sigma \not\geq_p \tau$ we typically investigate the shape of sequences $P(\sigma)$, the *transducts* of σ : then it remains to show that τ is not of the required shape. In the next section we give a number of results of this type.

3 Classifying Transducts

We want to classify permutation transducts of sequences of the shape $\langle f \rangle$ and $\llbracket f \rrbracket$ for well-known functions f , like polynomials. A key property of polynomials f that will be exploited is that the function $n \mapsto (f(n) \bmod m)$ is periodic for every $m > 0$. We start by a class of functions for which the analysis is slightly simpler, namely functions like $f(n) = n!$ for which $n \mapsto (f(n) \bmod m)$ is ultimately 0 for every $m > 0$.

Theorem 2. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a positive function for which for every $m > 0$ there exists $N \in \mathbb{N}$ such that $f(n) \equiv 0 \pmod m$ for all $n > N$. If $\llbracket f \rrbracket \geq_p \sigma$ then there exist $u, c, d \in \Sigma^*$ and $b, h \in \mathbb{N}$ such that*

$$\sigma = u \prod_{i=0}^{\infty} \left(c^{f(b+2i)/h} d^{f(b+2i+1)/h} \right) = uc^{f(b)/h} d^{f(b+1)/h} c^{f(b+2)/h} d^{f(b+3)/h} \dots$$

Proof. Let $P = (Q, q_0, \delta, \lambda)$ be a permutation transducer such that $P(\llbracket f \rrbracket) = \sigma$. By Lemma 1 there exists h such that $\delta(q, 0^h) = \delta(q, 1^h) = q$ for all q . Choose b even such that $f(i) \equiv 0 \pmod h$ for all $i \geq b$. Let u be the output of P of the initial part $v = 0^{f(0)} 1^{f(1)} \dots 1^{f(b-1)}$, and let $q = \delta(q_0, v)$. Let $c = \lambda(q, 0^h)$ and $d = \lambda(q, 1^h)$. Then the next blocks $0^{f(b)}$, $1^{f(b+1)}$, $0^{f(b+2)}$, $1^{f(b+3)}$, \dots produce output $c^{f(b)/h} d^{f(b+1)/h} c^{f(b+2)/h} d^{f(b+3)/h} \dots$, exactly the pattern claimed. \square

Corollary 1. $\llbracket n! \rrbracket \not\geq_p \llbracket n! - 1 \rrbracket$.

Proof. Suppose that $\llbracket n! \rrbracket \geq_p \llbracket n! - 1 \rrbracket$. Then by Theorem 2 we obtain

$$\llbracket n! - 1 \rrbracket = uc^{b!/h} d^{(b+1)!/h} c^{(b+2)!/h} d^{(b+3)!/h} \dots$$

Since in $\llbracket n! - 1 \rrbracket$ only groups of 0s and 1s occur of increasing size, both c and d either consist only of 0s or only of 1s. Since $\llbracket n! - 1 \rrbracket$ contains infinitely many 0s and infinitely many 1s, either c consists of 0s and d consists of 1s, or the other way around. But then the resulting consecutive groups of 0s and 1s have sizes $|c|b!/h$, $|d|(b+1)!/h$, $|c|(b+2)!/h$, $|d|(b+3)!/h \dots$, ultimately divisible by any number, which does not hold for the group sizes $n! - 1$, $(n+1)! - 1$, $(n+2)! - 1$, $(n+3)! - 1 \dots$ in $\llbracket n! - 1 \rrbracket$. This contradiction proves $\llbracket n! \rrbracket \not\geq_p \llbracket n! - 1 \rrbracket$. \square

Corollary 2. $\llbracket n! \rrbracket \not\geq_p \llbracket (2n)! \rrbracket$.

Proof. As in the previous proof, use the form of the transducts of $\llbracket n! \rrbracket$ given by Theorem 2: again c and d both consist of copies of a single symbol, different for the two. But now it will be impossible for such transduct to equal $\llbracket (2n)! \rrbracket$ because the growth of the groups in the transduct is like a multiple of $n!$, which is much slower than that of the groups in $\llbracket (2n)! \rrbracket$. \square

For the same class of functions we now give a characterization for transducts of $\langle f \rangle$ rather than $\llbracket f \rrbracket$.

Theorem 3. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function for which for every $m > 0$ there exists $N \in \mathbb{N}$ such that $f(n) \equiv 0 \pmod{m}$ for all $n > N$. If $\langle f \rangle \geq_p \sigma$ then there exist $k > 0$, $a \geq 0$ and $u, p_0, \dots, p_{k-1}, c_0, \dots, c_{k-1} \in \Sigma^*$ such that

$$\sigma = u \prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i c_i^{f(a+i+jk)/k} \right) = up_0 c_0^{f(a)/k} p_1 c_1^{f(a+1)/k} \dots$$

Proof. Assume that $P(\langle f \rangle) = \sigma$ for a permutation transducer $P = (Q, q_0, \delta, \lambda)$. Choose k by Lemma 1 such that $\delta(q, 0^k) = \delta(q, 1^k) = q$ for all $q \in Q$. By the assumption on f there exists a such that $f(n) \equiv 0 \pmod{k}$ for all $n \geq a$. Let $v = 10^{f(0)} 10^{f(1)} 1 \dots 10^{f(a-1)}$, which is a prefix of $\langle f \rangle$. Let $u = \lambda(q_0, v)$, and $r_0 = \delta(q_0, v)$. Define $r_i = \delta(r_0, 1^i)$ for $i = 1, 2, \dots, k$; since $\delta(r_0, 1^k) = r_0$ we have $r_k = r_0$. Since $f(a+i) \equiv 0 \pmod{k}$ we obtain $r_{i+1} = \delta(r_i, 10^{f(a+i)})$ for $i = 0, \dots, k-1$. Write $p_i = \lambda(r_i, 1)$ and $c_i = \lambda(r_{i+1}, 0^k)$, then by using $\delta(r_{i+1}, 0^k) = r_{i+1}$ for $i = 0, \dots, k-1$, we obtain the desired result that σ equals

$$u \lambda(r_0, 10^{f(a)}) \lambda(r_1, 10^{f(a+1)}) \lambda(r_2, 10^{f(a+2)}) \dots = u \prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i c_i^{f(a+i+jk)/k} \right).$$

□

Corollary 3. $\langle n! \rangle \not\geq_p \langle n! - 1 \rangle$.

Proof. Suppose that $\langle n! \rangle \geq_p \langle n! - 1 \rangle$. Then by Theorem 3 we obtain

$$\langle n! - 1 \rangle = up_0 c_0^{a!/k} p_1 c_1^{(a+1)!/k} p_2 c_2^{(a+2)!/k} \dots$$

Since in $\langle n! - 1 \rangle$ only increasing groups of 0s occur between consecutive 1s, every p_i contains at most one 1 and every c_i only consists of 0s. By possibly doubling k , we may assume that two distinct p_i s contain a 1; let p_g and p_h be the first two containing a 1. For $i = 0, \dots, h-g-1$ define d_i by $c_i = 0^{d_i}$. Then for every $j \geq 0$ the string $p_g c_g^{(a+g+jk)!/k} p_{g+1} c_{g+1}^{(a+g+1+jk)!/k} \dots p_h$ is a part of $\langle n! - 1 \rangle$ containing exactly two 1s, with exactly $C + \sum_{i=0}^{h-g-1} \frac{(a+g+jk)! d_i}{k}$ separating 0s, for some constant $C \geq 0$. Choose $N > 2C + 2$. Then for j large enough all of these groups of 0s have size $C \pmod{N}$, contradicting the fact that after a finite part $\langle n! - 1 \rangle$ only contains groups of 0s of size $-1 \pmod{N}$. □

Next we switch to functions f , like polynomials, for which $n \mapsto (f(n) \pmod{m})$ is periodic for every $m > 0$. For transducts of $\langle f \rangle$ under permutation transducers the following characterization was given in [2].

Theorem 4. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function for which $n \mapsto (f(n) \pmod{m})$ is periodic for every $m > 0$. Then $\langle f \rangle \geq_p \sigma$ for $\sigma : \mathbb{N} \rightarrow \Sigma$ if and only if there exist $k, h > 0$ and $p_0, \dots, p_{k-1}, c_0, \dots, c_{k-1} \in \Sigma^*$ such that

$$\sigma = \prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i c_i^{\lfloor f(i+jk)/h \rfloor} \right) = p_0 c_0^{\lfloor f(0)/h \rfloor} p_1 c_1^{\lfloor f(1)/h \rfloor} \dots$$

We give a similar description of transducts of $\llbracket f \rrbracket$.

Theorem 5. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function for which $n \mapsto (f(n) \bmod m)$ is periodic for every $m > 0$. If $\llbracket f \rrbracket \geq_p \sigma$ for $\sigma : \mathbb{N} \rightarrow \Sigma$ then there exist $k, h > 0$ and $p_0, \dots, p_{k-1}, c_0, \dots, c_{k-1} \in \Sigma^*$ such that*

$$\sigma = \prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i(c_i p_i)^{\lfloor f(i+jk)/h \rfloor} \right) = p_0(c_0 p_0)^{\lfloor f(0)/h \rfloor} p_1(c_1 p_1)^{\lfloor f(1)/h \rfloor} \dots,$$

with the additional constraint that $p_i = \epsilon$ if $f(i) \equiv 0 \pmod h$, and c_i is non-empty.

Proof. Let f be as in the statement, and suppose that $\llbracket f \rrbracket \geq_p \sigma$, for some $\sigma \in \Sigma^\omega$, given by a permutation transducer $P = (Q, q_0, \delta, \lambda)$. Let $m = \#Q$, and let h be the positive integer N from Lemma 1, so $\delta^h(q, x) = q$ for every $q \in Q, x \in \Sigma^+$. We will make use of the assumption that $\llbracket f \bmod h \rrbracket$ is periodic, as $n \mapsto (f(n) \bmod h)$ is; replacing u by u^h if necessary, we may assume that $\llbracket f \bmod h \rrbracket = \lambda(u)^\omega$. By k we will denote the (minimal) period of $f \bmod h$; however, if k is odd, we replace it by $2k$, doubling h if necessary.

Starting at state q_0 , the transducer P on input $\llbracket f \bmod h \rrbracket$ will read $f(0) \bmod h$ 0's and arrive at a state we will call q_1 . Let h_0 be the least positive integer for which $\delta(q_0, 0^{h_0}) = q_0$; note that h_0 is a divisor of h by construction. If we write $f(0) = x_0 + y_0 \cdot h$, with $0 \leq x_0 < h$, then by assumption $f(j \cdot k) = x_0 + z_j \cdot h$ for every $j \geq 1$. Define $p_0 = \lambda(q_0, 0_0^x)$ and $c_0 p_0 = \lambda(q_1, 0^h)$, where the state $q_1 = \delta(q_0, 0_0^x)$; this can be done since h is a multiple of h_0 and hence $\delta(q_1, 0^h) = q_1$ and p_0 is a postfix of $\lambda(q_1, 0^h)$. Note that $q_1 = q_0$ if and only if x_0 is a multiple of h_0 ; this holds in particular when $x_0 = 0$, in which case $p_0 = \epsilon$.

We proceed by similarly defining p_1 and c_1 : with $f(1) = x_1 + y_1 \cdot h$ we obtain $q_2 = \delta(q_1, 0_1^x)$ and put $p_1 = \lambda(q_1, 0_1^x)$ and $c_1 p_1 = \lambda(q_2, 0^h)$. This is repeated, to obtain x_i, q_{i+1}, p_i, c_i for $i = 0, 1, \dots, k-1$; by our choices, $q_k = \delta(q_{k-1}, 1_{k-1}^x) = q_0$.

By periodicity of $f \bmod h$ with period k , it is now easily seen that $\sigma = P(\llbracket f \rrbracket)$ is of the desired form. \square

4 Basic Function Operations

In this section we investigate how $\langle f(n) \rangle$ relates to $\langle f(n+k) \rangle$, $\langle f(n) + k \rangle$, $\langle kf(n) \rangle$ and $\langle f(kn) \rangle$, and similarly for $\llbracket \cdot \rrbracket$. For completeness we do not only consider \sim_p, \geq_p, \leq_p based on permutation transducers, but also \sim, \geq, \leq based on ordinary transducers.

Theorem 6. *When linear operations on sequences of the form $\langle f \rangle$ or $\llbracket f \rrbracket$ are performed, the general relation between the original sequence and its image by ordinary or permutation is given by an entry in the following two tables:*

$\langle f(n) \rangle \leq, \geq, \not\leq_p, \geq_p \langle f(n+k) \rangle$
$\langle f(n) \rangle \leq, \geq, \not\leq_p, ? \langle f(n+k) \rangle$
$\langle f(n) \rangle \leq, \geq, \leq_p, \geq_p \langle kf(n) \rangle$
$\langle f(n) \rangle \not\leq, \geq, \not\leq_p, \geq_p \langle f(kn) \rangle$

$\llbracket f(n) \rrbracket \leq, \geq, \not\leq_p, \not\geq_p \llbracket f(n+k) \rrbracket$
$\llbracket f(n) \rrbracket \leq, \geq, \not\leq_p, \not\geq_p \llbracket f(n+k) \rrbracket$
$\llbracket f(n) \rrbracket \leq, \geq, \leq_p, \geq_p \llbracket kf(n) \rrbracket$
$\llbracket f(n) \rrbracket \not\leq, \geq, \not\leq_p, \not\geq_p \llbracket f(kn) \rrbracket$

An entry of the form $\sigma \geq \tau$ indicates that for every f, k a transducer exists transforming σ to τ , while $\sigma \not\geq \tau$ indicates that f, k exist for which such a transducer does not exist, and similar for \leq, \leq_p, \geq_p . The question mark '?' states that this question is open for $\langle f(n) \rangle \geq_p \langle f(n+k) \rangle$.

The correctness of these statements is given by the proofs of the following four propositions.

Proposition 2. For every positive function $f : \mathbb{N} \rightarrow \mathbb{N}$:

- $\langle f(n) \rangle \sim \langle f(n+1) \rangle$;
- $\langle f(n) \rangle \geq_p \langle f(n+1) \rangle$ but $\langle f(n) \rangle \leq_p \langle f(n+1) \rangle$ does not hold generally;
- $\llbracket f(n) \rrbracket \sim \llbracket f(n+1) \rrbracket$;
- neither $\llbracket f(n) \rrbracket \geq_p \llbracket f(n+1) \rrbracket$ nor $\llbracket f(n) \rrbracket \leq_p \llbracket f(n+1) \rrbracket$ holds generally.

Proof. For any f the permutation transducer on the left



implies that $\langle f(n) \rangle \geq_p \langle f(n+1) \rangle$, and hence \geq too. Conversely, the (partial) ordinary transducer on the right implies $\langle f(n+1) \rangle \geq \langle f(n) \rangle$ for every positive f . Thus $\langle f \rangle \sim \langle f+1 \rangle$ and since both $\llbracket f \rrbracket \sim \langle f \rangle$ and $\llbracket f+1 \rrbracket \sim \langle f+1 \rangle$ by Proposition 1, we immediately get $\llbracket f \rrbracket \sim \llbracket f+1 \rrbracket$.


In Corollary 3 we have seen that $\langle f(n+1) \rangle \geq_p \langle f(n) \rangle$ does not hold for $f(n) = n! - 1$.

Finally, neither $\llbracket f \rrbracket \geq_p \llbracket f+1 \rrbracket$ nor $\llbracket f \rrbracket \leq_p \llbracket f+1 \rrbracket$ hold in general, as we will see later in Section 5 in Corollary 4 (for $f(n) = n+1$) and we have seen in Corollary 1 (for $f(n) = n! - 1$).

Proposition 3. For every positive function $f : \mathbb{N} \rightarrow \mathbb{N}$:

- $\langle f(n) \rangle \sim \langle f(n+1) \rangle$;
- $\langle f(n) \rangle \leq_p \langle f(n+1) \rangle$ does not hold generally;
- $\llbracket f(n) \rrbracket \sim \llbracket f(n+1) \rrbracket$; does not hold generally;
- neither $\llbracket f(n) \rrbracket \geq_p \llbracket f(n+1) \rrbracket$ nor $\llbracket f(n) \rrbracket \leq_p \llbracket f(n+1) \rrbracket$ holds generally.

Proof. The ordinary transducers



prove the first claim.

As an example for which no permutation transducer maps $\langle f(n+1) \rangle$ to $\langle f(n) \rangle$, take $f(0) = 2$ and $f(n) = 1$ for $n \geq 1$; then $\langle f(n+1) \rangle = (10)^\omega$ is periodic, but $\langle f(n) \rangle = 10^2(10)^\omega$ is not; hence no permutation transducer will transform $\langle f(n+1) \rangle$ to $\langle f(n) \rangle$. See also Corollary 6.

For the opposite case for permutation transducers, it is conjectured that $\langle f(n) \rangle \geq_p \langle f(n+1) \rangle$ does not generally hold, but the proof is missing as yet.

These transducers



map $\llbracket f(n) \rrbracket$ to $\llbracket f(n+1) \rrbracket$ and vice versa, for positive functions f .


The claim that no permutation transducer will map $\llbracket f(n) \rrbracket$ to $\llbracket f(n+1) \rrbracket$ in general, is proven in Corollary 4 (for $f(n) = n+1$).

In the opposite direction, there will be no permutation transducer in general to map $\llbracket f(n+1) \rrbracket$ to $\llbracket f(n) \rrbracket$; explicitly, again the function f defined by $f(0) = 2$ and $f(n) = 1$ for $n \geq 1$ we see that $\llbracket f(n+1) \rrbracket = (10)^\omega$ is periodic but $\llbracket f(n) \rrbracket = 0^2(10)^\omega$ is not, hence not $\llbracket f(n+1) \rrbracket \geq_p \llbracket f(n) \rrbracket$.

Proposition 4. For every positive function $f : \mathbb{N} \rightarrow \mathbb{N}$ and all $k \geq 1$:


- $\langle f(n) \rangle \sim \langle k \cdot f(n) \rangle$.
- $\langle f(n) \rangle \sim_p \langle k \cdot f(n) \rangle$.
- $\llbracket f(n) \rrbracket \sim \llbracket k \cdot f(n) \rrbracket$.
- $\llbracket f(n) \rrbracket \sim_p \llbracket k \cdot f(n) \rrbracket$.

Proof. The permutation transducer on the left in the following picture shows that $\langle f(n) \rangle \geq_p \langle k \cdot f(n) \rangle$ (and hence $\langle f(n) \rangle \geq \langle k \cdot f(n) \rangle$) for every positive f and every $k \geq 1$; notice that for $k = 1$ this is the identity transducer.



For the converse, we describe a partial permutation transducer (the picture on the right shows the case $k = 4$): it consists of k states, connected by a directed cycle of k arrows, all with rule $0 | \epsilon$ except for (say) the first one, which has $0 | 0$. Moreover, on the initial state there is a directed loop with rule $1 | 1$.

The one-state permutation transducer on the left establishes one direction of the fourth (and hence third) claim.




For the direction $\llbracket k \cdot f(n) \rrbracket \geq_p \llbracket f(n) \rrbracket$ (implying \geq as well) we adapt the k -cycle described above: this time to the initial node on the main cycle we attach a new directed k -cycle, each arrow carrying the rule $1 \mid \epsilon$ with a single exception again, on the first arrow (say), which has $1 \mid 1$ instead. This way k consecutive like symbols are replaced by just one of them. The picture above shows on the right the case $k = 4$.

Proposition 5. *For every positive function $f : \mathbb{N} \rightarrow \mathbb{N}$ and all $k \geq 1$:*

- $\langle f(n) \rangle \geq \langle f(k \cdot n) \rangle$ but not conversely, generally;
- $\langle f(n) \rangle \geq_p \langle f(k \cdot n) \rangle$ but not conversely, generally;
- $\llbracket f(n) \rrbracket \geq \llbracket f(k \cdot n) \rrbracket$, but not conversely, generally;
- neither $\llbracket f(n) \rrbracket \geq_p \llbracket f(k \cdot n) \rrbracket$ nor $\llbracket f(n) \rrbracket \leq_p \llbracket f(k \cdot n) \rrbracket$ holds generally.

Proof. Again, we combine the cases $\langle f(n) \rangle \geq \langle f(k \cdot n) \rangle$ and $\langle f(n) \rangle \geq_p \langle f(k \cdot n) \rangle$ by describing a (partial) permutation transducer, mapping $\langle f(n) \rangle$ to $\langle f(k \cdot n) \rangle$: it will have k nodes connected by a directed cycle of arrows with rule $1 \mid \epsilon$ on them, except for the first arrow, which has $1 \mid 1$. Moreover, in each node there will be a directed loop with rule $0 \mid \epsilon$, except for the node following the initial node on the cycle, where the rule on the loop will be $0 \mid 0$. We depict the case $k = 4$.




That $\langle f(k \cdot n) \rangle \geq_p \langle f(n) \rangle$ does not hold in general can be seen from taking as an example the case $k = 2$ and $f(n) = 1$ if $n \neq 1$ and $f(1) = 2$: then $\langle f(2n) \rangle = (10)^\omega$ is periodic, but $\langle f(n) \rangle = 1010^2(10)^\omega$ is not.

Here we describe the ordinary transducer that shows $\llbracket f(n) \rrbracket \geq \llbracket f(k \cdot n) \rrbracket$. We distinguish the cases k even and k odd.

If k is odd, the transducer will consist of a directed $2k$ -cycle with directed loops attached to all $2k$ states; the arrows on the cycle will read alternately a symbol 0 or a symbol 1. The arrow to the initial state has rule $0 \mid 0$, and all other arrows reading 0 have rule $0 \mid \epsilon$ attached. The arrow diagonally opposite

to this arrow on the cycle carries rule $1 \mid 1$ and all other arrows reading 1 have rule $1 \mid \epsilon$. The loops read alternately symbols 0 and 1, starting with 0 at the initial node, and all output ϵ , with exception of the loop on the initial node, where $0 \mid 0$ is used and the loop diagonally opposite, where where $1 \mid 1$ applies. We show the resulting transducer for $k = 3$:



If k is even there will be a main cycle of size k ; but note that for a positive function f the result $\llbracket f(n) \rrbracket$ will be periodic, namely 0^ω or 1^ω .

The result upon input $f(n)$ will be that only the strings $b^{f(kn)}$ will be copied, for $n \geq 0$, all others ignored.

By Corollary 2 we have that $\llbracket n! \rrbracket \not\geq_p \llbracket (2n)! \rrbracket$, so not always $\llbracket f(n) \rrbracket \geq_p \llbracket f(2n) \rrbracket$.

Finally, in general not $\llbracket f(kn) \rrbracket \geq \llbracket f(n) \rrbracket$ (and thus certainly not \geq_p); explicitly now, the function g defined by $g(1) = 2$ and $g(n) = 1$ for $n \neq 1$ is not periodic while $g(2n)$ is.

5 Classes of Linear Functions

For linear polynomial functions $f_{k,l}(n) = kn + l$, the relations between $\langle f_{k,l} \rangle$ were already dealt with in [2].

Theorem 7. *For all $k, l \in \mathbb{N}$ with $k \geq 1$: $\langle kn + l \rangle \sim_p \langle n \rangle$; in other words: all (non-constant) linear functions are equivalent under \sim_p .*

The situation is markedly different for $\llbracket f_{k,l} \rrbracket$, as we will see in the next theorem. Note that by Proposition 1 and Theorem 1 we already know $\llbracket n \rrbracket <_p \langle n \rangle$, so $\langle n \rangle$ is not atomic. We do not yet know whether $\llbracket n \rrbracket$ is atomic or not.

Theorem 8. *Under permutation transduction there is an infinite, strictly ascending sequence of equivalence classes containing $\llbracket f \rrbracket$ for linear polynomials f , in between $\llbracket n \rrbracket$ and $\langle n \rangle$; in particular*


$$\llbracket n \rrbracket <_p \llbracket n + 2 \rrbracket <_p \llbracket n + 4 \rrbracket <_p \llbracket n + 8 \rrbracket <_p \dots \leq_p \langle n \rangle.$$

Before we give the proof, we state a corollary that settles a question from the previous section.

Corollary 4. *No permutation transducer P exists such that $P(\llbracket n+1 \rrbracket) = \llbracket n+2 \rrbracket$.*

Proof. This is the content of the first strict inequality in Theorem 8, in combination with $\llbracket n \rrbracket \sim_p \llbracket n+1 \rrbracket$. The latter follows from $\overline{\llbracket n+1 \rrbracket} = \llbracket n \rrbracket$, in which $\bar{\sigma}$ denotes the complement of σ , obtained by permuting the two symbols 0, 1; it will be clear that $\bar{\sigma}$ can be obtained from σ by a one-state permutation transducer. \square

The proof of Theorem 8 is given in two parts. The first (existence) part is immediate from the following lemma.



Lemma 2. *Let $k \geq 0$ be an integer; then $\llbracket n+k \rrbracket \geq_p \llbracket n + \lfloor \frac{k+1}{2} \rfloor \rrbracket$.*

Proof. For $k = 0, 1$ the statement is trivial; so assume that $k \geq 2$. Consider the four-state permutation transducer given above. For even k it will remove any 1's from the input sequence $\llbracket n+k \rrbracket$ and alternately divide by 2 or divide by 2 and complement, any sequence of 0's; the result is $\overline{\llbracket n + \frac{k}{2} \rrbracket}$. For odd k it converts $\overline{\llbracket n+k \rrbracket}$ into $\llbracket n + \frac{k+1}{2} \rrbracket$. Taking complements is easily achieved by a permutation transducer, and the permutation property is transitive. \square

In fact this lemma yields a stronger result than required for Theorem 8, as stated in the following corollary.

Corollary 5. *For every $a > 0$ we have $\llbracket n+a \rrbracket \geq_p \llbracket n \rrbracket$.*

Proof. Starting by $\llbracket n+a \rrbracket$ repeat applying Lemma 2 until $\llbracket n+1 \rrbracket$ is obtained. Then the corollary follows from transitivity of \geq_p and $\llbracket n+1 \rrbracket \sim_p \llbracket n \rrbracket$. \square

To complete the proof of Theorem 8 we have to prove that for none of the strict steps a backward transduction is possible. This is immediate from the following stronger result.

Proposition 6. *For $a \geq 0$, $b \geq a+2$ no permutation transducer P exists satisfying $P(\llbracket n+a \rrbracket) = \llbracket n+b \rrbracket$.*

Proof. The permutation transducts of $\llbracket n+a \rrbracket$, according to Theorem 5, are of the form

$$\prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i (c_i p_i)^{\lfloor \frac{i+jk+a}{h} \rfloor} \right).$$

Replace the period k by a multiple (if necessary) in order for h to be a divisor of k , and write $m = \frac{k}{h}$ and $a_i = \lfloor \frac{i+a}{k} \rfloor$ for $i = 0, \dots, k-1$. Write $b_i = \lfloor \frac{i+a}{h} \rfloor - a_i m$ for $i = 0, \dots, k-1$, note that $0 \leq b_i < m$. Then the transduct is of the shape $\prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i (c_i p_i)^{b_i + (j+a_i)m} \right)$. Writing $w_i = (c_i p_i)^m$ and replacing

p_i by $p_i(c_i p_i)^{b_i}$, we conclude that the transduct is of the shape $P(\llbracket n + a \rrbracket) = \prod_j \prod_{i=0}^{k-1} p_i(w_i)^{j+a_i}$.

Now suppose that this image equals $\llbracket n + b \rrbracket$, for some $b \geq a + 2$. It is impossible for w_i to contain both 0 and 1, since in that case $\prod_j \prod_i p_i w_i^{j+a_i}$ contains infinitely many pairs of the same symbol separated by a fixed number of copies of the other symbol, which $\llbracket n + b \rrbracket = 0^{b+1} 1^{b+2} 0^{b+3} \dots$ clearly does not.

Hence each w_i consist of copies of a single symbol; if w_{i+1} consists of the same symbol or equals ϵ , we can merge $p_i w_i p_{i+1} w_{i+1}$ and reduce k . Hence without loss of generality we may assume that w_i and w_{i+1} consist of different symbols; the same then holds for p_i and p_{i+1} (but they could equal ϵ). By multiplying the period k we may assume that k is even and $k > b$.

The linear growth of $\llbracket n + b \rrbracket$ implies that each w_i will consist of exactly k symbols; since $p_i w_i$ and $p_{i+1} w_{i+1}$ are consecutive blocks of different symbols, $\#p_{i+1} \bmod k = (\#p_i + 1) \bmod k$.

Since $k > b > a$ and $a_i = \lfloor \frac{i+a}{k} \rfloor$ we obtain $a_i = 0$ for $i < k - a$. So $p_0 p_1 \dots p_{k-a-1}$ is an initial part of $\llbracket n + b \rrbracket$, in which p_i alternately consist of 0s and 1s. If $p_0 = \epsilon$ then p_1 is the first group of 0s, being 0^b , contradicting $\#p_{i+1} \bmod k = (\#p_i + 1) \bmod k$. Hence $p_0 = 0^b$, and by $\#p_{i+1} \bmod k = (\#p_i + 1) \bmod k$ we obtain $\#p_i = b + i$ for $i = 0, 1, \dots, k - a - 1$. But since $\#(c_i p_i) = k$ we obtain $\#p_i \leq k$. But then we have $b + k - a - 1 = \#p_{k-a-1} \leq k$, contradicting $b \geq a + 2$. \square

6 Higher Degree Polynomials

Theorem 9. *Let $f, g : \mathbb{N} \rightarrow \mathbb{N}$ be two polynomials of degree $n > 1$ with the same leading coefficient such that*

- $f - g$ is not constant, and
- $\lim_{x \rightarrow \infty} (f(ax) - a^n g(x)) = \infty$ for every $a > 1$.

Then no permutation transducer P exists such that $P(\langle f \rangle) = \langle g \rangle$.

Proof. Assume that such a P exists. Then according to Theorem 4 there exist $k, h > 0$ and $p_0, \dots, p_{k-1}, c_0, \dots, c_{k-1} \in \Sigma^*$ such that

$$\langle g \rangle = \prod_{j=0}^{\infty} \left(\prod_{i=0}^{k-1} p_i c_i^{\lfloor \frac{f(i+jk)}{h} \rfloor} \right).$$

Since $\lim_{n \rightarrow \infty} g(n) = \infty$, for every $i = 0, \dots, k-1$ no 1 occurs in c_i , and at most one 1 occurs in p_i . Since $\langle g \rangle$ contains symbols 1, at least one of the p_i 's contains a symbol 1.

If there is only one such p_i , by doubling k we make it two.

Let p_a and p_b be the first two p_i 's containing a 1. Let q be the total number of 1's in p_0, \dots, p_{k-1} , and $c_i = 0^{a_i}$ for $i = 0, \dots, k-1$. Now we count the number of 0's right after the $qj + 1$ -th 1 of $\langle f \rangle$ in two ways, and obtain that there is

constant $c \geq 0$ (corresponding to the number of 0's occurring in some p_i 's) such that

$$c + \sum_{i=a}^{b-1} \lfloor \frac{f(i+jk)}{h} \rfloor a_i = g(jq)$$

for all $j \geq 0$.

First we consider the case $k = q$. Then $a = 0$, $b = 1$ and we have $c + a_0 \lfloor \frac{f(jk)}{h} \rfloor = g(jk)$ for all $j \geq 0$. This is only possible if $f - g$ is constant, which we assumed to be not. In the remaining case we have $0 < q < k$.

Write $A = \sum_{i=a}^{b-1} a_i$.

Using that f is ascending for sufficiently large arguments, we have $f(jk) \leq f(i+jk) \leq f((j+1)k)$ for $j > C$ for some C , and $a \leq i < b$.

Using this and $x - 1 \leq \lfloor x \rfloor \leq x$ for all x , we obtain

$$c + A \left(\frac{f(jk)}{h} - 1 \right) \leq g(jq) \leq c + A \frac{f((j+1)k)}{h}$$

for all $j \geq C$. Then for $j \rightarrow \infty$ in the above inequalities we obtain $Ak^n = hq^n$. Then the left inequality yields

$$c - A + \left(\frac{q}{k} \right)^n f(jk) = c + A \left(\frac{f(jk)}{h} - 1 \right) \leq g(jq)$$

for all $j \geq C$. This contradicts $\lim_{x \rightarrow \infty} (f(ax) - a^n g(x)) = \infty$ for $a = \frac{k}{q} > 1$. \square

Corollary 6. $\langle (n+1)^2 \rangle \not\leq_p \langle n^2 \rangle$ and $\langle n^2 \rangle \not\leq_p \langle (n-1)^2 \rangle$.


Lemma 3. For $k > 0$ there is no permutation transducer P such that $P : \langle (n-k)^2 \rangle \mapsto \langle (n-2k)^2 \rangle$.

Proof. Apply Theorem 9 directly to $f = (n-k)^2$ and $g = (n-2k)^2$. \square

Corollary 7. The following provides an infinite ascending chain of quadratic polynomial functions that are non-equivalent under permutation transducers:

$$\langle (n+1)^2 \rangle <_p \langle n^2 \rangle <_p \langle (n-1)^2 \rangle <_p \langle (n-2)^2 \rangle <_p \langle (n-4)^2 \rangle <_p \langle (n-8)^2 \rangle <_p \dots$$

Proof. Consider the three permutation transducers



based on the principle that the left cycle reduces $10^{4m}1$ to 10^m and the right cycle $10^{4m+1}1$ to 1. Here $0^4|u$ means that all four arrows consume 0, while only one has output u , the others have empty output. It is not hard to see that the first transduces $\langle (n-2k)^2 \rangle$ to $\langle (n-k)^2 \rangle$ for every $k > 0$, the second transduces $\langle (n-1)^2 \rangle \mapsto \langle n^2 \rangle$ and the third transduces $\langle n^2 \rangle$ to $\langle (n+1)^2 \rangle$. None of the arrows is reversible by Corollary 6 and Lemma 3. \square

Remark 1. Now consider the permutation transducers:



The one on the left (or the first of the three transducers in the previous picture) is easily seen to provide transition from $\langle(n + 2k)^2\rangle$ to $\langle(n + k)^2\rangle$, for $k > 0$, so

$$\langle(n + 1)^2\rangle \leq_p \langle(n + 2)^2\rangle \leq_p \langle(n + 4)^2\rangle \leq_p \langle(n + 8)^2\rangle \leq_p \dots;$$

but here transductions in the opposite direction are not ruled out by Theorem 9. The other transducer shows that $\langle(n + 2k - 1)^2\rangle \geq_p \langle(n + k)^2\rangle$ for $k > 0$, and puts $\langle(n + 2k - 1)^2\rangle$ in some infinite non-descending sequence. For example:

$$\langle(n + 1)^2\rangle \leq_p \langle(n + 2)^2\rangle \leq_p \langle(n + 3)^2\rangle \leq_p \langle(n + 5)^2\rangle \leq_p \langle(n + 9)^2\rangle \dots$$

References

1. J.-P. Allouche and J. Shallit. *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press, 2003.
2. W. Bosma and H. Zantema. Ordering sequences by permutation transducers. *Indagationes Mathematicae*, 38:38–54, 2017.
3. J. Endrullis, C. Grabmayer, D. Hendriks, and H. Zantema. The degree of squares is an atom. In F. Manea and D. Nowotka, editors, *Combinatorics on Words: 10th International Conference, WORDS 2015, Kiel, Germany, September 14-17*, volume 9304 of *LNCS*, pages 109–121. Springer International Publishing, 2015.
4. J. Endrullis, D. Hendriks, and J. W. Klop. Degrees of streams. *Integers, Electronic Journal of Combinatorial Number Theory*, 11B(A6):1–40, 2011. Proc. Leiden Numeration Conf. 2010.
5. J. Endrullis, J. Karhumäki, J.W. Klop, and A. Saarela. Degrees of infinite words, polynomials and atoms. In *Developments in Language Theory - 20th International Conference, DLT 2016, Montréal, Canada, July 25-28, 2016, Proceedings*, volume 9840, pages 164–176. Springer International Publishing, 2016.
6. J. Endrullis, J.W. Klop, A. Saarela, and M. Whiteland. Degree of transducibility. In F. Manea and D. Nowotka, editors, *Combinatorics on Words: 10th International Conference, WORDS 2015, Kiel, Germany, September 14-17*, volume 9304 of *LNCS*, pages 1–13. Springer International Publishing, 2015.
7. J.E. Pin. On reversible automata. In I. Simon, editor, *LATIN'92*, volume 583 of *LNCS*, pages 401–416. Springer, 1992.
8. J. Sakarovitch. *Elements of automata theory*. Cambridge University Press, 2009.