

# Integral trees homeomorphic to a double star

A. E. Brouwer

2009-09-17

## Abstract

Trees with two nonadjacent vertices of degree larger than two are not integral. This settles a question by Watanabe & Schwenk (1979).

## 1 Introduction

The spectrum of a graph is the spectrum of its (0,1)-adjacency matrix. A graph is called *integral* when its spectrum is integral, i.e., when all eigenvalues of its adjacency matrix are integers. A tree is a connected graph without circuits. In [4] Watanabe & Schwenk studied integral trees, and characterized those integral trees that have a single vertex of degree larger than two, or two adjacent vertices of degree larger than two.

They write

We would now like to examine the trees homeomorphic to a double star, that is, to a tree obtained by joining the centers of two stars with an edge. Unfortunately, the details are too involved to allow us to analyze trees in which the two vertices of high degree are nonadjacent.

The purpose of this note is to settle this question. The result is that there are no integral trees with precisely two vertices of degree larger than two when these two vertices are nonadjacent. This is shown using eigenvalue interlacing and Godsil's Lemma.

**Lemma 1** ('Interlacing') *Let  $T$  be a graph on  $n$  vertices with eigenvalues  $\theta_1 \geq \theta_2 \geq \dots \geq \theta_n$ , and let  $x$  be a vertex of  $T$ . Let  $T \setminus x$  have eigenvalues  $\eta_1 \geq \dots \geq \eta_{n-1}$ . Then  $\theta_1 \geq \eta_1 \geq \theta_2 \geq \eta_2 \geq \dots \geq \eta_{n-1} \geq \theta_n$ .*

**Lemma 2** (Godsil [2]) *Let  $T$  be a tree and  $\theta$  an eigenvalue of multiplicity  $m > 1$ . Let  $P$  be a path in  $T$ . Then  $\theta$  is eigenvalue of  $T \setminus P$  with multiplicity at least  $m - 1$ .*

## 2 Theorem

**Theorem 3** *Let  $T$  be a tree with integral spectrum and with precisely two vertices of degree larger than two. Then these vertices are adjacent.*

**Proof:** Let  $K_n$  denote the complete graph on  $n$  vertices. Let  $P_n$  denote the path of length  $n - 1$ , with  $n$  vertices. For a graph  $G$  on  $n$  vertices, let  $\theta_1(G), \dots, \theta_n(G)$  be its eigenvalues, where  $\theta_1(G) \geq \dots \geq \theta_n(G)$ .

Let  $u, v$  be the two vertices of degree larger than two in  $T$ , and assume that they are nonadjacent.

Apply interlacing twice to compare the spectra of  $T$  and  $T \setminus u, v$ . We find that  $\theta_1(T \setminus u, v) \geq \theta_2(T \setminus u) \geq \theta_3(T)$ . But  $T \setminus u, v$  is a union of paths, so all eigenvalues of  $T \setminus u, v$  are smaller than 2. It follows, since  $T$  is integral, that  $\theta_3(T) \leq 1$ .

Let  $w$  be a vertex separating  $u$  and  $v$ . Then  $\theta_3(T \setminus w) \leq \theta_3(T) \leq 1$ . Both components of  $T \setminus w$  have at least three vertices and hence contain the path  $P_3$  on 3 vertices which has largest eigenvalue  $\sqrt{2} > 1$ . So,  $T \setminus w$  has precisely two eigenvalues larger than one, and the same holds for  $T$ . So the spectrum of  $T$  is  $0^a \pm 1^b \pm m_1 \pm m_2$  where  $m_2 > m_1 > 1$ . (A tree is bipartite, so has symmetric spectrum.) It follows that  $T$  has at most 7 distinct eigenvalues.

As was shown by Watanabe [3], an integral tree has eigenvalue 0, so  $a$  is nonzero. If  $b = 0$ , then  $T$  has precisely 5 distinct eigenvalues, and by Godsil's Lemma cannot contain a path of length more than 4. In any case,  $T$  cannot contain a path of length more than 6.

## 2.1 Diameter 6

Suppose  $T$  contains a path  $P$  of length 6. Then  $T \setminus P$  is integral with spectrum  $0^{a-1} \pm 1^{b-1}$ . That is,  $T \setminus P$  consists of  $a - 1$  isolated points  $K_1$  and  $b - 1$  copies of  $K_2$ . It follows that  $P$  passes through  $u$  and  $v$ .

Let  $Q$  be the path between  $u$  and  $v$ , including  $u$  and  $v$ . The  $Q$  has at least 3 vertices, and  $Q$  is contained in  $P$ . Let  $P_u$  and  $P_v$  be the components of  $P \setminus u$  (resp.  $P \setminus v$ ) not containing  $v$  (resp.  $u$ ). Then  $P_u$  and  $P_v$  each contain at least one vertex, and together contain at most 4 vertices.

### 2.1.1 Pending $P_3$

Suppose first that  $P_u$  has at least three vertices. Then it has precisely three vertices, and all components of  $T \setminus v$  not containing  $u$  are single points. If some component  $P'_u$  of  $T \setminus u$  has two vertices, then let  $P'$  be the path obtained from  $P$  by replacing  $P_u$  by  $P'_u$ . Now  $T \setminus P'$  has spectrum  $0^a \pm 1^{b-2} \pm \sqrt{2}$ , violating Godsil's Lemma. It follows that  $b = 1$ . Writing down an eigenvector of  $T$  for the eigenvalue 1 yields a contradiction.

### 2.1.2 Pending $P_2$ on both sides

Suppose both  $P_u$  and  $P_v$  have 2 vertices. If  $T \setminus u$  (or  $T \setminus v$ ) has a single-point component  $S$ , then let  $P'$  be the path obtained from  $P$  by replacing  $P_u$  (or  $P_v$ ) by  $S$ . Now  $T \setminus P'$  has spectrum  $0^{a-2} \pm 1^b$  contradicting Godsil's Lemma. It follows that  $a = 1$ . But all integral trees with eigenvalue 0 of multiplicity 1 were determined in Brouwer [1], and these have only one vertex of degree more than two.

### 2.1.3 Pending $P_2$ on one side

Suppose  $P_u$  has 2 vertices and  $P_v$  is a single point. Then the path  $Q$  has 4 vertices. If  $T \setminus u$  has a single-point component  $S$  we find the same contradiction as before. Let  $P_u$  and  $P'_u$  be two components of  $T \setminus u$  with two vertices each, and let  $P''$  be the path on 5 vertices induced by  $P_u \cup \{u\} \cup P'_u$ . Then  $T \setminus P''$  has eigenvalue 1 of multiplicity  $b - 2$ , contradicting Godsil's Lemma.

### 2.1.4 Pending $K_1$ on both sides

Writing down an eigenvector of  $T$  for the eigenvalue 1 yields a contradiction except for the case of the tree  $\tilde{D}_8$  on 9 vertices, with spectrum  $0^3 \pm 1 \pm \sqrt{3} \pm 2$ , that has a nonintegral eigenvalue.

Since all subcases fail, the tree  $T$  cannot have diameter 6.

## 2.2 Diameter 5

With diameter only 5, there is no room to have pending  $P_2$ 's on both sides.

### 2.2.1 Pending $P_2$ on one side

Now the path  $Q$  has 3 vertices. Let  $T \setminus u$  have components  $pK_1 + qK_2$  not on  $v$ , and let  $T \setminus v$  have components  $rK_1$  not on  $u$ . With  $P$  a path of length 5,  $T \setminus P$  has spectrum  $0^{p+r-1} \pm 1^{q-1}$ . If  $p \neq 0$ , then let  $P'$  be a path of length 4 on  $u$  and  $v$ . Then  $T \setminus P'$  has spectrum  $0^{p+r-2} \pm 1^q$ . It follows that  $a = p + r - 1$  and  $b = q$ . Writing down an eigenvector for 1 we see that it must vanish at  $u$  and then also on the component of  $T \setminus u$  containing  $v$ , since that is a star that does not have eigenvalue 1, and the multiplicity of the eigenvalue 1 is  $q - 1$ , contradiction.

### 2.2.2 Pending $K_1$ on both sides

Now the path  $Q$  has 4 vertices, and  $T$  has a single eigenvalue 1. Writing down an eigenvector for 1 yields a contradiction.

Since both subcases fail, the tree  $T$  cannot have diameter 5.

## 2.3 Diameter 4

Now the path  $Q$  has 3 vertices. Let  $T \setminus u$  have components  $pK_1$  not on  $v$  and let  $T \setminus v$  have components  $rK_1$  not on  $u$ . Writing down the equation for a nonzero eigenvalue  $\lambda$  yields  $(\lambda^2 - p - 1)(\lambda^2 - r - 1) = 1$ . If all solutions  $\lambda$  are integers, then  $\lambda^2 - p - 1 = \lambda^2 - r - 1 = \pm 1$ , so that  $p = r$ , and the four roots are  $\pm\sqrt{p}$ ,  $\pm\sqrt{p+2}$ . But these cannot all be integers.  $\square$

## References

- [1] A. E. Brouwer, *Small integral trees*, Electr. J. Combin. **15** (2008) N1.
- [2] C. D. Godsil, *Spectra of trees*, Annals of Discrete Mathematics **20** (1984) 151–159.
- [3] M. Watanabe, *Note on integral trees*, Math. Rep. Toyama Univ. **2** (1979), 95–100.
- [4] Mamoru Watanabe & Allen J. Schwenk, *Integral starlike trees*, J. Austral. Math. Soc. (Ser A) **28** (1979) 120–128.