



Centrum voor Wiskunde en Informatica
Centre for Mathematics and Computer Science

A.E. Brouwer

Recursive constructions of mutually orthogonal latin squares

Department of Pure Mathematics

Note PM-N8501

June

The Centre for Mathematics and Computer Science is a research institute of the Stichting Mathematisch Centrum, which was founded on February 11, 1946, as a nonprofit institution aiming at the promotion of mathematics, computer science, and their applications. It is sponsored by the Dutch Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

Recursive Constructions of Mutually Orthogonal Latin Squares

A.E. Brouwer

Centre for Mathematics and Computer Science, Amsterdam.

1980 Mathematics Subject Classification: 05B15.

Key Words & Phrases: Latin squares, transversal designs.

0. INTRODUCTORY DEFINITIONS

Let S be fixed set (of 'symbols') of cardinality n . We say that two $n \times n$ matrices $A = (a_{ij})$ and $B = (b_{ij})$ with entries in S are *orthogonal* if for all $(s, t) \in S \times S$ there is a unique position (i, j) such that $a_{ij} = s$ and $b_{ij} = t$. In this chapter we shall be interested in constructing large sets of pairwise orthogonal matrices.

The connection with Latin squares is as follows: given a set $A^{(k)}$ ($k = 1, \dots, r$) of pairwise orthogonal matrices we may (after permuting the n^2 positions) assume that $A^{(1)}$ has constant rows and $A^{(2)}$ has constant columns. Now each $A^{(k)}$ ($k = 3, \dots, r$) is a Latin square, and we have found $r - 2$ mutually orthogonal Latin squares.

Conversely, given $r - 2$ mutually orthogonal Latin squares we can add two orthogonal matrices, one with constant rows and one with constant columns and get a set of r pairwise orthogonal matrices.

Clearly, for the concept of orthogonality the matrix structure does not play a rôle, that is, we might as well talk about orthogonal vectors of length n^2 . If we define an *orthogonal array* $OA(n, r)$ of order n and depth r to be an $r \times n^2$ matrix over S such that any two rows are orthogonal, then an $OA(n, r)$ is equivalent to a set of r pairwise orthogonal matrices of order n . (Similarly one might consider orthogonal arrays of strength t , that is $r \times n^t$ matrices A over S such that for any t rows i_1, \dots, i_t and any t symbols s_1, \dots, s_t there is a unique column j such that $A_{i_k j} = s_k (1 \leq k \leq t)$. Unless the contrary is explicitly mentioned, all our orthogonal arrays will have strength 2.)

Report PM-N8501

Centre for Mathematics and Computer Science

P.O. Box 4079, 1009 AB Amsterdam, The Netherlands

A more geometric picture is obtained by regarding the columns of an orthogonal array as the lines of a geometry. If R is the set of rows of the orthogonal array A , then take as point set the set $R \times S$, and let the line L_j corresponding to the j -th column be the set $L_j = \{(i, A_{ij}) \mid i \in R\}$. What we get is called a *transversal design*.

1. PAIRWISE BALANCED DESIGNS - DEFINITIONS

A *pairwise balanced design* is a set X (of *points*) together with a set \mathfrak{B} of subsets of X (called *blocks*) such that for some integer λ each 2-subset $\{x, y\}$ of X is contained in precisely λ blocks. The number λ is called the *index* of the design, and unless specified otherwise we shall always assume $\lambda = 1$. (In particular we need not worry about the possibility of repeated blocks.) When $\lambda = 1$ two blocks have at most one point in common, and the blocks are also called *lines* and the design a *linear space* (not to be confused with the linear spaces from linear algebra).

More generally, a *partial linear space* is a set X (of *points*) together with a set \mathcal{L} of subsets of X (called *lines*) such that two points are joined by at most one line.

A set of blocks \mathcal{C} of a pairwise balanced design is called a *clear set* if the elements of \mathcal{C} are pairwise disjoint. It is called a *parallel class* if \mathcal{C} is a partition of X .

If ∞ is an element not in X then there is a natural 1-1 correspondence between pairwise balanced designs (of index unity) on $X \cup \{\infty\}$ and pairwise balanced designs with designated parallel class \mathcal{C} on X : if the latter has block set \mathfrak{B} then the former has block set $(\mathfrak{B} \setminus \mathcal{C}) \cup \mathcal{C}^*$ where $\mathcal{C}^* = \{C \cup \{\infty\} \mid C \in \mathcal{C}\}$.

If a pairwise balanced design (X, \mathfrak{B}) has several pairwise disjoint parallel classes \mathcal{C}_j ($1 \leq j \leq k$) then one obtains a new pairwise balanced design by "adding points at infinity": find k new points $\infty_1, \dots, \infty_k$ and set $\bar{X} = X \cup \{\infty_1, \dots, \infty_k\}$, $\bar{\mathfrak{B}} = (\mathfrak{B} \setminus \bigcup_{j=1}^k \mathcal{C}_j) \cup \bigcup_{j=1}^k \mathcal{C}_j^* \cup \{\{\infty_1, \dots, \infty_k\}\}$, where $\mathcal{C}_j^* = \{C \cup \{\infty_j\} \mid C \in \mathcal{C}_j\}$.

(Example: construct the projective plane from the affine plane by adding 'a line at infinity').

A pairwise balanced design (X, \mathfrak{B}) is called *resolvable* if \mathfrak{B} can be partitioned into parallel classes.

A *group divisible design* is a set X (of *points*), a partition \mathcal{G} of X (the elements of which are called *groups*) and a collection \mathfrak{B} of subsets of X (the *blocks*) such that $(X, \mathfrak{B} \cup \mathcal{G})$ is a pairwise balanced design of index unity. (In other words, we have a pairwise balanced design with designated parallel class, and decide to

call the elements of this parallel class groups instead of blocks. There should be no confusion with the algebraic concept of group). (Many other concepts of group divisible design exist. A fairly standard definition says that a group divisible design with indices λ_1 and λ_2 is a set X , a partition \mathcal{G} of X and a collection \mathcal{B} of blocks such that if $x, y \in X$ are two points in the same group $G \in \mathcal{G}$ then $\{x, y\}$ is in λ_1 blocks, otherwise $\{x, y\}$ is in λ_2 blocks. HANANI [H 1975a] takes $\lambda_1 = 0, \lambda_2 = \lambda$. We take $\lambda_1 = 0, \lambda_2 = 1$. For the general concept and, more generally, for partially balanced incomplete block designs see RAGHAVARAO [RA].)

A pairwise balanced design with blocks of size k on $|X| = v$ points is called a $B(k; v)$. A group divisible design with blocks of size k and groups of size m on v points is called a $GD(k, m; v)$. A transversal design $TD(r; n)$ is a group divisible design $GD(r, n; rn)$.

If several blocks sizes may occur, we write $B(K; v)$ when each occurring block size is a member of K , and similarly $GD(K, M; v)$.

(For a study of transversal designs with index $\lambda > 1$ see HANANI [H 1975]).

2. SIMPLE CONSTRUCTIONS FOR TRANSVERSAL DESIGNS

As we have seen, the concepts of set of $(r-2)$ mutually orthogonal Latin squares and orthogonal array (of depth r) and transversal design (with r groups) are equivalent. We shall mostly use the language of transversal designs. Let $TD(r)$ be the set of all n such that a $TD(r; n)$ exists.

A. $0, 1 \in TD(r)$ for all r . If $1 < n \in TD(r)$ then $n \geq r-1$, and $r-1 \in TD(r)$ if and only if there exists a projective plane of order $r-1$.

PROOF A $TD(r; 0)$ is a design with no points and no blocks. A $TD(r; 1)$ is a design with r points, all in unique block. If $n > 1$ and $r \geq 2$ then let B be a fixed block, G a fixed group and x a point not in $B \cup G$. The $r-1$ blocks on x meeting B meet G in distinct points, so $n = |G| \geq r-1$. If $n = r-1$ then any two blocks meet, and adding a point at infinity to the parallel class formed by the groups produces a projective plane with lines of size r .

Conversely, given a projective plane with lines of size r , removal of a point yields a $TD(r; r-1)$. \square

B. [MACNEISH, BUSH] If $m, n \in TD(r)$ then $mn \in TD(r)$.

PROOF Given r pairwise orthogonal matrices $A^{(i)}$ over a symbol set S and r pairwise orthogonal matrices $B^{(i)}$ over a symbol set T ($1 \leq i \leq r$), the r matrices $A^{(i)} \times B^{(i)}$ over the symbol set $S \times T$ will be pairwise orthogonal. \square

B1. COROLLARY Let $n = \prod_i p_i^{e_i}$ be the factorization of n into prime powers. Then $n \in TD(r)$ for $r = \min_i (p_i^{e_i} + 1)$.

PROOF If q is a prime power then there exists a projective plane of order q and hence $q \in TD(r)$ for $r \leq q+1$. \square

C. [PARKER, BOSE & SHRIKHANDE] Let (X, \mathfrak{B}) be a pairwise balanced design such that for each $B \in \mathfrak{B}$ we have $|B| \in TD(r+1)$. Then $|X| \in TD(r)$.

PROOF If R is an r -set then construct a transversal design with point set $R \times X$ and groups $\{y\} \times X$ for $y \in R$ with subdesigns $TD(r; |B|)$ on $R \times B$ for each $B \in \mathfrak{B}$, taking care that each of these subdesigns contains the blocks $R \times \{b\}$ for $b \in B$. This will yield the required design. The $TD(r; |B|)$ with parallel class that we need are obtained from the given $TD(r+1; |B|)$ by throwing away one group and taking as parallel class the blocks that used to contain a fixed point of this thrown-away group. \square

This construction can be strengthened in many ways. First of all one can weaken the hypothesis " $|B| \in TD(r+1)$ " to "there exists a $TD(r; |B|)$ with a parallel class", and strengthen the conclusion to "there exists a $TD(r; |X|)$ with a parallel class".

Let us call a transversal design $TD(r; n)$ with e pairwise disjoint parallel classes a $TD_e(r; n)$. Adding points at infinity shows that a $TD_n(r; n)$ exists if and only if a $TD(r+1; n)$ exists.

A second variation on Theorem C requires $|B| \in TD_1(r)$ for all $B \in \mathfrak{B} \setminus \mathcal{C}$ where \mathcal{C} is a clear set of blocks, and $|B| \in TD(r)$ for $B \in \mathcal{C}$. (Now the conclusion is $|X| \in TD(r)$). It even suffices to ask that \mathcal{C} be an *almost clear* set, that is, that for each $C \in \mathcal{C}$ there is at most one $x \in C$ such that x is member of more than one block of \mathcal{C} .

A third version is the following.

D. [PARKER, BOSE & SHRIKHANDE] Let (X, \mathfrak{B}) be a pairwise balanced design such that \mathfrak{B} has a partition $\{\mathfrak{B}_j\}_j$, where each family \mathfrak{B}_j has blocks of constant size k_j and is either a partition of X or a symmetric 1-design on X . (Such a design is called *separable*).

Assume that $|B| \in TD(r)$ for each $B \in \mathfrak{B}$. Then $|X| \in TD(r)$.

PROOF By the previous we have $|X| \in TD(r-1)$. We shall show that a $TD(r-1; n)$ (where $n = |X|$) can be constructed so as to possess n pairwise disjoint parallel classes; then adding points at infinity will show that $n \in TD(r)$. Indeed, if B_j is a partition of X then for each $B \in \mathfrak{B}_j$ construct a transversal design $TD_k(r-1; k)$ (where $k = k_j$) with pointset $R \times B$ and groups $\{y\} \times B, y \in R$ where R is a fixed $(r-1)$ -set. If we number the parallel classes of each of these designs from 1 to k (and make sure that the 'verticals' $R \times \{b\}$ belong to parallel class 1 for all b) then the union of the parallel classes with a given number is a parallel class on $R \times X$.

On the other hand, if \mathfrak{B}_j is a symmetric 1-design on X then we cannot

construct the transversal designs on $R \times B$ ($B \in \mathfrak{B}_j$) independently. Instead, let N be the point-block incidence matrix of (X, \mathfrak{B}_j) and write N as the sum of $k = k_j$ permutation matrices N_t ($1 \leq t \leq k$). We may regard N_t as a 1-1 correspondence $\phi_t : \mathfrak{B}_j \rightarrow X$.

Let B_0 be a fixed block in \mathfrak{B}_j and construct a $TD_1(r-1, k)$ on $R \times B_0$ containing the verticals. For each non-vertical block T of this design construct a parallel class $\{T_B \mid B \in \mathfrak{B}_j\}$ with $T_{B_0} = T$ and such that for each $r \in R$ the transversal T_B contains the point $(r, \phi_t B)$, where t is determined by $(r, \phi_t B_0) \in T$.

In this way we 'transport' the transversal design on $R \times B_0$ and construct isomorphic copies on $R \times B$ for all $B \in \mathfrak{B}_j$, but in such a way that the entire collection of blocks is resolvable into parallel classes.

Taking all the blocks found in this way, and the groups $\{y\} \times X$, $y \in R$ yields the required design. \square

- E. As a modification to the previous idea of transporting a transversal design around a symmetric 1-design, suppose (X, \mathfrak{B}) is as in D. and that \mathfrak{B}_j is a symmetric 1-design. This time construct a $TD(r-1, k+1)$ on $R \times (B_0 \cup \infty)$ where ∞ is a new element. Repeating the previous construction we find for each point (r, ∞) k almost parallel classes of transversals (disjoint apart from the common point (r, ∞)); label this point now (r, ∞_i) ($1 \leq i \leq k$) so that different almost parallel classes have different points (r, ∞_i) in common. This yields:

Let (X, \mathfrak{B}) be a pairwise balanced design such that \mathfrak{B} contains pairwise disjoint families \mathfrak{B}_j such that (X, \mathfrak{B}_j) is a symmetric 1-design with block size k_j .

Suppose that $|B| \in TD(r+1)$ for $B \in \mathfrak{B} \setminus \bigcup_j \mathfrak{B}_j$ and that $|B+1| \in TD(r+1)$ for $B \in \bigcup_j \mathfrak{B}_j$. Finally suppose that $\sum_j k_j \in TD(r)$. Then $|X| + \sum_j k_j \in TD(r)$.

I shall call this construction "adding points at infinity to symmetric 1-design". Note that this terminology is misleading: we do not construct a pairwise balanced design on $n+k$ points, but only a transversal design with groups of that size.

2A. EXAMPLES

Let $N(v)$ be the maximum number of mutually orthogonal Latin squares of order v . We have $N(0) = N(1) = \infty$, $N(q) = q-1$ for prime powers q and $N(v) \leq v-1$ for arbitrary v . The statements $v \in TD(r)$ and $N(v) \geq r-2$ are equivalent.

- (i) We may apply C1 with a projective plane as design (X, \mathfrak{B}) . This yields for prime powers q that $N(q^2+q+1) \geq N(q+1)$. Usually this bound is bad, but when $q+1$ is also a prime power we get $N(q^2+q+1) \geq q$.

EXAMPLES: $N(21) \geq 4$, $N(57) \geq 7$, $N(273) \geq 16$, $N(993) \geq 31$.

- (ii) If q is a prime power then there exists a $2-(q^3+1, q+1, 1)$ design (a 'unital', the isotropic points and hyperbolic lines in the projective plane $PG(2, q^2)$ with a unitary polarity). This design is resolvable with q^2 parallel classes, and adding q^2 points at infinity yields a pairwise balanced design $B(\{q+2, q^2\}; q^3+q^2+1)$. In case also $q+2$ is a prime power, this yields $N(q^3+q^2+1) \geq q$.

EXAMPLE: $N(393) \geq 7$.

- (iii) If q_1 is an even prime power then there exists a resolvable $2-(\frac{1}{2}q(q-1), \frac{1}{2}q, 1)$ design (where points and blocks are the exterior lines and points of a hyperoval in $PG(2, q)$) with $q+1$ parallel classes. Thus we find a $B(\{\frac{1}{2}q, \frac{1}{2}q+1, x\}; \frac{1}{2}q(q-1)+x)$ by adding x points at infinity ($0 \leq x \leq q+1$), where blocksize $\frac{1}{2}q+1$ does not occur for $x=0$ and blocksize $\frac{1}{2}q$ not for $x=q+1$.

EXAMPLES:

$N(120) \geq 7$	$(q=16, \text{ design resolvable}),$
$N(136) \geq 7$	$(q=16, x=16, \text{ one parallel class of blocks of size } 8),$
$N(504) \geq 7$	$(q=32, x=8),$
$N(528) \geq 15$	$(q=32, x=32, \text{ one parallel class of blocks of size } 16),$
$N(2016) \geq 31$	$(q=64, \text{ design resolvable}).$

- (iv) Useful pairwise balanced designs can often be constructed from a projective or affine plane by throwing away a suitably chosen set of points.

Throwing away one point from $PG(2, q)$ we find a $B(\{q, q+1\}; q^2+q)$ where the blocks of size q form a parallel class. If $q+1$ is a prime power then it follows that $N(q^2+q) \geq q-1$. Examples: $N(20) \geq 3$, $N(72) \geq 7$, $N(272) \geq 15$, $N(992) \geq 30$.

Starting with $AG(2, q)$ instead we find (if $q-1, q$ are prime powers) $N(q^2-1) \geq q-2$. Examples: $N(24) \geq 3$, $N(63) \geq 6$, $N(80) \geq 7$, $N(288) \geq 15$, $N(1023) \geq 30$.

Throwing away x points from one line we find a $B(\{q+1-x, q, q+1\}; q^2+q+1-x)$ or $B(\{q-x, q-1, q\}; q^2-x)$. In this way one gets

$N(54) \geq 4$ ($q=7, x=3$), $N(280) \geq 7$ ($q=17, x=9$),
 $N(264) \geq 7$, $N(265) \geq 8$, $N(267) \geq 10$ ($q=16, x=9, 8, 6$),
 $N(285) \geq 12$ ($q=17, x=4$),
 $N(993-x) \geq 31-x$ ($q=31, x=3, 5, 7, 13, 15, 19, 21, 23$),
 $N(1024-x) \geq 31-x$ ($q=32, x=7, 9, 13, 16, 24$).

If $q \equiv 0$ or $1 \pmod{3}$ then $PG(2, q)$ contains a subconfiguration isomorphic to $AG(2, 3)$, and removing that yields a $B(\{q-2, q, q+1\}; v-9)$.

For $q=31$ this shows $N(984) \geq 27$.

Throwing away x points from a (hyper)oval in $PG(2, q)$ or $AG(2, q)$ yields a $B(\{q-1, q, q+1\}; q^2+q+1-x)$ or $B(\{q-2, q-1, q\}; q^2-x)$ for $x \leq q+1$ (or $x \leq q+2$ if q is even). Since 7, 8, 9 are three consecutive prime powers we find with $q=8$: $N(66) \geq 5$, $N(68) \geq 5$, $N(69) \geq 6$, $N(70) \geq 6$, and with $q=9$: $N(74) \geq 5$, $N(75) \geq 5$, $N(76) \geq 5$, $N(78) \geq 6$.

Note that the blocks of size 7 form a clear set in $B(\{7, 8, 9\}; 70)$ and $B(\{7, 8, 9\}; 78)$ and an almost clear set in $B(\{7, 8, 9\}; 69)$. (After writing this I found that L. ZHU (1984) had made the same observation).

- (v) Continuing in this vein we note that $PG(2, q^2)$ has a partition into Baer subplanes, and taking t of those produces a $B(\{t, q+t\}; t(q^2+q+1))$ where the collection of blocks of size $q+t$ forms a symmetric 1-design and the collection of blocks of size t is resolvable into $q^2-q+1-t$ parallel classes.

This yields many useable pairwise balanced designs

EXAMPLES: $N(189) \geq 8$ ($q=4, t=9$),
 $N(253) \geq 12$ ($q=4, t=12$, add one point at infinity to get an almost clear set of blocks of size 13),
 $N(357) \geq 9$ ($q=5, t=11$, add 16 points at infinity to the symmetric 1-design).

[For more details, see BROUWER [Br 1980]].

- (vi) Adding $q+1$ points at infinity to the symmetric 2-design $PG(2, q)$ we find if both $q+1$ and $q+2$ are prime powers: $N((q+1)^2+1) \geq q$. Examples: $N(10) \geq 2$, $N(65) \geq 7$.
- (vii) From a Singer difference set we find a separable subdesign $B(\{9, 13, 16\}; 469)$ in $PG(2, 37)$. It follows that $N(469) \geq 8$. [Again, see BROUWER, [Br 1980]].

In these examples I have listed virtually all instances I know of where the pairwise balanced design construction yields the best known bound on $N(v)$, and where the pairwise balanced design was not a (truncated) transversal design itself. (In fact, under (iv) it was a truncated transversal design). In the next section we shall see that one can do better with a transversal design as ingredient than with a general pairwise balanced design as starting point.

3. WILSON'S CONSTRUCTION

Applying the PBD construction to a transversal design $TD(m+1; t)$ of which one group has been truncated to size h (so that we have a $GD(\{m, m+1\}, \{t, u\}; mt+h)$) we find

- If $N(t) \geq m-1$ then for $0 < h < t$:
 $N(mt+h) \geq \min \{N(t), N(h), N(m)-1, N(m+1)-1\}$

- If $N(t) \geq m-2$ then $N(mt) \geq \min \{N(m)-1, N(t)\}$.

Clearly the second bound is worse than MacNeish's bound. The first one is always worse (or at least : not better) than Wilson's bound [WILSON, 1974 Thm. 2.3]

F. - If $0 < h < t$ then

$$N(mt+h) \geq \min \{N(m), N(m+1), N(t)-1, N(h)\}.$$

(For: if $m-1 \leq N(t)$ then $N(m)-1 < N(t)$, so in the first bound the minimum cannot be $N(t)$).

This bound, together with Wojtas's bound [WOJTAS, 1977]

G. - If $0 < h < t$ then

$$N(mt+h) \geq \min \{N(m), N(m+1), N(m+h), N(t)-h\}$$

account for the majority of the best lower bounds for $N(v)$ known. Both bounds follow from special cases of Wilson's construction, which we shall now describe.

Construction Ingredients: (1) A transversal design $TD(k+l; t)$ of which l groups have been truncated, so that k groups have size t and the remaining groups size h_i ($1 \leq i \leq l$) where clearly $0 \leq h_i \leq t$. Denote the union of the l truncated groups by H (so that $h := |H| = \sum_{i=1}^l h_i$). (2) Transversal designs $TD(k; h_i)$ for $1 \leq i \leq l$. (3) Transversal designs $TD(k; m + |B \cap H|)$ for each block B from the $TD(k+l; t)$ with $|B \cap H|$ pairwise disjoint blocks. We construct a $TD(k; mt+h)$ in the obvious way (given ingredients and result):

Let the $TD(k+l; t)$ have groups $G_1, \dots, G_k, H_1, \dots, H_l$, then the constructed design will have groups $(G_j \times M) \cup (H \times \{j\})$ ($j=1, 2, \dots, k$), all of size $mt+h$ (M is an arbitrary set of cardinality m); put ingredients (2) on $H_i \times K$ ($1 \leq i \leq l$) where $K = \{1, 2, \dots, k\}$; for each block B from ingredient (1) the set $(B \setminus H) \times M \cup (B \cap H) \times K$ has cardinality $k(m + |B \cap H|)$; put on this set ingredient (3) in such a way that the groups of this design are subsets of the design to be constructed and for each $b \in B \cap H$ the set $\{b\} \times K$ is a block. It is straightforward to check that this works. \square

Bound F is obtained by taking $l=1, h_1=h$. Bound G is obtained by taking $l=h, h_i=1$ ($i \leq l$). Taking $l=2, h_1=u, h_2=v$ one gets

H. - If $0 < u, v < t$ then

$$N(mt+u+v) \geq \min \{N(m), N(m+1), N(m+2), N(u), N(v), N(t)-2\}.$$

[WILSON 1974, Thm. 2.4]

3A. EXAMPLES

- (i) $N(95) \geq 6$ follows from the PBD construction using a truncated $TD(9;11)$ since $95=8.11+7$.
- (ii) $N(33) \geq 3$ follows from F . since $33=4.8+1$.
 $N(84) \geq 6$ follows from F . since $84=7.11+7$.
- (iii) $N(91) \geq 7$ follows from G . since $91=8.11+3$.
- (iv) $N(94) \geq 6$ follows from H . since $94=7.11+8+9$.
- (v) $N(90) \geq 6$ [WOLTAJ 1980 a] follows since $90=6.11+8+8+8$ and we can truncate a $TD(9;11)$ in such a way that each block meets the set H in at least one point. (In general with $l=3$ one can obtain the condition that $B \cap H \neq \emptyset$ for all B certainly when $h_1 \leq h_2$ and $(t-h_1)(t-h_2) < h_3$). Another example is $N(796) \geq 7$ since $796 = 70.11+8+8+8$.
- (vi) $N(135) \geq 7$ [BROUWER 1978] follows since $135 = 8.16+7$ and we can truncate a $TD(15;16)$ in such a way that each block meets the set H in 0,1 or 3 points, and $h_i = 1$ ($1 \leq i \leq 7$) - in fact we may take H to be a Fano subplane of $PG(2,16)$.
- (vii) $N(164) \geq 6$ follows since $164 = 7.23+3$ and we can take $h=3, h_1=h_2=h_3=1, |B \cap U| \leq 2$. In fact, for $h \leq t$ and t a prime power we can take H to be part of an oval in $PG(2,t)$ and obtain $N(mt+h) \geq \min\{N(m), N(m+1), N(m+2), N(t)-h\}$. [WILSON 1974, thm 2.5 - BROUWER 1979]
- (viii) Continuing the previous construction: if we take v points, no 3 on a line, all on different groups and $t > \binom{v}{2}$ then we can add w points all on one group and get $N(mt+w+v) \geq \min\{N(m), N(m+1), N(m+2), N(w), N(t)-v-1\}$ for $t \geq w + \binom{v}{2}$. [VAN REES]

EXAMPLES:

n	m	t	w	v	lower bound for $N(n)$
1554	81	19	13	2	8
1884	81	23	19	2	8
2046	81	25	19	2	8
2298	99	23	19	2	8
2694	99	27	19	2	8
4622	271	17	13	2	12
4776	207	23	13	2	8

I know of no examples with $v \neq 2$ where this construction yields the best known lower bound.

4. WEIGHTING AND HOLES

As was noted by WOJTAS [1980] and STINSON [1979 a] in certain special cases, and by BROUWER & VAN REES [1982] in general, one may generalize Wilson's construction by giving weights to the points of H .

In this way one constructs a transversal design $TD(k; mt + \sum_{h \in H} m_h)$, where m_h is the weight of $h (h \in H)$. Ingredient (1) is unchanged, and (2) and (3) now read:

(2') Transversal designs $TD(k; \sum_{h \in H_i} m_h)$ for $1 \leq i \leq l$.

(3') For each block B from the $TD(k+l; t)$ a transversal design $TD(k; m + \sum_{h \in B \cap H} m_h)$ with pairwise disjoint subdesigns $TD(k; m_h)$ ($h \in B \cap H$).

(The construction is entirely analogous to that in Section 3.)

But one may go further: all one needs the subdesigns in (3') for, is to throw them out in order not to cover certain pairs twice; in other words, what actually is needed is a transversal designs with holes

(3'') $TD(k; m + \sum_{h \in B \cap H} m_h) - \sum_{h \in B \cap H} TD(k; m_h)$

and (3'') may well exist while (3) does not.

Let us formally define the concept of 'transversal design with holes' - the above discussion shows that what we have in mind looks like a transversal design from which a collection of pairwise disjoint subdesigns has been removed.

A transversal design with holes $TD(k; v) - \sum_{i=1}^r TD(k; u_i)$ consists of a set X of cardinality kv (the set of points), a partition \mathcal{G} of X into k groups of v elements each, pairwise disjoint subsets Y_i of X ($1 \leq i \leq r$) of cardinality ku_i (the holes) such that $|Y_i \cap G| = u_i$ for each $G \in \mathcal{G}$ and each i , $1 \leq i \leq r$, and a collection \mathcal{B} of subsets of X of cardinality k (the blocks) such that no block meets a group or a hole in more than one point, and any two points not in the same group or hole are in a unique block.

It follows that $|\mathcal{B}| = v^2 - \sum_{i=1}^r u_i^2$. For $r=0$ the concept 'transversal design with zero holes' coincides with the usual transversal design. In case $u_i = 1$ for all i , $1 \leq i \leq r$, then $TD(k; v) - rTD(k; 1)$ (in an obvious extension of the notation) exists iff a $TD(k; v)$ with r pairwise disjoint blocks exists - showing that (3'') generalizes (3). If $TD(k; u_i)$ exists we may put it on Y_i and thus 'plug' the hole Y_i , obtaining a transversal design with $r-1$ holes. Conversely, if a transversal design (with holes) has a subdesign (disjoint from all the holes) we can unplug it and obtain a transversal design with $r+1$ holes. Not all holes can be filled: HORTON [1974], who introduced the concept 'transversal design with one hole' under the name 'incomplete array', constructs a $TD(4; 6) - TD(4; 2)$, but neither $TD(4; 6)$ nor $TD(4; 2)$ exist. (Also, BROUWER [1978] constructs $TD(6; 10) - TD(6; 2)$, while not even $TD(5; 10)$ is known.) As most important special case we find (with $l=1$) [BROUWER, 1979]

- I. - If $t = \sum_{j=1}^p h_j$ and $TD(k+1;t)$, $TD(k; \sum_{j=1}^p m_j h_j)$ and (for $j=1, \dots, p$) $TD(k; m+m_j) - TD(k; m_j)$ all exist, then also a $TD(k; mt + \sum_{j=1}^p m_j h_j)$ exists.

Instead of making holes $TD(k; m_h)$ in the ingredients (3'') corresponding to all blocks B on the point $h \in H$ we may leave one such ingredient alone and make a hole in the ingredient (2') corresponding to the group containing h . For the general formulation of this construction see BROUWER & VAN REE [1982], Theorem 1.2. The most important special case is [BROUWER, 1980 a]

- J. - If $w = \sum_{i=1}^l w_i$ and $TD(k+l;t)$, $TD(k; m)$, $TD(k; m+w)$ and (for $j=1, \dots, l$) $TD(k; m+w_j) - TD(k; w_j)$ all exist, then also $TD(k; mt+w)$ exists.

For more details about construction of transversal designs with holes, see BROUWER & VAN REES [1982].

4A. EXAMPLES

- (i) We show $N(5467) \geq 15$. The construction uses a distribution of holes as discussed above before J. Noting that $5467 = 19 \cdot 271 + 289 + 29$ we apply the construction with $k=17$, $t=19$, $m=271$, $l=2$, $h_1=17$, $h_2=13$; $289=17 \cdot 17$: the points in H_1 all get weight 17; $29 = 1 \cdot 17 + 12 \cdot 1$: one point x_0 in H_2 gets weight 17, the twelve others weight 1. We need the following ingredients:

- (i) $TD(19;19)$ exists since 19 is prime.
- (2) $TD(17;289) - 17TD(17;17)$ exists, e.g. by the PBD construction on the affine plane $AG(2,17)$.
 $TD(17;29)$ exists since 29 is prime.
- (3) $TD(17;271)$ exists since 271 is prime.
 $TD(17;272) - TD(17;1)$ exists by MacNeish: $272 = 16 \cdot 17$.
 $TD(17;288) - TD(17;17)$ exists since Wilson's construction for $TD(17;288)$ using $288 = 16 \cdot 17 + 16$ yields a design with subdesign $TD(17;17)$.
 $TD(17;289) - TD(17;17) - TD(17;1)$ exists, and is found from $AG(2,17)$.
 $TD(17;305) - TD(17;17)$ exists since Wilson's construction for $TD(17;305)$ using $305 = 16 \cdot 17 + 1$ yields a design with subdesign $TD(17;17)$.

For the standard distribution of holes we would have needed $TD(17;305) - 2TD(17;17)$, but it is not obvious how to obtain this ingredient. Therefore we cover the pairs in the km_h -subsets corresponding to points $h \in H_1$ in the designs corresponding to the (unique) block B containing h and x_0 . This yields the required $TD(17;5467)$.

- (ii) We show $N(4738) \geq 8$. (This was the largest unknown value for 8 squares; it follows that $n_8 \leq 4242$).

$$4738 = 271 \cdot 17 + (125 = 7 \cdot 17 + 6) + 6 \times 1$$

Apply the construction with $k=10$, $t=17$, $m=271$, $l=7$, $h_1=13$, $h_2=h_3=h_4=h_5=h_6=h_7=1$; give in H_1 seven points weight 17 and six points weight 1. Give all other points in H weight 1. Choose the six points on $H \setminus H_1$ on a single block B where $B \cap H_1 = \emptyset$.

- (iii) We show $N(10618) \geq 15$ and $N(10632) \geq 15$. (These were the largest unknown values for 15 squares; it follows that $n_{15} < 10000$).

$$10618 = 435 \cdot 23 + (293 = 2 \cdot 16 + 9 \cdot 29) + (320 = 20 \cdot 16)$$

$$10632 = 435 \cdot 23 + (128 = 8 \cdot 16) + (499 = 4 \cdot 16 + 15 \cdot 29)$$

Ingredients:	23, 128, 293, 499 are prime powers.
$320 = 16 \cdot 19 + 16$	shows $N(320) \geq 15$.
$435 = 16 \cdot 27 + 3 \times 1$	shows $N(435) \geq 15$.
$451 = 16 \cdot 27 + 19$	shows the existence of $TD(17; 451) - TD(17; 16)$.
$464 = 16 \cdot 29$	shows the existence of $TD(17; 464) - TD(17; 29)$.
$467 = 16 \cdot 29 + 3 \times 1$	shows the existence of $TD(17; 467) - 2TD(17; 16)$.

5. ASYMPTOTIC RESULTS

CHOWLA, ERDÖS & STRAUS [CES] showed that $\lim_{v \rightarrow \infty} N(v) = \infty$. Consequently we may define

$$n_r := \max\{v \mid N(v) < r\} \quad (\text{for } r \geq 2).$$

In fact they showed that $n_r < (3r)^{91}$, a result that was improved by ROGERS [Ro] to $n_r < r^{42}$, by WANG YUAN [WY] to $n_r < r^{26}$, by WILSON [W 1974] to $n_r < r^{17}$ and by BETH [Be] to $n_r < r^{14.8}$, all for sufficiently large r .

For small values of r explicit upper bounds for n_r have been obtained. The current state of affairs is:

$n_2 = 6$	(BOSE, SHRIKHANDE & PARKER [BSP]),
$n_3 \leq 14$	(WANG & WILSON [WaW]),
$n_4 \leq 52$	(GUÉRIN [G]),
$n_5 \leq 62$	(HANANI [H 1979]),
$n_6 \leq 76$	(WOJTAS [Wo 1980a]),
$n_7 \leq 780$,	$n_9 \leq 5842$, $n_{10} \leq 7222$ (BROUWER & VAN REES [Br vR]),
$n_8 \leq 4216$,	$n_{11} \leq 7222$, $n_{12} \leq 7286$, $n_{13} \leq 7288$,
$n_{14} \leq 7874$,	$n_{15} \leq 8360$ (BROUWER, unpublished),
$n_{30} \leq 52502$,	(BROUWER, unpublished, cf. [Br 1980a]).

The proofs are by the constructions given above (together with some explicit constructions for small v) coupled with some number theory (trivial for fixed r , sieve methods for large r) required to show that sufficiently large numbers can be written in a suitable form.

REFERENCES

- [Be] THOMAS BETH, Eine Bemerkung zur Abschätzung der Anzahl orthogonaler lateinischer Quadrate mittels Siebverfahren, *Abh. Math. Semin. Univ. Hamb.* **53** (1983) 284-288.
- [BS] R.C. BOSE & S.S. SHRIKHANDE, On the construction of sets of mutually orthogonal Latin squares and the falsity of a conjecture of Euler, *Trans. Amer. Math. Soc.* **95** (1960) 191-209.
- [BSP] R.C. BOSE, S.S. SHRIKHANDE & E.T. PARKER, Further results on the construction of mutually orthogonal Latin squares and the falsity of Euler's conjecture, *Canad. J. Math.* **12** (1960) 189-203.
- [Br 1978] A.E. BROUWER, Mutually orthogonal Latin squares, *Math. Centr. report ZN 81*, August 1978.
- [Br 1979] A.E. BROUWER, The number of mutually orthogonal Latin squares - a table up to order 10000, *Math. Centr. report ZW 123*, June 1979.
- [Br 1980] A.E. BROUWER, A series of separable designs with application to pairwise orthogonal Latin squares, *European J. Combinatorics* **1** (1980) 39-41.
- [Br 1980a] A.E. BROUWER, On the existence of 30 mutually orthogonal Latin squares, *Math. Centr. report ZW 136*, Jan 1980.
- [BrvR] A.E. BROUWER & G.H.J. VAN REES, More mutually orthogonal Latin squares, *Discrete Math.* **39** (1982) 263-281.
- [Bu] K.A. BUSH, A generalization of a theorem due to MacNeish, *Ann. Math. Stat.* **23** (1952) 293-295.
- [CES] S. CHOWLA, P. ERDÖS & E.G. STRAUS, On the maximal number of pairwise orthogonal Latin squares of a given order, *Canad. J. Math.* **12** (1960) 204-208.
- [DS] J.H. DINITZ & D.R. STINSON, MOLS with holes, *Discr. Math.* **44** (1983) 145-154.
- [DJM] A.L. DULMAGE, D.M. JOHNSON & N.S. MENDELSON, Orthomorphisms of groups and orthogonal Latin squares, I, *Canad. J. Math* **13** (1961) 356-372.
- [G] R. GUÉRIN, Existence et propriétés des carrés latins orthogonaux II, *Publ. Inst. Statist. Univ. Paris* **15** (1966) 215-293. MR **35** (1968) # 4118.
- [H 1970] H. HANANI, On the number of orthogonal Latin squares, *J. Combinatorial Theory* **8** (1970) 247-271.
- [H 1975] H. HANANI, On transversal designs, In: *Combinatorics, Proceedings of the Advanced Study Institute on Combinatorics held at Nijenrode Castle, Breukelen, 1974*, M. Hall, jr. & J.H. van Lint (eds.) pp. 42-52.
- [H 1975a] H. HANANI, Balanced incomplete block designs and related designs, *Discr. Math.* **11** (1975) 255-369.
- [Ho] J.D. HORTON, Sub-Latin squares and incomplete orthogonal arrays, *J. Combinatorial Theory (A)* **16** (1974) 23-33.
- [vL] J.H. VAN LINT, *Combinatorial Theory Seminar, Lecture Notes in Math.* **382**, Springer, Berlin, 1974.

- [MacN] H.F. MACNEISH, Euler Squares, *Ann. Math.* **23** (1922) 221-227.
- [Mi] W.H. MILLS, Some mutually orthogonal Latin squares, *Proc. 8th S-E Conf. on Combinatorics, Graph Theory and Computing* (1977) 473-487.
- [MuSSV 1978] R.C. MULLIN, P.J. SCHELLENBERG, D.R. STINSON & S.A. VANSTONE, On the existence of 7 and 8 mutually orthogonal Latin squares, *Dept. of Combinatorics & Optimization Research Report CORR 78-14* (1978), Univ. of Waterloo.
- [MuSSV 1980] R.C. MULLIN, P.J. SCHELLENBERG, D.R. STINSON & S.A. VANSTONE, Some results on the existence of squares, *Ann. Discr. Math.* **6** (1980) 257-274.
- [P] E. PARKER, Construction of some sets of mutually orthogonal Latin squares, *Proc. Amer. Math. Soc.* **10** (1959) 946-949.
- [Ra] D. RAGHAVARAO, *Constructions and combinatorial problems in design of experiments*, J. Wiley & Sons, New York, 1971.
- [Ro] K. ROGERS, A note on orthogonal Latin squares, *Pacific J. Math.* **14** (1964) 1395-1397.
- [S 1978] D.R. STINSON, A note on the existence of 7 and 8 mutually orthogonal Latin squares, *Ars Combinatoria* **6** (1978) 113-115.
- [S 1979] D.R. STINSON, On the existence of 30 mutually orthogonal Latin squares, *Ars Combinatoria* **7** (1979) 153-170.
- [S 1979a] D.R. STINSON, A generalization of Wilson's construction for mutually orthogonal Latin squares, *Ars Combinatoria* **8** (1979) 95-105.
- [Sz] K. SZAJOWSKI, The number of orthogonal Latin squares, *Applicationes Mathematicae* **15** (1976) 85-102.
- [vR] G.H.J. VAN REES, A corollary to a theorem of Wilson, *Dept of Combinatorics & Optimization Research Report CORR 78-15* (1978), Univ. of Waterloo.
- [WaW] S.M.P. WANG & R.M. WILSON, A few more squares II, *Proc. 9th S-E Conf. on Combinatorics, Graph Theory & Computing* (1978) p.688 (Abstract).
- [W 1974] R.M. WILSON, Concerning the number of mutually orthogonal Latin squares, *Discr. Math.* **9** (1974) 181-198.
- [W 1974a] R.M. WILSON, A few more squares, *Proc. 5th S-E Conf. on Combinatorics, Graph Theory & Computing* (1974) 675-680.
- [Wo 1977] M. WOJTAS, On seven mutually orthogonal Latin squares, *Discr. Math.* **20** (1977) 193-201.
- [Wo 1980] M. WOJTAS, New Wilson-type constructions of mutually orthogonal Latin squares, *Discr. Math.* **32** (1980) 191-199.
- [Wo 1980a] M. WOJTAS, A note on mutually orthogonal Latin squares, *Analiza Dyskretna, Prace Nauk. Inst. Mat. Politech. Wrocław* **19** (1980) 11-14.
- [Wo 1981] M. WOJTAS, letter dated 810305.
- [WY] WANG YUAN, On the maximal number of pairwise orthogonal Latin squares of order s ; an application of the sieve method, *Chinese Math.* **8** (1966) 422-432.

[Z] L. ZHU, Six pairwise orthogonal Latin squares of order 69, J. Austral. Math. Soc. (Series A) 37 (1984) 1-3.

