

Final examination Logic & Set Theory (2IT61/2IT07)

Thursday October 30, 2014, 9:00–12:00 hrs.

You are **not** allowed to use any books, notes, or other course material. Your solutions to the problems have to be formulated and written down in a clear and precise manner.

- (2) 1. Prove that the formulas

$$a \Rightarrow (b \vee c) \quad \text{and} \quad (b \Rightarrow a) \wedge \neg c$$

are incomparable.

- (1) 2. Prove with a *calculation* (i.e., using the formal system based on standard equivalences described in *Part I* of the book) that

$$P \Rightarrow \neg Q \stackrel{val}{=} \neg(P \wedge Q) .$$

3. Let \mathbb{P} be the set of all people, let Anna denote a particular person in \mathbb{P} , and let \mathbb{B} be the set of all books. Furthermore, let R and L be predicates on $\mathbb{P} \times \mathbb{B}$ with the following interpretations for all $p \in \mathbb{P}$ and $b \in \mathbb{B}$:

$R(p, b)$ means ‘ p has read b ’, and
 $L(p, b)$ means ‘ p liked b ’.

Give formulas of predicate logic that express the following statements:

- (1) (a) Everybody has read a book.
(1) (b) Anna has only read books she liked.

- (3) 4. Prove with a *derivation* (i.e., using the methods described in *Part II* of the book) that the formula

$$(\forall x[P(x) \Rightarrow \neg Q(x)] \wedge \exists y[P(y) : Q(y) \vee R(y)]) \Rightarrow \exists z[P(z) \wedge R(z)]$$

is a tautology.

Final examination Logic & Set Theory (2IT61/2IT07)

Thursday October 30, 2014, 9:00–12:00 hrs.

- (2) 5. Prove that the following formula holds for all sets A , B and C :

$$A \cap B = A \cap C \Rightarrow A \cap (B \setminus C) = \emptyset .$$

- (3) 6. Let the sequence a_0, a_1, a_2, \dots be inductively defined by

$$\begin{aligned} a_0 &:= 0 \\ a_1 &:= 1 \\ a_{i+2} &:= 3a_{i+1} - 2a_i \quad (i \in \mathbb{N}). \end{aligned}$$

Prove that $a_n = 2^n - 1$ for all $n \in \mathbb{N}$.

7. Let A and B be sets, and let $F : A \rightarrow B$ and $G : B \rightarrow A$ be mappings such that $\forall x[x \in A : G(F(x)) = x]$.

- (2) (a) Prove that F is an injection.
(1) (b) Show, with a counterexample, that F is not necessarily a surjection.

8. We define

$$V := \mathcal{P}(\{0, 1, 2\}) \setminus \{\{0, 1\}, \{1, 2\}, \{0, 1, 2\}\} .$$

- (1) (a) Determine V .
(2) (b) Make a Hasse diagram of $\langle V, \subseteq \rangle$.
(1) (c) What are the minimal elements of V in $\langle V, \subseteq \rangle$?
What are the maximal elements of V in $\langle V, \subseteq \rangle$?

The number between parentheses in front of a problem indicates how many points you score with a correct answer to it. A partially correct answer is sometimes awarded with a fraction of those points. The grade for this examination will be determined by dividing the total number of scored points by 2.