

Remco van der Hofstad · Gordon Slade

# A generalised inductive approach to the lace expansion

Received: 13 September 2000 / Revised version: 16 May 2001 /  
Published online: 20 December 2001 – © Springer-Verlag 2001

**Abstract.** The lace expansion is a powerful tool for analysing the critical behaviour of self-avoiding walks and percolation. It gives rise to a recursion relation which we abstract and study using an adaptation of the inductive method introduced by den Hollander and the authors. We give conditions under which the solution to the recursion relation behaves as a Gaussian, both in Fourier space and in terms of a local central limit theorem. These conditions are shown elsewhere to hold for sufficiently spread-out models of networks of self-avoiding walks in dimensions  $d > 4$ , and for sufficiently spread-out models of critical oriented percolation in dimensions  $d + 1 > 5$ , providing a unified approach and an essential ingredient for a detailed analysis of the branching behaviour of these models.

---

## 1. The recursion relation and results

### 1.1. The recursion relation

When applied to self-avoiding walks or oriented percolation, the lace expansion gives rise to a convolution recursion relation of the form

$$f_{n+1}(k; z) = \sum_{m=1}^{n+1} g_m(k; z) f_{n+1-m}(k; z) + e_{n+1}(k; z) \quad (n \geq 0), \quad (1.1)$$

with  $f_0(k; z) = 1$ . Here,  $k \in [-\pi, \pi]^d$  is a parameter dual to a spatial lattice variable  $x \in \mathbb{Z}^d$ , and  $z$  is a positive parameter. The functions  $g_m$  and  $e_m$  are to be regarded as given, and the goal is to understand the behaviour of the solution  $f_n(k; z)$  of (1.1). We begin with three examples of (1.1).

#### 1.1.1. Example: Random walk

Let  $D$  be a non-negative function on  $\mathbb{Z}^d$  which respects the lattice symmetries of reflection and rotation and obeys  $\sum_{x \in \mathbb{Z}^d} D(x) = 1$ . Let  $z > 0$ , and, for  $n \geq 1$ , let  $q_n(x; z)$  denote the  $n$ -step transition function of the random walk on  $\mathbb{Z}^d$  with

---

R. van der Hofstad: Stieltjes Institute for Mathematics, Delft University, Mekelweg 4, 2628 CD Delft, The Netherlands. e-mail: R.W.vanderHofstad@its.tudelft.nl

G. Slade: Department of Mathematics, University of British Columbia, Vancouver, BC V6T 1Z2, Canada. e-mail: slade@math.ubc.ca

*Mathematics Subject Classification (2000):* 60K35, 82B41, 82B43

1-step transition function  $D(x)$ , multiplied by  $z^n$ . In particular,  $q_1(x; z) = zD(x)$ . Set  $q_0(x; z) = \delta_{0,x}$ . Then  $q_n$  obeys

$$q_{n+1}(x; z) = \sum_{y \in \mathbb{Z}^d} zD(y)q_n(x - y; z) \quad (n \geq 0). \tag{1.2}$$

Given an absolutely summable function  $h$  on  $\mathbb{Z}^d$ , we denote its Fourier transform by

$$\hat{h}(k) = \sum_{x \in \mathbb{Z}^d} h(x)e^{ik \cdot x} \quad (k \in [-\pi, \pi]^d). \tag{1.3}$$

Then (1.2) implies that

$$\hat{q}_{n+1}(k; z) = z\hat{D}(k)\hat{q}_n(k; z) \quad (n \geq 0). \tag{1.4}$$

The recursion relation (1.4) is in the form (1.1), with  $f_n(k; z) = \hat{q}_n(k; z)$ ,  $g_1(k; z) = z\hat{D}(k)$ ,  $g_m = 0$  for  $m \geq 2$ , and  $e_n = 0$  for  $n \geq 1$ . It is readily solved to give  $\hat{q}_n(k; z) = z^n \hat{D}(k)^n$ . Assuming that  $D$  has a finite second moment  $\sigma^2 = \sum_x |x|^2 D(x)$ , Taylor expansion of  $\hat{D}$  then gives the central limit theorem

$$\lim_{n \rightarrow \infty} \hat{q}_n\left(\frac{k}{\sqrt{\sigma^2 n}}; 1\right) = e^{-k^2/2d}. \tag{1.5}$$

Our goal in this paper is to prove that the solutions of certain more elaborate recursion relations still exhibit this Gaussian behaviour for a critical value of  $z$ . In (1.5), the value  $z = 1$  plays the role of a critical point. For general  $z > 0$ , the right side of (1.5) includes a factor  $z^n$ , which grows exponentially for  $z > 1$  and decays exponentially for  $z < 1$ . The introduction of  $z$  in this example is artificial, but in the following two examples the role of the critical point will be more profound.

### 1.1.2. Example: Self-avoiding walk

An  $n$ -step walk is a mapping  $\omega : \{0, \dots, n\} \rightarrow \mathbb{Z}^d$ . An  $n$ -step self-avoiding walk is an  $n$ -step walk with  $\omega(i) \neq \omega(j)$  for all  $i \neq j$ . Let  $\mathcal{R}_n(x)$  denote the set of  $n$ -step walks with  $\omega(0) = 0$  and  $\omega(n) = x$ , and let  $\mathcal{C}_n(x)$  denote the subset of  $\mathcal{R}_n(x)$  consisting of  $n$ -step self-avoiding walks. For  $\omega \in \mathcal{R}_n(x)$ , let

$$W(\omega) = \prod_{i=1}^n D(\omega(i) - \omega(i - 1)), \tag{1.6}$$

where  $D$  is as in Section 1.1.1. We define  $c_0(x) = \delta_{0,x}$ , and, for  $n \geq 1$ ,

$$c_n(x) = \sum_{\omega \in \mathcal{C}_n(x)} W(\omega). \tag{1.7}$$

Without loss of generality, we require here that  $D(0) = 0$ .

The lace expansion [6, 17] gives rise to an identity

$$c_{n+1}(x) = \sum_{y \in \mathbb{Z}^d} c_1(y)c_n(x - y) + \sum_{y \in \mathbb{Z}^d} \sum_{m=2}^{n+1} \pi_m(y)c_{n+1-m}(x - y) \quad (n \geq 0), \tag{1.8}$$

where  $\pi_m(y)$  is an explicit function. The identity (1.8) is discussed in more detail in Section 1.4.1, where its derivation via repeated application of inclusion-exclusion is described. Multiplying by  $z^{n+1}$ , using  $c_1(x) = D(x)$ , and applying the Fourier transform, (1.8) becomes

$$\hat{c}_{n+1}(k)z^{n+1} = z\hat{D}(k)\hat{c}_n(k)z^n + \sum_{m=2}^{n+1} \hat{\pi}_m(k)z^m\hat{c}_{n+1-m}(k)z^{n+1-m} \quad (n \geq 0). \tag{1.9}$$

This is in the form of (1.1), with  $f_n(k; z) = \hat{c}_n(k)z^n$ ,  $g_1(k; z) = z\hat{D}(k)$ ,  $g_m(k; z) = \hat{\pi}_m(k)z^m$  for  $m \geq 2$ , and  $e_n(k; z) = 0$ .

An important feature of the lace expansion is that  $\pi_m(x)$  can be estimated in terms of  $c_j(x)$  for  $j$  strictly less than  $m$ . This allows the right side of (1.8) to be analysed using information about  $c_j(x)$  for  $j \leq n$ , which opens up the possibility of an inductive analysis of (1.8). Such an inductive analysis was introduced by den Hollander and the authors [13] to study weakly self-avoiding walk for  $d > 4$ . Our inductive analysis of (1.1) extends and generalises the analysis of [13]. A small parameter is introduced by considering the ‘‘spread-out’’ models defined below. Consequently  $\pi_m$  turns out to be small for  $m \geq 2$  when  $d > 4$ , both uniformly in  $m$  and in terms of its rapid decay as  $m$  becomes large. The recursion relation (1.8) can thus be considered to be a small perturbation of the simple equation (1.4).

We will make a set of assumptions on (1.1), motivated by (1.9) and related identities for other models. Under these assumptions, we will prove that solutions to (1.1) exhibit Gaussian behaviour at a critical value  $z_c$  of  $z$ . Our assumptions on (1.1) are verified for sufficiently spread-out models of self-avoiding walk, in [15]. Our results below then imply, in particular, the result of [17] that for  $d > 4$  and sufficiently spread-out models there are positive  $A, v$  and  $z_c$  such that

$$\lim_{n \rightarrow \infty} \hat{c}_n\left(\frac{k}{\sqrt{\sigma^2 vn}}\right)z_c^n = Ae^{-k^2/2d}. \tag{1.10}$$

The critical value  $z_c$  is closely related to the connective constant for self-avoiding walk. This is best seen for the example where  $D$  is uniform on  $\mathbb{Z}^d \cap [-L, L]^d \setminus \{0\}$ . Let  $\Lambda_L$  denote the cardinality of this set, and let  $N_n$  denote the number of  $n$ -step self-avoiding walks with steps in  $\Lambda_L$ . Then  $\hat{c}_n(0) = N_n \Lambda_L^{-n}$ . The connective constant for the spread-out self-avoiding walk is  $\mu_L = \lim_{n \rightarrow \infty} N_n^{1/n}$ , where the limit exists by a well-known subadditivity argument [17]. This is consistent with (1.10) if and only if  $z_c = \Lambda_L \mu_L^{-1}$ .

### 1.1.3. Example: Oriented percolation

Consider the graph with vertices  $\mathbb{Z}^d \times \mathbb{Z}_+$  and directed bonds  $((x, n), (y, n + 1))$ , for  $n \geq 0$  and  $x, y \in \mathbb{Z}^d$ . Let  $z \in [0, \|D\|_\infty^{-1}]$ , so that  $zD(y - x) \leq 1$ . We associate to each directed bond  $((x, n), (y, n + 1))$  an independent random variable taking the value 1 with probability  $zD(y - x)$  and 0 with probability  $1 - zD(y - x)$ . We say a bond is *occupied* when the corresponding random variable takes the value 1, and *vacant* when the random variable is 0. The joint probability distribution of the bond variables will be denoted  $\mathbb{P}_z$ . We say that  $(x, n)$  is *connected* to  $(y, m)$ , and write  $(x, n) \rightarrow (y, m)$ , when there is an oriented path from  $(x, n)$  to  $(y, m)$  consisting of occupied bonds, or if  $(x, n) = (y, m)$ . Given  $z \in [0, \|D\|_\infty^{-1}]$ ,  $n \geq 0$  and  $x \in \mathbb{Z}^d$ , we define the two-point function:

$$\tau_n(x; z) = \mathbb{P}_z((0, 0) \rightarrow (x, n)). \tag{1.11}$$

As we explain in more detail in Section 1.4.2, the lace expansion gives rise to a function  $\hat{\pi}_m(k; z)$  (not equal to the function  $\hat{\pi}_m(k)$  for self-avoiding walk) for which

$$\begin{aligned} \hat{\tau}_{n+1}(k; z) &= z\hat{D}(k)\tau_n(k; z) + z\hat{D}(k) \sum_{m=2}^n \hat{\pi}_m(k; z)\hat{\tau}_{n-m}(k; z) \\ &+ \hat{\pi}_{n+1}(k; z) \quad (n \geq 0). \end{aligned} \tag{1.12}$$

Equation (1.12) is a special case of (1.1) with the choices  $f_n(k; z) = \hat{\tau}_n(k; z)$ ,  $e_n(k; z) = \hat{\pi}_n(k; z)$ ,  $g_1(k; z) = z\hat{D}(k)$ ,  $g_2(k; z) = 0$ ,  $g_m(k; z) = z\hat{D}(k) \times \hat{\pi}_{m-1}(k; z)$  ( $m \geq 3$ ). (By definition,  $\hat{\tau}_0(0; z) = 1$  and  $\hat{\pi}_1(k; z) = 0$ .) Our assumptions on (1.1) are verified in [14] for sufficiently spread-out models of oriented percolation when  $d > 4$ . Our results below then imply, in particular, the result of [20] that for  $d > 4$  and for sufficiently spread-out models, there are positive constants  $z_c$ ,  $A$  and  $\nu$  such that

$$\lim_{n \rightarrow \infty} \hat{\tau}_n\left(\frac{k^2}{\sqrt{\sigma^2 \nu n}}\right) = Ae^{-k^2/2d}. \tag{1.13}$$

Here  $z_c$  is the critical probability for spread-out oriented percolation, in the sense that the origin is connected to an infinite cluster of occupied bonds with probability 0 when  $z < z_c$  and with positive probability when  $z > z_c$ .

### 1.2. Assumptions on the recursion relation

We will prove Gaussian behaviour of  $f_n$  under certain assumptions on  $e_n$  and  $g_n$ , as well as some modest assumptions on  $f_n$ . Our assumptions are shown elsewhere to apply for sufficiently spread-out models of self-avoiding walks on  $\mathbb{Z}^d$  [15] and critical oriented bond percolation on  $\mathbb{Z}^d \times \mathbb{Z}_+$  [14], in dimensions  $d > 4$ . Our results form the basis for a proof of Gaussian behaviour of sufficiently spread-out polymer networks for  $d > 4$ , and for a proof that the moment measures of sufficiently spread-out critical oriented percolation converge to those of super-Brownian

motion for  $d > 4$ . In this paper, we extract an important model-independent aspect of the proof of these results. The precise form of the functions  $e_n$  and  $g_n$  is model-dependent, and the verification of the assumptions on them does depend on the model.

Before stating the assumptions, we first introduce some notation. For  $x = (x_1, \dots, x_d)$  in  $\mathbb{Z}^d$  or  $\mathbb{R}^d$ , we write  $\|x\|_p = (\sum_{i=1}^d |x_i|^p)^{1/p}$  for  $p \in [1, \infty)$ , and  $\|x\|_\infty = \max_{i \in \{1, \dots, d\}} |x_i|$ . We also write  $x^2 = |x|^2 = \|x\|_2^2$  for the square of the Euclidean norm of  $x$ . For a function  $h$  on  $[-\pi, \pi]^d$ , we write  $\|h\|_p = [(2\pi)^{-d} \int_{[-\pi, \pi]^d} |h(k)|^p d^d k]^{1/p}$ .

Throughout this paper, we make the following four assumptions, Assumptions S, D, E, G, on the quantities appearing in the recursion equation (1.1).

The first assumption, Assumption S, requires that the functions appearing in (1.1) respect the lattice symmetries of reflection and rotation, and that  $f_n$  remains bounded in a weak sense.

**Assumption S.** For every  $n \in \mathbb{N}$  and  $z > 0$ , the mapping  $k \mapsto f_n(k; z)$  is symmetric under replacement of any component  $k_i$  of  $k$  by  $-k_i$ , and under permutations of the components of  $k$ . The same holds for  $e_n(\cdot; z)$  and  $g_n(\cdot; z)$ . In addition, for each  $n$ ,  $|f_n(k; z)|$  is bounded uniformly in  $k \in [-\pi, \pi]^d$  and  $z$  in a neighbourhood of 1 (which may depend on  $n$ ).

Our next assumption, Assumption D, incorporates a ‘‘spread-out’’ aspect to the recursion equation. It introduces a function  $D$  which defines the underlying random walk model, about which (1.1) is a perturbation. The assumption involves a non-negative parameter  $L$ , which will be taken to be large, and which serves to spread out the steps of the random walk over a large set. We write  $D = D_L$  in the statement of Assumption D to emphasise this dependence, but the subscript will not be retained elsewhere. An example of a family of  $D$ ’s obeying the assumption is given following its statement. Assumption D implies, in particular, that  $D$  has a finite second moment, and we define

$$\sigma^2 = -\nabla^2 \hat{D}(0). \tag{1.14}$$

Our assumptions will involve a parameter  $d$ , which corresponds to the spatial dimension in our applications. Our theorems will assume that  $d > 4$ . We will write

$$a(k) = 1 - \hat{D}(k). \tag{1.15}$$

**Assumption D.** We assume that

$$f_1(k; z) = z \hat{D}_L(k), \quad e_1(k; z) = 0. \tag{1.16}$$

In particular, this implies that  $g_1(k; z) = z \hat{D}_L(k)$ . As part of Assumption D, we also assume:

(i)  $D_L$  is normalised so that  $\hat{D}_L(0) = 1$ , and has  $2 + 2\epsilon$  moments for some  $\epsilon > 0$ , i.e.,

$$\sum_{x \in \mathbb{Z}^d} |x|^{2+2\epsilon} D_L(x) < \infty. \tag{1.17}$$

(ii) There is a constant  $C$  such that, for all  $L \geq 1$ ,

$$\|D_L\|_\infty \leq CL^{-d}, \quad \sigma^2 = \sigma_L^2 \leq CL^2, \tag{1.18}$$

(iii) There exist constants  $\eta, c_1, c_2 > 0$  such that

$$c_1L^2k^2 \leq a_L(k) \leq c_2L^2k^2 \quad (\|k\|_\infty \leq L^{-1}), \tag{1.19}$$

$$a_L(k) > \eta \quad (\|k\|_\infty \geq L^{-1}), \tag{1.20}$$

$$a_L(k) < 2 - \eta \quad (k \in [-\pi, \pi]^d). \tag{1.21}$$

*Example.* Let  $h$  be a non-negative bounded function on  $\mathbb{R}^d$  which is almost everywhere continuous, and which is symmetric under the lattice symmetries of reflection in coordinate hyperplanes and rotations by ninety degrees. Assume that there is an integrable function  $H$  on  $\mathbb{R}^d$  with  $H(te)$  non-increasing in  $t \geq 0$  for every unit vector  $e \in \mathbb{R}^d$ , such that  $h(x) \leq H(x)$  for all  $x \in \mathbb{R}^d$ . Assume that  $\int_{\mathbb{R}^d} h(x)d^d x = 1$  and  $\int_{\mathbb{R}^d} |x|^{2+2\epsilon} h(x)d^d x < \infty$  for some  $\epsilon > 0$ . The monotonicity and integrability hypotheses on  $H$  imply that  $\sum_{x \in \mathbb{Z}^d} h(x/L) < \infty$  for all  $L$ . We will verify in Appendix A that the function

$$D_L(x) = \frac{h(x/L)}{\sum_{x \in \mathbb{Z}^d} h(x/L)} \tag{1.22}$$

obeys the conditions of Assumption D, when  $L$  is large enough. A good example is given by the function  $h(x) = 2^{-d}$  for  $x \in [-1, 1]^d$ ,  $h(x) = 0$  otherwise. In this case,  $D_L$  is uniform on the cube  $[-L, L]^d \cap \mathbb{Z}^d$ .

We will take  $L$  large, yielding a small parameter

$$\beta = L^{-d}. \tag{1.23}$$

Each of  $e_n, f_n$  and  $g_n$  depend on  $L$ , but we do not make this dependence explicit in the notation.

To help motivate the remaining two assumptions we will make on (1.1), it is useful to discuss the bounds on  $\pi_m(x)$  for self-avoiding walk in more detail. As we explain in Section 1.4.1, one contribution to  $\pi_m(x)$  is

$$\pi_m^{(2)}(x) = \sum_{\substack{m_1, m_2, m_3 \geq 1 \\ m_1 + m_2 + m_3 = m}} \prod_{j=1}^3 \sum_{\omega_j \in \mathcal{C}_{m_j}(x)} W(\omega_j) I(\omega_1, \omega_2, \omega_3), \tag{1.24}$$

where  $I(\omega_1, \omega_2, \omega_3)$  is equal to 1 if the  $\omega_i$  are pairwise mutually avoiding apart from their common endpoints, and otherwise equals 0. The mutual avoidance of the three walks in (1.24), and the more complicated pattern of avoidances occurring in higher order contributions to  $\pi_m(x)$ , make  $\pi_m(x)$  difficult to analyse exactly.

However, upper bounds can be obtained by neglecting the mutual avoidance of the  $\omega_j$ , yielding

$$\begin{aligned}
 |\hat{\pi}_m^{(2)}(k)| &\leq \sum_x \pi_m^{(2)}(x) \leq \sum_x \sum_{\substack{m_1, m_2, m_3 \geq 1 \\ m_1 + m_2 + m_3 = m}} \prod_{j=1}^3 c_{m_j}(x) \\
 &\leq 3! \sum_{\substack{m_1 \geq m_2 \geq m_3 \geq 1 \\ m_1 + m_2 + m_3 = m}} \|c_{m_1}\|_\infty \|c_{m_2}\|_\infty \|c_{m_3}\|_1. \tag{1.25}
 \end{aligned}$$

Here we used the notation  $\|c_l\|_p = [\sum_{x \in \mathbb{Z}^d} |c_l(x)|^p]^{1/p}$  (for  $1 \leq p < \infty$ ) and  $\|c_l\|_\infty = \sup_x |c_l(x)|$  for functions on  $\mathbb{Z}^d$ . Thus  $\hat{\pi}_m^{(2)}$  can be bounded in terms of norms of  $c_l$  with  $l < m$ . Similar estimates are possible for  $\hat{\pi}_m$ . The fact that  $l$  here is strictly less than  $n$  is what enables the inductive approach to succeed.

Suppose that bounds  $\hat{c}_l(0) \leq K$  and  $\|\hat{c}_l\|_1 \leq K\beta l^{-d/2}$  were true, for  $l < m$ . Then  $\|c_l\|_1 = \hat{c}_l(0) \leq K$  and  $\|c_l\|_\infty \leq \|\hat{c}_l\|_1 \leq K\beta l^{-d/2}$  for  $l < m$ . These bounds are consistent with the Gaussian behaviour of ordinary random walk. Applying these bounds to (1.25) then gives

$$|\hat{\pi}_m^{(2)}(k)| \leq 3! K^3 \beta^2 \sum_{\substack{m_1 \geq m_2 \geq m_3 \geq 1 \\ m_1 + m_2 + m_3 = m}} m_1^{-d/2} m_2^{-d/2} \leq CK^3 \beta^2 m^{-d/2}, \tag{1.26}$$

where the last inequality uses  $d > 4$ .

We will assume the existence of a similar structure for the general recursion relation (1.1). The next two assumptions summarise the manner in which bounds on  $f_m$  are assumed to imply bounds on  $e_n$  and  $g_n$ . Verification of these assumptions requires a model-dependent analysis, which is carried out for self-avoiding walks in [15] and for oriented percolation in [14]. Further discussion of this is given in Section 1.4. However, given the assumptions, the remaining inductive analysis, which we present in this paper, is model-independent.

The relevant bounds on  $f_m$ , which *a priori* may or may not be satisfied, are

$$\|\hat{D}^2 f_m(\cdot; z)\|_1 \leq K\beta m^{-d/2}, \quad |f_m(0; z)| \leq K, \quad |\nabla^2 f_m(0; z)| \leq K\sigma^2 m, \tag{1.27}$$

for some positive constant  $K$ . The factor  $\hat{D}(k)^2$  appearing in the first bound can be understood from the fact that  $\|\hat{D}^2\|_1$  is the probability that a random walk taking steps with step distribution  $D$  returns to the origin after two steps, which is bounded by  $\|D\|_\infty \leq \mathcal{O}(\beta)$ . Because of this, the factor  $\hat{D}(k)^2$  is helpful in extracting the small factor  $\beta$  appearing in the upper bound.

**Assumption E.** There is an  $L_0$ , an interval  $I \subset [1 - \alpha, 1 + \alpha]$  with  $\alpha \in (0, 1)$ , and a function  $K \mapsto C_e(K)$ , such that if (1.27) holds for some  $K > 1$ ,  $L \geq L_0$ ,  $z \in I$  and for all  $1 \leq m \leq n$ , then for that  $L$  and  $z$ , and for all  $k \in [-\pi, \pi]^d$  and  $2 \leq m \leq n + 1$ , the following bounds hold:

$$|e_m(k; z)| \leq C_e(K)\beta m^{-d/2}, \quad |e_m(k; z) - e_m(0; z)| \leq C_e(K)a(k)\beta m^{-(d-2)/2}.$$

**Assumption G.** There is an  $L_0$ , an interval  $I \subset [1 - \alpha, 1 + \alpha]$  with  $\alpha \in (0, 1)$ , and a function  $K \mapsto C_g(K)$ , such that if (1.27) holds for some  $K > 1$ ,  $L \geq L_0$ ,  $z \in I$  and for all  $1 \leq m \leq n$ , then for that  $L$  and  $z$ , and for all  $k \in [-\pi, \pi]^d$  and  $2 \leq m \leq n + 1$ , the following bounds hold:

$$|g_m(k; z)| \leq C_g(K)\beta m^{-d/2}, \quad |\nabla^2 g_m(0; z)| \leq C_g(K)\sigma^2 \beta m^{-(d-2)/2},$$

$$|\partial_z g_m(0; z)| \leq C_g(K)\beta m^{-(d-2)/2},$$

$|g_m(k; z) - g_m(0; z) - a(k)\sigma^{-2}\nabla^2 g_m(0; z)| \leq C_g(K)\beta a(k)^{1+\epsilon'} m^{-(d-2-2\epsilon')/2}$ , with the last bound valid for any  $\epsilon' \in [0, \epsilon]$ , with  $\epsilon$  given in (1.17).

The last bound of Assumption G is effectively a bound on the error in a second order Taylor expansion of  $g_m$ , since  $a(k)\sigma^{-2}$  is asymptotic to  $k^2/(2d)$  as  $k \rightarrow 0$ .

We emphasize that: (1) we do not assume that the bounds (1.27) are true, but rather that they imply the bounds of Assumptions E and G, and (2) it is the assumption that bounds on  $f_m$  for  $m \leq n$  imply bounds on  $g_m$  and  $e_m$  for  $m \leq n + 1$  that permits an inductive analysis.

### 1.3. Main results

Our first theorem shows that there is a value  $z_c$  for which  $f_n$  behaves as a Gaussian. Recall that  $\sigma$  and  $\epsilon$  were defined in (1.14) and (1.17).

**Theorem 1.1.** *Let  $d > 4$ , and assume that Assumptions S, D, E and G all hold. There exist positive  $L_0 = L_0(d, \epsilon)$ ,  $z_c = z_c(d, L)$ ,  $A = A(d, L)$ , and  $v = v(d, L)$ , such that for  $L \geq L_0$ , the following statements hold.*

(a) Fix  $\gamma \in (0, 1 \wedge \frac{d-4}{4} \wedge \epsilon)$  and  $\delta \in (0, (1 \wedge \frac{d-4}{4} \wedge \epsilon) - \gamma)$ . Then

$$f_n\left(\frac{k}{\sqrt{v\sigma^2 n}}; z_c\right) = Ae^{-\frac{k^2}{2d}}[1 + \mathcal{O}(k^2 n^{-\delta}) + \mathcal{O}(\beta n^{-(d-4)/2})], \tag{1.28}$$

with the error estimate uniform in  $\{k \in \mathbb{R}^d : a(k/\sqrt{v\sigma^2 n}) \leq \gamma n^{-1} \log n\}$ .  
(b)

$$-\frac{\nabla^2 f_n(0; z_c)}{f_n(0; z_c)} = v\sigma^2 n[1 + \mathcal{O}(\beta n^{-\delta})]. \tag{1.29}$$

(c)

$$\|\hat{D}^2 f_n(\cdot; z_c)\|_1 \leq \text{const.} \beta n^{-d/2}. \tag{1.30}$$

(d) The constants  $z_c$ ,  $A$  and  $v$  obey

$$1 = \sum_{m=1}^{\infty} g_m(0; z_c), \tag{1.31}$$

$$A = \frac{1 + \sum_{m=1}^{\infty} e_m(0; z_c)}{\sum_{m=1}^{\infty} m g_m(0; z_c)}, \tag{1.32}$$

$$v = -\frac{\sum_{m=1}^{\infty} \nabla^2 g_m(0; z_c)}{\sigma^2 \sum_{m=1}^{\infty} m g_m(0; z_c)}. \tag{1.33}$$

Part (b) is an expression of diffusive behaviour. Part (c) is the first bound of (1.27), and the factor  $n^{-d/2}$  is associated with the probability of return to the origin of simple random walk after  $n$  steps. Since  $a(k)$  is asymptotic to  $\sigma^2 k^2 / (2d)$  as  $k \rightarrow 0$ , the domain of  $k$  in part (a) can be regarded as  $\{k : k^2 \leq 2d\nu\gamma \log n\}$ .

In the proof of Theorem 1.1, we will establish the bounds of (1.27) for all  $m \in \mathbb{N}$ , with  $z$  in an  $m$ -dependent interval containing  $z_c$ . Consequently, all bounds appearing in Assumptions E and G follow as a corollary, for  $z = z_c$  and all  $m \in \mathbb{N}$ .

It follows immediately from Theorem 1.1(d) and the bounds of Assumptions E and G that

$$z_c = 1 + \mathcal{O}(\beta), \quad A = 1 + \mathcal{O}(\beta), \quad v = 1 + \mathcal{O}(\beta). \quad (1.34)$$

Equation (1.31) states that at the critical value  $z_c$  of the recursion equation (1.1), with  $k = 0$ , the coefficients sum up to 1 as in renewal theory. However, unlike the standard situation in renewal theory, here the coefficients  $g_m$  depend on an additional variable  $k$ , and are not assumed to be non-negative. For the case of spread-out self-avoiding walk, the above bound for  $z_c$  improves on the error estimates of [9, 17, 21].

With modest additional assumptions, the critical point  $z_c$  can be characterised in terms of the *susceptibility*

$$\chi(z) = \sum_{n=0}^{\infty} f_n(0; z). \quad (1.35)$$

**Theorem 1.2.** *Let  $d > 4$  and assume that Assumptions S, D, E and G all hold. Let  $L$  be sufficiently large. Suppose there is a  $p_c > 0$  such that the susceptibility is absolutely convergent for  $z \in (0, p_c)$ , with  $\lim_{z \uparrow p_c} \chi(z) = \infty$ . Suppose also that the bounds of (1.27) for  $z = z_c$  and all  $m \geq 1$  imply the bounds of Assumptions E and G for all  $m \geq 2$ , uniformly in  $z \in [0, z_c]$ . Then  $z_c = p_c$ .*

Next, we state a local central limit theorem. Let

$$p_n(x; z) = \int_{[-\pi, \pi]^d} \frac{d^d k}{(2\pi)^d} e^{-ik \cdot x} f_n(k; z) \quad (1.36)$$

denote the inverse Fourier transform of  $f_n(k; z)$ . The pointwise form

$$p_n(x\sqrt{v\sigma^2 n}; z_c) \approx A \left( \frac{d}{2\pi n v \sigma^2} \right)^{d/2} e^{-dx^2/2} \quad (1.37)$$

of the local central limit theorem cannot hold for all solutions of the recursion (1.1). For example, for self-avoiding walks starting at the origin,  $p_n(0; z) = c_n(0)z^n = 0$  for every  $n \geq 1$ . This local effect will disappear only if we average over a region that grows with  $n$ . For the averaging, we denote the cube of radius  $R$  centred at  $x \in \mathbb{Z}^d$  by

$$C_R(x) = \{y \in \mathbb{Z}^d : \|x - y\|_{\infty} \leq R\}. \quad (1.38)$$

We use  $\lfloor x \rfloor$  to denote the closest lattice point in  $\mathbb{Z}^d$  to  $x \in \mathbb{R}^d$  (with an arbitrary rule to break ties).

**Theorem 1.3.** *Let  $d > 4$  and assume that Assumptions S, D, E and G all hold. Let  $R_n$  be any sequence with  $\lim_{n \rightarrow \infty} R_n = \infty$  and  $R_n = o(n^{1/2})$ . Then for all  $x \in \mathbb{R}^d$  with  $x^2[\log R_n]^{-1}$  sufficiently small,*

$$\begin{aligned} & \frac{1}{(2R_n + 1)^d} \sum_{y \in C_{R_n}(\lfloor x\sqrt{v\sigma^2 n} \rfloor)} p_n(y; z_c) \\ &= A \left( \frac{d}{2\pi n v \sigma^2} \right)^{d/2} e^{-dx^2/2} [1 + o(1)] \quad \text{as } n \rightarrow \infty. \end{aligned} \tag{1.39}$$

An explicit error bound for (1.39) in terms of  $R$  and  $n$  is given in (4.23). Note that (1.39) is not an immediate consequence of the convergence of the Fourier transform in Theorem 1.1. The statement that sums over sets of volume  $n^{d/2}$  converge to integrals of the Gaussian density over the scaled set follows from Theorem 1.1. However, since we allow arbitrarily slow growth of  $R_n$  in Theorem 1.3, we are investigating  $p_n$  on a smaller scale.

A lower bound on  $p_n(x; z_c)$  is given in the following elementary corollary to Theorem 1.1(a). In Section 1.4, we will show that the  $L$  and  $n$  dependence of the lower bound is sharp in our applications to self-avoiding walk and oriented percolation. A corresponding upper bound will be discussed in Section 1.4 in the contexts of self-avoiding walk and oriented percolation.

**Corollary 1.4.** *Let  $d > 4$  and assume that Assumptions S, D, E and G all hold. There is an  $L_0(d)$  and a  $C_1(d)$  (independent of  $L$ ) such that for  $L \geq L_0$*

$$\sup_{x \in \mathbb{Z}^d} p_n(x; z_c) \geq C_1 \sigma^{-d} n^{-d/2}. \tag{1.40}$$

*Proof.* By the  $k = 0$  case of Theorem 1.1(a), it suffices to prove (1.40) when the left side is replaced by  $p_n(x; z_c) / \sum_{y \in \mathbb{Z}^d} p_n(y; z_c)$ . Let  $X_n$  denote a random variable with probability mass function

$$\mathbb{P}(X_n = y) = \frac{p_n(yd^{-1/2}(v\sigma^2 n)^{1/2}; z_c)}{\sum_{x \in \mathbb{Z}^d} p_n(x; z_c)} \quad (y \in d^{1/2}(v\sigma^2 n)^{-1/2}\mathbb{Z}^d), \tag{1.41}$$

and let  $Z = (Z_1, \dots, Z_d)$  denote a vector of independent standard normal random variables. Then Theorem 1.1(a) asserts that  $X_n$  converges weakly to  $Z$ .

Let  $B = \{x \in \mathbb{R}^d : |x| \leq 1\}$ , let  $B_n = d^{1/2}(v\sigma^2 n)^{-1/2}\mathbb{Z}^d \cap B$ , and let  $g$  denote the density of  $Z$ . Then  $\mathbb{P}(X_n \in B) = \mathbb{P}(X_n \in B_n)$  and  $\lim_{n \rightarrow \infty} \mathbb{P}(X_n \in B_n) = \int_B g(x) d^d x$ . It follows that

$$\begin{aligned} \sup_{x \in \mathbb{Z}^d} \frac{p_n(x; z_c)}{\sum_{y \in \mathbb{Z}^d} p_n(y; z_c)} &\geq \frac{1}{|B_n|} \sum_{y \in B_n} \frac{p_n(yd^{-1/2}(v\sigma^2 n)^{1/2}; z_c)}{\sum_{x \in \mathbb{Z}^d} p_n(x; z_c)} \\ &= \frac{1}{|B_n|} \left[ \int_B g(x) d^d x + o(1) \right], \end{aligned} \tag{1.42}$$

where  $o(1)$  denotes a quantity that goes to zero as  $n \rightarrow \infty$ . The desired result then follows from the fact that  $|B_n|$  is order  $\sigma^d n^{d/2}$ , since  $v$  is order 1.  $\square$

### 1.4. Self-avoiding walks and oriented percolation

In this section, we discuss two examples where our assumptions, and hence our main results, apply. The examples are spread-out self-avoiding walks on  $\mathbb{Z}^d$ , and critical spread-out oriented percolation on  $\mathbb{Z}^d \times \mathbb{Z}_+$ , in dimensions  $d > 4$ . Both models are based on a function  $D = D_L$  obeying the assumptions of Assumption D. Assumption S is immediate for both models. We have two goals in this section. The first goal is to provide a short sketch of the lace expansion, in order to indicate how a recursion relation of the form (1.1) arises. The second goal is to give further motivation for Assumptions E and G, as well as for our main results.

#### 1.4.1. Self-avoiding walks

Recall the definition of self-avoiding walks in Section 1.1.2. If the sum in (1.7) were over  $\mathcal{R}_n(x)$ , rather than  $\mathcal{C}_n(x)$ , we would have the recursion  $c_{n+1}(x) = \sum_{y \in \mathbb{Z}^d} c_1(y)c_n(x - y)$ . However, the right side of this equation includes contributions in which the origin is visited twice, and these are not present in  $c_{n+1}(x)$ . There should therefore be a correction term  $F_{n+1}^{(1)}(x)$ , defined by the equation

$$c_{n+1}(x) = \sum_{y \in \mathbb{Z}^d} c_1(y)c_n(x - y) - F_{n+1}^{(1)}(x). \tag{1.43}$$

The lace expansion is an expansion of  $F_{n+1}^{(1)}(x)$  that treats self-avoiding walk as a small perturbation of ordinary random walk for  $d > 4$ . It was originally introduced in [6] using a graph theoretic notion of ‘‘lace.’’ Here, we follow instead the equivalent inclusion-exclusion approach of [24] (see also [17]), for which the laces do not explicitly appear.

To analyse the correction term  $F_{n+1}^{(1)}(x)$ , we define  $\mathcal{P}_{n+1}(x)$  to be the set of  $\omega \in \mathcal{R}_{n+1}(x)$  for which there exists an  $l \in \{2, \dots, n + 1\}$  (depending on  $\omega$ ) with  $\omega(l) = 0$  and  $\omega(i) \neq \omega(j)$  for all  $i \neq j$  with  $\{i, j\} \neq \{0, l\}$ . For the special case  $x = 0$ ,  $\mathcal{P}_{n+1}(0)$  is the set of  $(n + 1)$ -step self-avoiding polygons. For general  $x$ ,  $\mathcal{P}_{n+1}(x)$  is the set of self-avoiding polygons followed by a self-avoiding walk from 0 to  $x$ , with the total length being  $n + 1$  and with the walk and polygon mutually avoiding. The set  $\mathcal{P}_{n+1}(x)$  is exactly the set of walks that contribute to  $\sum_{y \in \mathbb{Z}^d} c_1(y)c_n(x - y)$  but do not contribute to  $c_{n+1}(x)$ . Therefore

$$F_{n+1}^{(1)}(x) = \sum_{\omega \in \mathcal{P}_{n+1}(x)} W(\omega). \tag{1.44}$$

Equation (1.43) can then be understood as just the inclusion-exclusion relation: The first term on the right side includes all walks from 0 to  $x$  which are self-avoiding *after* the first step, and the second subtracts the contribution due to those which are not self-avoiding from the beginning, i.e., walks that return to the origin.

The inclusion-exclusion relation can now be applied to  $F_{n+1}^{(1)}(x)$ . We neglect the mutual avoidance of the polygon and self-avoiding walk portions of an

$\omega \in \mathcal{P}_{n+1}(x)$ , and then correct by excluding the configurations which included intersections of these two portions. For  $y \in \mathbb{Z}^d$ , let

$$\pi_m^{(1)}(y) = \delta_{0,y} \sum_{\omega \in \mathcal{P}_m(0)} W(\omega). \tag{1.45}$$

Then we define  $F_{n+1}^{(2)}(x)$  by

$$F_{n+1}^{(1)}(x) = \sum_{y \in \mathbb{Z}^d} \sum_{m=2}^{n+1} \pi_m^{(1)}(y) c_{n+1-m}(x-y) - F_{n+1}^{(2)}(x). \tag{1.46}$$

The correction term  $F_{n+1}^{(2)}(x)$  involves configurations consisting of a self-avoiding polygon and a self-avoiding walk from 0 to  $x$ , of total length  $n + 1$ , and with an intersection required between the self-avoiding polygon and the self-avoiding walk (in addition to their intersection at the origin). If we define  $\Theta_{n+1}(x)$  to be the corresponding subset of  $\mathcal{R}_{n+1}(x)$ , then we have

$$F_{n+1}^{(2)}(x) = \sum_{\omega \in \Theta_{n+1}(x)} W(\omega). \tag{1.47}$$

The walk and polygon may intersect more than once, and we focus on the first time an intersection occurs, measuring time according to the walk. We then perform inclusion-exclusion again, neglecting the avoidance between the portions of the self-avoiding walk before and after this first intersection, and then subtracting a correction term. The process is continued indefinitely. Defining  $\pi_m^{(2)}(y)$  as in (1.24), we are led to

$$c_{n+1}(x) = \sum_{y \in \mathbb{Z}^d} c_1(y) c_n(x-y) + \sum_{y \in \mathbb{Z}^d} \sum_{m=2}^{n+1} \pi_m(y) c_{n+1-m}(x-y). \tag{1.48}$$

Here

$$\pi_m(y) = \sum_{N=1}^{\infty} (-1)^N \pi_m^{(N)}(y), \tag{1.49}$$

with expressions for the  $\pi^{(N)}$  for  $N \geq 3$  that are given in [6, 17].

By definition,  $c_1(x) = D(x)$ . Using this in the first term of (1.48), we then multiply through by  $z^{n+1}$ , with  $z \geq 0$ , and take the Fourier transform to obtain

$$\hat{c}_{n+1}(k) z^{n+1} = z \hat{D}(k) \hat{c}_n(k) z^n + \sum_{m=2}^{n+1} \hat{\pi}_m(k) z^m \hat{c}_{n+1-m}(k) z^{n+1-m}. \tag{1.50}$$

As was noted already in Section 1.1.2, (1.50) is a special case of (1.1), with  $e_n = 0$  and

$$\begin{aligned} f_n(k; z) &= \hat{c}_n(k) z^n \quad (n \geq 0), & g_1(k; z) &= z \hat{D}(k), \\ g_n(k; z) &= \hat{\pi}_n(k) z^n \quad (n \geq 2). \end{aligned} \tag{1.51}$$

Assumption S is a consequence of the lattice symmetry of self-avoiding walks and the fact that  $\hat{c}_n(k) \leq 1$ . We are already assuming that  $D$  obeys the requirements of Assumption D, so Assumption D is immediate from (1.51). Assumption E is vacuous, and the remaining assumption, Assumption G, is the substantial one.

An analysis corresponding to the verification of some version of Assumption G is present in one form or another in all lace expansion analyses [5–7, 9, 11, 13, 16, 17, 23]. In [15], we prove that Assumption G is satisfied, as part of a larger analysis proving Gaussian behaviour for networks of mutually-avoiding spread-out self-avoiding walks in dimensions  $d > 4$ .

As a consequence of the verification of Assumption G in [15], Theorems 1.1 and 1.3 hold for  $f_n(k; z_c) = \hat{c}_n(k)z_c^n$ , with  $z_c$ ,  $A$  and  $D$  given by Equations (1.31)–(1.33). Theorem 1.1(a,b) were proved in [17] with weaker error bounds, via a different approach using generating functions. Theorem 1.3 is the first time a local central limit theorem has been proved for the self-avoiding walk. (As this work was completed, an alternate approach to the local central limit theorem appeared in [5].) The limit  $\lim_{n \rightarrow 0} \hat{c}_n(0)^{1/n}$  exists by a well-known subadditivity argument [17]. Thus the critical value  $z_c$  appearing in Theorem 1.1(a) must be given by  $z_c = \lim_{n \rightarrow 0} \hat{c}_n(0)^{-1/n}$ , consistent with the conclusion of Theorem 1.2.

The first bound assumed in (1.27), which asserts that  $\|\hat{D}^2 \hat{c}_m\|_1 \leq \mathcal{O}(\beta z_c^{-m} m^{-d/2})$ , actually implies a bound on  $\|c_m\|_\infty$ . In fact, if  $m \geq 2$ , by ignoring the self-avoidance constraint for the first two steps we have

$$\|c_m\|_\infty \leq \sup_{x \in \mathbb{Z}^d} \sum_{u, v \in \mathbb{Z}^d} D(u)D(v-u)c_{m-2}(x-v) \leq \|\hat{D}^2 \hat{c}_{m-2}\|_1. \tag{1.52}$$

Since  $z_c^{-1} \leq \mathcal{O}(1)$  by (1.34), the first bound of (1.27) implies that

$$\|c_m\|_\infty \leq \mathcal{O}(\beta z_c^{-m} m^{-d/2}). \tag{1.53}$$

The above argument gives (1.53) when  $m \geq 2$ , but (1.53) clearly also holds for  $m = 1$  since  $\|c_1\|_\infty = \|D\|_\infty \leq \mathcal{O}(\beta)$  and  $z_c^{-1}$  is bounded away from zero. The inequality (1.53) was useful to estimate (1.25). After having verified Assumption G, the bound  $\|\hat{D}^2 \hat{c}_m\|_1 \leq \mathcal{O}(\beta z_c^{-m} m^{-d/2})$  then follows from Theorem 1.1(c). We conclude that (1.53), which is of independent interest, does in fact hold for all  $m$ . A completely different proof of this bound, using generating functions and a finite memory cut-off, is given in [17] for the uniform  $D$ . With Corollary 1.4, (1.53) gives

$$C_1 \beta z_c^{-m} m^{-d/2} \leq \|c_m\|_\infty \leq C_2 \beta z_c^{-m} m^{-d/2} \tag{1.54}$$

for  $d > 4$  and  $L$  sufficiently large.

### 1.4.2. Oriented percolation

The lace expansion for oriented percolation is an expansion for the two-point function (1.11). It was derived specifically in the context of oriented percolation by Nguyen and Yang in [19, 20], using laces as in [6]. An expansion for ordinary percolation based on inclusion-exclusion was derived in [10], which applies also

to oriented percolation and which is utilised in [14]. Here, we briefly outline a derivation of the Nguyen–Yang expansion using inclusion-exclusion. This approach, which is different than the inclusion-exclusion approach of [10], is developed also in [22].

Given a bond configuration, we define a bond to be *pivotal* for  $(x, n) \rightarrow (y, m)$  if  $(x, n) \rightarrow (y, m)$  when this bond is made occupied, whereas  $(x, n) \not\rightarrow (y, m)$  when this bond is made vacant. We say  $(x, n)$  is *doubly connected* to  $(y, m)$ , denoted  $(x, n) \Rightarrow (y, m)$  when  $x \rightarrow y$  but there is no pivotal bond for the connection from  $(x, n)$  to  $(y, m)$ . We write  $(x, n) \not\Rightarrow (y, m)$  for the complement of  $(x, n) \Rightarrow (y, m)$ . Given a configuration in which  $(0, 0) \rightarrow (x, n)$ , either  $(0, 0) \Rightarrow (x, n)$  or  $(0, 0) \not\Rightarrow (x, n)$ . For  $n \geq 0$ , let

$$\rho_n^{(0)}(x; z) = \mathbb{P}_z((0, 0) \Rightarrow (x, n)), \tag{1.55}$$

$$\sigma_n^{(0)}(x; z) = \mathbb{P}_z(\{(0, 0) \rightarrow (x, n)\} \cap \{(0, 0) \not\Rightarrow (x, n)\}). \tag{1.56}$$

Then

$$\tau_{n+1}(x; z) = \rho_{n+1}^{(0)}(x; z) + \sigma_{n+1}^{(0)}(x; z) \quad (n \geq 0). \tag{1.57}$$

A configuration contributing to  $\sigma_{n+1}^{(0)}(x; z)$  contains at least one pivotal bond, and hence a first pivotal bond  $((u, m), (v, m + 1))$ . Let  $E$  denote the event that  $(0, 0) \Rightarrow (u, m)$ , that  $((u, m), (v, m + 1))$  is occupied, and that  $(v, m + 1) \rightarrow (x, n + 1)$ . Let  $F$  be the event that the bond  $((u, m), (v, m + 1))$  is pivotal for  $(0, 0) \rightarrow (x, n + 1)$ . Then

$$\sigma_{n+1}^{(0)}(x; z) = \sum_{u,v} \sum_{m=0}^n \mathbb{P}_z(E \cap F). \tag{1.58}$$

We use inclusion-exclusion to write  $\mathbb{P}_z(E \cap F) = \mathbb{P}_z(E) - \mathbb{P}_z(E \cap F^c)$ . Oriented percolation enjoys a Markov property (unlike non-oriented percolation) which implies that

$$\mathbb{P}_z(E) = \rho_m^{(0)}(u; z) zD(v - u) \tau_{n-m}(x - v; z). \tag{1.59}$$

Thus we have

$$\begin{aligned} \tau_{n+1}(x; z) &= \rho_{n+1}^{(0)}(x; z) + \sum_{u,v} \sum_{m=0}^n \rho_m^{(0)}(u; z) zD(v - u) \tau_{n-m}(x - v; z) \\ &\quad - \sigma_{n+1}^{(1)}(x; z), \end{aligned} \tag{1.60}$$

where

$$\sigma_{n+1}^{(1)}(x; z) = \sum_{u,v} \sum_{m=0}^n \mathbb{P}_z(E \cap F^c). \tag{1.61}$$

Further applications of inclusion-exclusion require more detail than we can enter into here. Details can be found in [22]. The result is as follows. Since a

site is always doubly connected to itself, we have  $\rho_0^{(0)}(x; z) = \delta_{0,x}$ . We define  $\pi_m^{(0)}(x; z) = \rho_m^{(0)}(x; z)$  for  $m \geq 2$ , and  $\pi_m^{(0)}(x; z) = 0$  for  $m = 0, 1$ . An appropriate repeated application of inclusion-exclusion then leads, after taking the Fourier transform, to

$$\hat{\tau}_{n+1}(k; z) = z\hat{D}(k)\tau_n(k; z) + z\hat{D}(k) \sum_{m=2}^n \hat{\pi}_m(k; z)\hat{\tau}_{n-m}(k; z) + \hat{\pi}_{n+1}(k; z) \quad (n \geq 0). \tag{1.62}$$

The function  $\pi_m(x; z)$  is given by an explicit alternating series [14, 19, 22]

$$\pi_m(x; z) = \sum_{N=0}^{\infty} (-1)^N \pi_m^{(N)}(x; z). \tag{1.63}$$

Equation (1.62) is a special case of (1.1) with the choices

$$f_n(k; z) = \hat{\tau}_n(k; z), \quad e_n(k; z) = \hat{\pi}_n(k; z), \tag{1.64}$$

and

$$g_1(k; z) = z\hat{D}(k), \quad g_2(k; z) = 0, \quad g_m(k; z) = z\hat{D}(k)\hat{\pi}_{m-1}(k; z) \quad (m \geq 3). \tag{1.65}$$

For oriented percolation, Assumption S follows from the lattice symmetry of the model and the fact that  $|\hat{\tau}_n(k; z)| \leq \hat{\tau}_n(0; z) \leq \hat{\tau}_1(0; z)^n = z^n$ . Assumption D follows from (1.64)–(1.65) and the fact that  $\hat{\pi}_1(k; z) = 0$ . Assumptions E and G are very closely related. They can be verified, for  $d > 4$  and  $L$  sufficiently large, by using the BK inequality to bound  $\pi^{(N)}$  in terms of the two-point function itself. For example, the BK inequality implies that  $\pi_m^{(0)}(x; z) \leq \tau_m(x; z)^2$  ( $m \geq 2$ ). This is carried out in detail in [14], where the verification of the assumptions is an essential part of a larger analysis relating the scaling limit of critical oriented percolation for  $d > 4$  to super-Brownian motion.

Given the result of [14] that Assumptions E and G do hold, the conclusions of Theorems 1.1 and 1.3 therefore apply at  $z = z_c$ . Theorems 1.1(a,b) were proved in [20], with weaker error bounds. Theorems 1.1(c) and 1.3 are new, and the former plays an essential role in the analysis of [14]. As was the case for self-avoiding walks in (1.53), it is possible to conclude from the assumed bounds (1.27) that  $\|\tau_n(\cdot; z_c)\|_{\infty} \leq CK\beta n^{-d/2}$ . See [14] for details. With Corollary 1.4, this gives

$$C_1\beta n^{-d/2} \leq \|\tau_n(\cdot; z_c)\|_{\infty} \leq C_2\beta n^{-d/2} \tag{1.66}$$

for  $d > 4$  and  $L$  sufficiently large.

The hypotheses of Theorem 1.2 concerning the susceptibility are well-known [2], and the hypothesis concerning Assumptions E and G is established in [14]. It therefore follows that  $z_c$  corresponds to the critical oriented percolation threshold. This can be understood directly from the conclusion of Theorem 1.1(a), which asserts in particular that  $\lim_{n \rightarrow \infty} \hat{\tau}_n(0; z_c) = A$ , with  $A$  positive and finite. The quantity  $\hat{\tau}_n(0; z)$  represents the expected number of sites in the intersection of the

connected cluster of the origin with  $\mathbb{Z}^d \times \{n\}$ . This goes to zero for  $z$  below the percolation threshold, by exponential decay of connectivities [1, 2, 18]. On the other hand, for  $z$  above the percolation threshold, it can be expected that  $\hat{\tau}_n(0; z) \rightarrow \infty$ . This latter statement is a consequence of the shape theorem, which has been recently extended from the nearest-neighbour model results of [4] to more general long-range models [8].

The triangle condition [2, 3] is a diagrammatic sufficient condition for the existence of several critical exponents. This is discussed in detail in [19], where the triangle condition is verified, in particular, for sufficiently spread-out models of oriented percolation when  $d + 1 > 5$ . The upper bound of (1.66), together with the bound  $\sum_x \tau_n(x; z_c) < \infty$  of Theorem 1.1(a), can be combined to give an alternate proof of the triangle condition. This is discussed in detail in [14].

### 1.5. Organisation

The remainder of this paper is organised as follows. In Section 2, we introduce the induction hypotheses on  $f_n$  that will be used to prove our main results, and derive some consequences of the induction hypotheses. The induction is advanced in Section 3. In Section 4, the main results stated in Section 1.3 are proved. We conclude in Appendix A, by showing that the function  $D$  defined in (1.22) obeys the properties listed in Assumption D.

## 2. Induction hypotheses

We will analyse the recursion relation (1.1) using induction on  $n$ . Our method is a generalisation of the inductive method introduced by den Hollander and the authors [13] to study weakly self-avoiding walk for  $d > 4$ . Here, we are considering long-range or “spread-out” models. This necessitates a number of modifications to the analysis of [13], as the small parameter is less explicit for the spread-out model. In addition, the analysis of [13] has been simplified in some respects. Our choice of induction hypotheses has been strongly motivated by those used in [13], which in turn were motivated by earlier work. The inductive approach leads to more detailed results than have been obtained previously for self-avoiding walk or oriented percolation using generating functions. In a different vein, it has also been adapted to prove ballistic behaviour for weakly self-avoiding walk for  $d = 1$  [12].

In this section, we introduce the induction hypotheses, verify that they hold for  $n = 1$ , discuss their motivation, and derive some of their consequences.

### 2.1. Statement of induction hypotheses (H1–H4)

The induction hypotheses involve a sequence  $v_n$ , which is defined as follows. We set  $v_0 = b_0 = 1$ , and for  $n \geq 1$  we define

$$b_n = -\frac{1}{\sigma^2} \sum_{m=1}^n \nabla^2 g_m(0; z), \quad c_n = \sum_{m=1}^n (m-1)g_m(0; z), \quad v_n = \frac{b_n}{1 + c_n}. \tag{2.1}$$

The  $z$ -dependence of  $b_n, c_n, v_n$  will usually be left implicit in the notation. We will often simplify the notation by dropping  $z$  also from  $e_n, f_n$  and  $g_n$ , and write, e.g.,  $f_n(k) = f_n(k; z)$ .

As we will explain in more detail in Section 2.3, the diffusion constant  $\sigma^2 v$  of Theorem 1.1 will turn out to be given by  $\sigma^2 v_\infty(z_c)$ . However, at this stage we have not yet proved that the series in the definition of  $v_\infty$  converge. Neither have we yet identified  $z_c$ , other than as a solution to (1.31), which involves a series whose convergence has not yet been established.

The induction hypotheses also involve several constants. Let  $d > 4$ , and recall that  $\epsilon$  was specified in (1.17). We fix  $\gamma, \delta, \rho > 0$  according to

$$0 < \frac{d-4}{2} - \rho < \gamma < \gamma + \delta < 1 \wedge \frac{d-4}{2} \wedge \epsilon. \tag{2.2}$$

This can be done by first fixing  $\gamma \in (0, 1 \wedge \frac{d-4}{2} \wedge \epsilon)$  and then choosing  $\delta$  and  $\rho$  accordingly. We also introduce constants  $K_1, \dots, K_5$ , which are independent of  $\beta$ . We define

$$K'_4 = \max\{C_e(cK_4), C_g(cK_4), K_4\}, \tag{2.3}$$

where  $c$  is a constant determined in Lemma 2.5 below. To advance the induction, we will need to assume that

$$K_3 \gg K_1 > K'_4 \geq K_4 \gg 1, \quad K_2 \geq K_1, 3K'_4, \quad K_5 \gg K_4. \tag{2.4}$$

Here  $a \gg b$  denotes the statement that  $a/b$  is sufficiently large. The amount by which, for instance,  $K_3$  must exceed  $K_1$  is independent of  $\beta$  and will be determined during the course of the advancement of the induction in Section 3.

Let  $z_0 = z_1 = 1$ , and define  $z_n$  recursively by

$$z_{n+1} = 1 - \sum_{m=2}^{n+1} g_m(0; z_n), \quad n \geq 1. \tag{2.5}$$

For  $n \geq 1$ , we define intervals

$$I_n = [z_n - K_1 \beta n^{-(d-2)/2}, z_n + K_1 \beta n^{-(d-2)/2}]. \tag{2.6}$$

Recall the definition  $a(k) = 1 - \hat{D}(k)$  from (1.15). Our induction hypotheses are that the following four statements hold for all  $z \in I_n$  and all  $1 \leq j \leq n$ .

- (H1)  $|z_j - z_{j-1}| \leq K_1 \beta j^{-d/2}$ .
- (H2)  $|v_j - v_{j-1}| \leq K_2 \beta j^{-(d-2)/2}$ .
- (H3) For  $k$  such that  $a(k) \leq \gamma j^{-1} \log j$ ,  $f_j(k; z)$  can be written in the form

$$f_j(k; z) = \prod_{i=1}^j [1 - v_i a(k) + r_i(k)],$$

with  $r_i(k) = r_i(k; z)$  obeying

$$|r_i(0)| \leq K_3 \beta i^{-(d-2)/2}, \quad |r_i(k) - r_i(0)| \leq K_3 \beta a(k) i^{-\delta}.$$

(H4) For  $k$  such that  $a(k) > \gamma j^{-1} \log j$ ,  $f_j(k; z)$  obeys the bounds

$$|f_j(k; z)| \leq K_4 a(k)^{-2-\rho} j^{-d/2}, \quad |f_j(k; z) - f_{j-1}(k; z)| \leq K_5 a(k)^{-1-\rho} j^{-d/2}.$$

Note that, for  $k = 0$ , (H3) reduces to  $f_j(0) = \prod_{i=1}^j [1 + r_i(0)]$ .

### 2.2. Initialisation of the induction

We now verify that the induction hypotheses hold when  $n = 1$ . Fix  $z \in I_1$ .

(H1) We simply have  $z_1 - z_0 = 1 - 1 = 0$ .

(H2) We simply have  $|v_1 - v_0| = |z - 1|$ , so that (H2) is satisfied provided  $K_2 \geq K_1$ .

(H3) We are restricted to  $a(k) = 0$ . By (1.19), this means  $k = 0$ . By Assumption D,  $f_1(0; z) = z$ , so that  $r_1(0) = z - z_1$ . Thus (H3) holds provided we take  $K_3 \geq K_1$ .

(H4) We note that  $|f_1(k; z)| \leq z \leq 2$  for  $\beta$  sufficiently small,  $|f_1(k; z) - f_0(k; z)| \leq 3$ , and  $a(k) \leq 2$ . The bounds of (H4) therefore hold provided we take  $K_4 \geq 2^{3+\rho}$  and  $K_5 \geq 3 \cdot 2^{1+\rho}$ .

### 2.3. Discussion of induction hypotheses

**(H1) and the critical point.** The critical point can be formally identified as follows. We set  $k = 0$  in (1.1), then sum over  $n$ , and solve for the susceptibility  $\chi(z)$  of (1.35). The result is

$$\chi(z) = \frac{1 + \sum_{m=2}^{\infty} e_m(0; z)}{1 - \sum_{m=1}^{\infty} g_m(0; z)}. \tag{2.7}$$

The critical point should correspond to the smallest zero of the denominator and hence should obey the equation

$$1 - \sum_{m=1}^{\infty} g_m(0; z_c) = 1 - z_c - \sum_{m=2}^{\infty} g_m(0; z_c) = 0. \tag{2.8}$$

However, we do not know *a priori* that the series in (2.7) or (2.8) converge. We therefore approximate (2.8) with the recursion (2.5), which bypasses the convergence issue by discarding the  $g_m(0)$  for  $m > n + 1$  that cannot be handled at the  $n^{\text{th}}$  stage of the induction argument. The sequence  $z_n$  will ultimately converge to  $z_c$ . Equation (2.8) is identical to (1.31).

In dealing with the sequence  $z_n$ , it is convenient to formulate the induction hypotheses for a small interval  $I_n$  approximating  $z_c$ . As we will see in Section 2.4, (H1) guarantees that the intervals  $I_j$  are decreasing:  $I_1 \supset I_2 \supset \dots \supset I_n$ . Because the length of these intervals is shrinking to zero, their intersection  $\bigcap_{j=1}^{\infty} I_j$  is a single point, namely  $z_c$ . Hypothesis (H1) drives the convergence of  $z_n$  to  $z_c$  and gives some control on the rate. The rate is determined from (2.5) and the ansatz that the difference  $z_j - z_{j-1}$  is approximately  $-g_{j+1}(0, z_c)$ , with  $|g_j(k; z_c)| = \mathcal{O}(\beta j^{-d/2})$  as in Assumption G.

**(H2).** The formula for  $v_n$  in (2.1) can be motivated by the following rough argument. Differentiating (1.1) twice with respect to  $k$ , setting  $k = 0$ , and using the fact that odd derivatives vanish by Assumption S, we obtain

$$\nabla^2 f_{n+1}(0) = \sum_{m=1}^{n+1} \left[ g_m(0) \nabla^2 f_{n+1-m}(0) + \nabla^2 g_m(0) f_{n+1-m}(0) \right] + \nabla^2 e_{n+1}(0). \quad (2.9)$$

We will use ‘ $\approx$ ’ to denote an uncontrolled approximation in a rough argument. In (2.9), we make the approximations  $f_{n+1-m}(0) \approx f_n(0)$  and  $\nabla^2 e_{n+1} \approx 0$ , subtract  $\nabla^2 f_n(0)$  from both sides, and recall the definition of  $b_n$  from (2.1), to obtain

$$\nabla^2 f_{n+1}(0) - \nabla^2 f_n(0) \approx -\sigma^2 b_{n+1} f_n(0) + \sum_{m=1}^{n+1} g_m(0) \nabla^2 f_{n+1-m}(0) - \nabla^2 f_n(0). \quad (2.10)$$

In view of (2.5), we insert the approximation  $1 \approx \sum_{m=1}^{n+1} g_m(0)$  to obtain

$$\begin{aligned} \nabla^2 f_{n+1}(0) - \nabla^2 f_n(0) &\approx -\sigma^2 b_{n+1} f_n(0) \\ &+ \sum_{m=1}^{n+1} g_m(0) \left[ \nabla^2 f_{n+1-m}(0) - \nabla^2 f_n(0) \right]. \end{aligned} \quad (2.11)$$

On the other hand, applying  $\nabla^2$  to (H3) with  $j = n$  and then using  $r_n(0) \approx 0$  and  $\nabla^2 r_n(0) \approx 0$  gives

$$\nabla^2 f_n(0) - \nabla^2 f_{n-1}(0) \approx -\sigma^2 v_n f_{n-1}(0). \quad (2.12)$$

In view of this, we assume  $\nabla^2 f_{n+1-m}(0) - \nabla^2 f_n(0) \approx (m-1)\sigma^2 v_{n+1} f_n(0)$ . Recalling the definition of  $c_{n+1}$  from (2.1), (2.11) then gives

$$\nabla^2 f_{n+1}(0) - \nabla^2 f_n(0) \approx -\sigma^2 (b_{n+1} - v_{n+1} c_{n+1}) f_n(0). \quad (2.13)$$

Putting the right-hand side equal to  $-\sigma^2 v_{n+1} f_n(0)$ , as in (2.12), leads to the formula for  $v_{n+1}$  of (2.1).

The assumed bound on  $|v_j - v_{j-1}|$  can then be guessed by writing  $v_j$  and  $v_{j-1}$  in terms of the  $b_i$  and  $c_i$ , and assuming the bounds of Assumption G to estimate the resulting expression. The calculations will be carried out in detail when we advance (H2) in Section 3.2.

**(H3).** The bound on  $r_i(0)$  ensures that the limit  $A = \lim_{j \rightarrow \infty} f_j(0; z_c) = \prod_{i=1}^{\infty} [1 + r_i(0; z_c)]$  exists. The expression for  $f_j$  in (H3) can be approximately rewritten as

$$f_j(k) \approx f_j(0) \exp \left[ \sum_{i=1}^j \left( -v_i a(k) + r_i(k) - r_i(0) \right) \right].$$

The bound on  $r_i(k) - r_i(0)$  indicates that it is a small perturbation of the leading term  $-v_i a(k)$ . The limit  $v = \lim_{n \rightarrow \infty} v_n$  exists, if we assume the bounds on  $g_m$  of

Assumption G. Also,  $a(k) \sim \sigma^2 k^2 / 2d$  as  $k \rightarrow 0$ . Thus we can understand (H3) as a precise version of the approximation

$$f_j(k) \approx f_j(0) \exp \left[ - \frac{v \sigma^2 k^2 j}{2d} \right], \tag{2.14}$$

which we expect to be valid at least for  $\sigma^2 k^2 j$  of order 1. This is consistent with Theorem 1.1. For large  $j$ , the restriction  $a(k) \leq \gamma j^{-1} \log j$  is essentially the restriction  $\sigma^2 k^2 j \leq 2d\gamma \log j$ , which includes the region where  $\sigma^2 k^2 j$  is order 1, plus some additional room to manoeuvre.

**(H4).** For  $\sigma^2 k^2 j > 2d\gamma \log j$ , we require (and can prove) less accurate control of  $f_j(k)$ , as expressed in (H4). The form of (H4) has been chosen in part to be less stringent than (H3) for  $a(k) = \gamma j^{-1} \log j$ , where the transition from (H3) to (H4) takes place. In fact, inserting  $\sigma^2 k^2 / 2d = \gamma j^{-1} \log j$  into (2.14) gives an expression which grows like  $j^{-\gamma v}$ . The bounds on  $g_m$  of Assumption G would imply that  $z_c = 1 + \mathcal{O}(\beta)$  and  $v = 1 + \mathcal{O}(\beta)$ , so that  $j^{-\gamma v} = j^{-\gamma(1+\mathcal{O}(\beta))}$ . On the other hand, putting  $a(k) = \gamma j^{-1} \log j$  in the first bound of (H4) gives a bound which grows like  $j^{-(d-4)/2+\rho}$  times a power of a logarithm. By (2.2), this is a weaker bound than the (H3) bound.

#### 2.4. Consequences of induction hypotheses

In this section we derive important consequences of the induction hypotheses. The key result is that the induction hypotheses imply (1.27) for all  $1 \leq m \leq n$ , from which the bounds of Assumptions E and G then follow, for  $2 \leq m \leq n + 1$ .

Here, and throughout the rest of this paper:

- $C$  denotes a strictly positive constant that may depend on  $d, \gamma, \delta, \rho$ , but *not* on the  $K_i$ , *not* on  $k$ , *not* on  $n$ , and *not* on  $\beta$  (provided  $\beta$  is sufficiently small, possibly depending on the  $K_i$ ). The value of  $C$  may change from line to line.
- We frequently assume  $\beta \ll 1$  without explicit comment.

The first lemma shows that the intervals  $I_j$  are nested, assuming (H1).

**Lemma 2.1.** *Assume (H1) for  $1 \leq j \leq n$ . Then  $I_1 \supset I_2 \supset \dots \supset I_n$ .*

*Proof.* Suppose  $z \in I_j$ , with  $2 \leq j \leq n$ . Then by (H1) and (2.6),

$$|z - z_{j-1}| \leq |z - z_j| + |z_j - z_{j-1}| \leq \frac{K_1 \beta}{j^{(d-2)/2}} + \frac{K_1 \beta}{j^{d/2}} \leq \frac{K_1 \beta}{(j-1)^{(d-2)/2}}, \tag{2.15}$$

and hence  $z \in I_{j-1}$ . □

By Lemma 2.1, if  $z \in I_j$  for  $1 \leq j \leq n$ , then  $z \in I_1$  and hence, by (2.6),

$$|z - 1| \leq K_1 \beta. \tag{2.16}$$

It also follows from (H2) that, for  $z \in I_n$  and  $1 \leq j \leq n$ ,

$$|v_j - 1| \leq CK_2\beta. \tag{2.17}$$

The induction hypothesis (H3) has the useful alternate form

$$f_j(k) = f_j(0) \prod_{i=1}^j [1 - v_i a(k) + s_i(k)], \tag{2.18}$$

with  $s_i(0) = 0$  and

$$|s_i(k)| \leq K_3(1 + C(K_2 + K_3)\beta)\beta a(k)i^{-\delta}. \tag{2.19}$$

In fact, (2.18) is an identity if we define

$$s_i(k) = [1 + r_i(0)]^{-1}[v_i a(k)r_i(0) + (r_i(k) - r_i(0))], \tag{2.20}$$

and (2.19) then follows from (2.17) and the bounds on  $r_i$  of (H3).

The next lemma provides an important upper bound on  $f_j(k; z)$ , for  $k$  small depending on  $j$ , as in (H3).

**Lemma 2.2.** *Let  $z \in I_n$  and assume (H2–H3) for  $1 \leq j \leq n$ . Then for  $k$  with  $a(k) \leq \gamma j^{-1} \log j$ ,*

$$|f_j(k; z)| \leq e^{CK_3\beta} e^{-(1-C(K_2+K_3)\beta)ja(k)}. \tag{2.21}$$

*Proof.* We use (2.18), and conclude from the bound on  $r_i(0)$  of (H3) that  $|f_j(0)| = \prod_{i=1}^j |1 + r_i(0)| \leq e^{CK_3\beta}$ , using  $1 + x \leq e^x$  for each factor. Then we use (2.17) and (2.19) to obtain

$$\prod_{i=1}^j |1 - v_i a(k) + s_i(k)| \leq \prod_{i=1}^j |1 - (1 - CK_2\beta)a(k) + CK_3\beta a(k)i^{-\delta}|. \tag{2.22}$$

The desired bound then follows, again using  $1 + x \leq e^x$  for each factor on the right side. □

The middle bound of (1.27) follows, for  $1 \leq m \leq n$  and  $z \in I_m$ , directly from Lemma 2.2. We next prove two lemmas which provide the other two bounds of (1.27). This will supply the hypothesis (1.27) for Assumptions E and G, and therefore plays a crucial role in advancing the induction.

**Lemma 2.3.** *Let  $z \in I_n$  and assume (H2), (H3) and (H4). Then for  $1 \leq j \leq n$ ,*

$$\|\hat{D}^2 f_j(\cdot; z)\|_1 \leq C(1 + K_4)\beta j^{-d/2}. \tag{2.23}$$

*Proof.* Fix  $z \in I_n$  and  $1 \leq j \leq n$ , and define

$$\begin{aligned} R_1 &= \{k \in [-\pi, \pi]^d : a(k) \leq \gamma j^{-1} \log j, \|k\|_\infty \leq L^{-1}\}, \\ R_2 &= \{k \in [-\pi, \pi]^d : a(k) \leq \gamma j^{-1} \log j, \|k\|_\infty > L^{-1}\}, \\ R_3 &= \{k \in [-\pi, \pi]^d : a(k) > \gamma j^{-1} \log j, \|k\|_\infty \leq L^{-1}\}, \\ R_4 &= \{k \in [-\pi, \pi]^d : a(k) > \gamma j^{-1} \log j, \|k\|_\infty > L^{-1}\}. \end{aligned} \tag{2.24}$$

The set  $R_2$  is empty if  $j$  is sufficiently large. Then

$$\|\hat{D}^2 f_j\|_1 = \sum_{i=1}^4 \int_{R_i} \hat{D}(k)^2 |f_j(k)| \frac{d^d k}{(2\pi)^d}. \tag{2.25}$$

We will treat each of the four terms on the right side separately.

On  $R_1$ , we use (1.19) in conjunction with Lemma 2.2 and the fact that  $\hat{D}^2 \leq 1$ , to obtain

$$\int_{R_1} \hat{D}(k)^2 |f_j(k)| \frac{d^d k}{(2\pi)^d} \leq \int_{R_1} C e^{-cj(Lk)^2} \frac{d^d k}{(2\pi)^d} \leq \frac{C}{L^d j^{d/2}}. \tag{2.26}$$

On  $R_2$ , we use Lemma 2.2 and (1.20) to conclude that there is an  $\alpha > 1$  such that

$$\int_{R_2} \hat{D}(k)^2 |f_j(k)| \frac{d^d k}{(2\pi)^d} \leq \int_{R_2} \alpha^{-j} \frac{d^d k}{(2\pi)^d} = \alpha^{-j} |R_2|, \tag{2.27}$$

where  $|R_2|$  denotes the volume of  $R_2$ . This volume is maximal when  $j = 3$ , so that

$$\begin{aligned} |R_2| &\leq |\{k : a(k) \leq 1 - \frac{\gamma \log 3}{3}\}| \leq |\{k : \hat{D}(k) \geq \frac{\gamma \log 3}{3}\}| \\ &\leq (\frac{3}{\gamma \log 3})^2 \|\hat{D}^2\|_1 \leq (\frac{3}{\gamma \log 3})^2 \beta, \end{aligned} \tag{2.28}$$

using (1.18) in the last step. Therefore  $\alpha^{-j} |R_2| \leq C\beta j^{-d/2}$  and

$$\int_{R_2} \hat{D}(k)^2 |f_j(k)| \frac{d^d k}{(2\pi)^d} \leq C\beta j^{-d/2}. \tag{2.29}$$

On  $R_3$  and  $R_4$ , we use (H4). As a result, the contribution from these two regions is bounded above by

$$\frac{K_4}{j^{d/2}} \sum_{i=3}^4 \int_{R_i} \frac{\hat{D}(k)^2}{a(k)^{2+\rho}} \frac{d^d k}{(2\pi)^d}. \tag{2.30}$$

On  $R_3$ , we use  $\hat{D}(k)^2 \leq 1$  and (1.19) to obtain the upper bound

$$\frac{CK_4}{j^{d/2} L^{4+2\rho}} \int_{\|k\|_\infty < L^{-1}} \frac{1}{k^{4+2\rho}} d^d k \leq \frac{CK_4}{j^{d/2} L^{4+2\rho}} \left(\frac{1}{L}\right)^{d-4-2\rho} = \frac{CK_4\beta}{j^{d/2}}, \tag{2.31}$$

where the integral is finite since  $\rho < \frac{d-4}{2}$  by (2.2). On  $R_4$ , we use (1.18) and (1.20) to obtain the bound

$$\frac{CK_4}{j^{d/2}} \int_{[-\pi, \pi]^d} \hat{D}(k)^2 \frac{d^d k}{(2\pi)^d} \leq \frac{CK_4\beta}{j^{d/2}}. \tag{2.32}$$

This completes the proof. □

**Lemma 2.4.** *Let  $z \in I_n$  and assume (H2) and (H3). Then, for  $1 \leq j \leq n$ ,*

$$|\nabla^2 f_j(0; z)| \leq (1 + C(K_2 + K_3)\beta)\sigma^2 j. \tag{2.33}$$

*Proof.* Fix  $z \in I_n$  and  $j$  with  $1 \leq j \leq n$ . By (2.18) and Assumption S,

$$\nabla^2 f_j(0) = f_j(0) \sum_{i=1}^j [-\sigma^2 v_i + \nabla^2 s_i(0)]. \tag{2.34}$$

By (2.17),  $|v_i - 1| \leq CK_2\beta$ . For the second term on the right side, we let  $e_1, \dots, e_d$  denote the standard basis vectors in  $\mathbb{R}^d$ . By (2.19), for all  $i \leq n$  we have

$$\begin{aligned} |\nabla^2 s_i(0)| &= \left| \sum_{l=1}^d \lim_{t \rightarrow 0} \frac{s_i(te_l) - s_i(0)}{t^2} \right| \\ &\leq CK_3\beta i^{-\delta} \sum_{l=1}^d \lim_{t \rightarrow 0} \frac{a(te_l)}{t^2} = CK_3\sigma^2\beta i^{-\delta}. \end{aligned} \tag{2.35}$$

Therefore, by Lemma 2.2,

$$|\nabla^2 f_j(0)| \leq e^{CK_3\beta} \sigma^2 j \left(1 + C(K_2 + K_3)\beta\right). \tag{2.36}$$

This completes the proof. □

The next lemma is the key to advancing the induction, as it provides bounds for  $e_{n+1}$  and  $g_{n+1}$ .

**Lemma 2.5.** *Let  $z \in I_n$ , and assume (H2), (H3) and (H4). For  $k \in [-\pi, \pi]^d$ ,  $2 \leq j \leq n + 1$ , and  $\epsilon' \in [0, \epsilon]$ , the following hold:*

- (i)  $|g_j(k; z)| \leq K'_4\beta j^{-d/2}$ ,
- (ii)  $|\nabla^2 g_j(0; z)| \leq K'_4\sigma^2\beta j^{-(d-2)/2}$ ,
- (iii)  $|\partial_z g_j(0; z)| \leq K'_4\beta j^{-(d-2)/2}$ ,
- (iv)  $|g_j(k; z) - g_j(0; z) - a(k)\sigma^{-2}\nabla^2 g_j(0; z)| \leq K'_4\beta a(k)^{1+\epsilon'} j^{-(d-2-2\epsilon')/2}$ ,
- (v)  $|e_j(k; z)| \leq K'_4\beta j^{-d/2}$ ,
- (vi)  $|e_j(k; z) - e_j(0; z)| \leq K'_4 a(k)\beta j^{-(d-2)/2}$ .

*Proof.* The bounds (1.27) for  $1 \leq m \leq n$  follow from Lemmas 2.2–2.4, with  $K = cK_4$  (this defines  $c$ ), assuming that  $\beta$  is sufficiently small. The bounds of the lemma then follow immediately from Assumptions E and G, with  $K'_4$  given in (2.3). □

### 3. The induction advanced

In this section we advance the induction hypotheses (H1–H4) from  $n$  to  $n + 1$ . Throughout this section, in accordance with the uniformity condition on (H2–H4), we fix  $z \in I_{n+1}$ . We frequently assume  $\beta \ll 1$  without explicit comment.

#### 3.1. Advancement of (H1)

By (2.5) and the mean-value theorem,

$$\begin{aligned} z_{n+1} - z_n &= - \sum_{m=2}^n [g_m(0; z_n) - g_m(0; z_{n-1})] - g_{n+1}(0; z_n) \\ &= -(z_n - z_{n-1}) \sum_{m=2}^n \partial_z g_m(0; y_n) - g_{n+1}(0; z_n), \end{aligned} \tag{3.1}$$

for some  $y_n$  between  $z_n$  and  $z_{n-1}$ . By (H1) and (2.6),  $y_n \in I_n$ . Using Lemma 2.5 and (H1), it then follows that

$$\begin{aligned} |z_{n+1} - z_n| &\leq K_1 \beta n^{-d/2} \sum_{m=2}^n K'_4 \beta m^{-(d-2)/2} + K'_4 \beta (n+1)^{-d/2} \\ &\leq K'_4 \beta (1 + CK_1 \beta) (n+1)^{-d/2}. \end{aligned} \tag{3.2}$$

Thus (H1) holds for  $n + 1$ , for  $\beta$  small and  $K_1 > K'_4$ .

Having advanced (H1) to  $n + 1$ , it then follows from Lemma 2.1 that  $I_1 \supset I_2 \supset \dots \supset I_{n+1}$ .

For  $n \geq 0$ , define

$$\zeta_{n+1} = \zeta_{n+1}(z) = \sum_{m=1}^{n+1} g_m(0; z) - 1 = \sum_{m=2}^{n+1} g_m(0; z) + z - 1. \tag{3.3}$$

The following lemma, whose proof makes use of (H1) for  $n + 1$ , will be needed in what follows.

**Lemma 3.1.** *For all  $z \in I_{n+1}$ ,*

$$|\zeta_{n+1}| \leq CK_1 \beta (n+1)^{-(d-2)/2}. \tag{3.4}$$

*Proof.* By (2.5) and the mean-value theorem,

$$\begin{aligned} |\zeta_{n+1}| &= \left| (z - z_{n+1}) + \sum_{m=2}^{n+1} [g_m(0; z) - g_m(0; z_n)] \right| \\ &= \left| (z - z_{n+1}) + (z - z_n) \sum_{m=2}^{n+1} \partial_z g_m(0; y_n) \right|, \end{aligned} \tag{3.5}$$

for some  $y_n$  between  $z$  and  $z_n$ . Since  $z \in I_{n+1} \subset I_n$  and  $z_n \in I_n$ , we have  $y_n \in I_n$ . Therefore, by Lemma 2.5,

$$\begin{aligned} |\zeta_{n+1}| &\leq K_1\beta(n+1)^{-(d-2)/2} + K_1\beta n^{-(d-2)/2} \sum_{m=2}^{n+1} K'_4\beta m^{-(d-2)/2} \\ &\leq K_1\beta(1 + CK'_4\beta)(n+1)^{-(d-2)/2}. \end{aligned} \tag{3.6}$$

The lemma then follows, for  $\beta$  sufficiently small. □

### 3.2. Advancement of (H2)

Let  $z \in I_{n+1}$ . As observed in Section 3.1, this implies that  $z \in I_j$  for all  $j \leq n + 1$ . The definitions in (2.1) imply that

$$v_{n+1} - v_n = \frac{1}{1 + c_{n+1}}(b_{n+1} - b_n) - \frac{b_n}{(1 + c_n)(1 + c_{n+1})}(c_{n+1} - c_n), \tag{3.7}$$

with

$$b_{n+1} - b_n = -\frac{1}{\sigma^2}\nabla^2 g_{n+1}(0), \quad c_{n+1} - c_n = n g_{n+1}(0). \tag{3.8}$$

By Lemma 2.5, both differences in (3.8) are bounded by  $K'_4\beta(n+1)^{-(d-2)/2}$ , and, in addition,

$$|b_j - 1| \leq CK'_4\beta, \quad |c_j| \leq CK'_4\beta \tag{3.9}$$

for  $1 \leq j \leq n + 1$ . Therefore

$$|v_{n+1} - v_n| \leq K_2\beta(n+1)^{-(d-2)/2}, \tag{3.10}$$

provided we assume  $K_2 \geq 3K'_4$ . This advances (H2).

### 3.3. Advancement of (H3)

#### 3.3.1. The decomposition

The advancement of the induction hypotheses (H3–H4) is the most technical part of the proof. For (H3), we fix  $k$  with  $a(k) \leq \gamma(n+1)^{-1} \log(n+1)$ , and  $z \in I_{n+1}$ . The induction step will be achieved as soon as we are able to write the ratio  $f_{n+1}(k)/f_n(k)$  as

$$\frac{f_{n+1}(k)}{f_n(k)} = 1 - v_{n+1}a(k) + r_{n+1}(k), \tag{3.11}$$

with  $r_{n+1}(0)$  and  $r_{n+1}(k) - r_{n+1}(0)$  satisfying the bounds required by (H3).

To begin, we divide the recursion relation (1.1) by  $f_n(k)$ , and use (3.3), to obtain

$$\frac{f_{n+1}(k)}{f_n(k)} = 1 + \sum_{m=1}^{n+1} \left[ g_m(k) \frac{f_{n+1-m}(k)}{f_n(k)} - g_m(0) \right] + \zeta_{n+1} + \frac{e_{n+1}(k)}{f_n(k)}. \tag{3.12}$$

By (2.1),

$$v_{n+1} = b_{n+1} - v_{n+1}c_{n+1} = -\sigma^{-2} \sum_{m=1}^{n+1} \nabla^2 g_m(0) - v_{n+1} \sum_{m=1}^{n+1} (m-1)g_m(0). \tag{3.13}$$

Thus we can rewrite (3.12) as

$$\frac{f_{n+1}(k)}{f_n(k)} = 1 - v_{n+1}a(k) + r_{n+1}(k), \tag{3.14}$$

where

$$r_{n+1}(k) = X(k) + Y(k) + Z(k) + \zeta_{n+1} \tag{3.15}$$

with

$$X(k) = \sum_{m=2}^{n+1} \left[ (g_m(k) - g_m(0)) \frac{f_{n+1-m}(k)}{f_n(k)} - a(k)\sigma^{-2}\nabla^2 g_m(0) \right], \tag{3.16}$$

$$Y(k) = \sum_{m=2}^{n+1} g_m(0) \left[ \frac{f_{n+1-m}(k)}{f_n(k)} - 1 - (m-1)v_{n+1}a(k) \right], \tag{3.17}$$

$$Z(k) = \frac{e_{n+1}(k)}{f_n(k)}. \tag{3.18}$$

The  $m = 1$  terms in  $X$  and  $Y$  vanish and have not been included.

We will prove that

$$|r_{n+1}(0)| \leq \frac{C(K_1 + K'_4)\beta}{(n+1)^{(d-2)/2}}, \quad |r_{n+1}(k) - r_{n+1}(0)| \leq \frac{CK'_4\beta a(k)}{(n+1)^\delta}. \tag{3.19}$$

This gives (H3) for  $n + 1$ , provided we assume that  $K_3 \gg K_1$  and  $K_3 \gg K'_4$ . To prove the bounds on  $r_{n+1}$  of (3.19), it will be convenient to make use of some elementary convolution bounds, as well as some bounds on ratios involving  $f_j$ . These preliminary bounds are given in Section 3.3.2, before we present the proof of (3.19) in Section 3.3.3.

### 3.3.2. Convolution and ratio bounds

The proof of (3.19) will make use of the following elementary convolution bounds. To keep the discussion simple, we do not obtain optimal bounds.

**Lemma 3.2.** *For  $n \geq 2$ ,*

$$\sum_{m=2}^n \frac{1}{m^a} \sum_{j=n-m+1}^n \frac{1}{j^b} \leq \begin{cases} Cn^{-(a \wedge b)+1} & \text{for } a, b > 1 \\ Cn^{-(a-2) \wedge b} & \text{for } a > 2, b > 0 \\ Cn^{-(a-1) \wedge b} & \text{for } a > 2, b > 1 \\ Cn^{-a \wedge b} & \text{for } a, b > 2. \end{cases} \tag{3.20}$$

*Proof.* Since  $m + j \geq n$ , either  $m$  or  $j$  is at least  $\frac{n}{2}$ . Therefore

$$\sum_{m=2}^n \frac{1}{m^a} \sum_{j=n-m+1}^n \frac{1}{j^b} \leq \left(\frac{2}{n}\right)^a \sum_{m=2}^n \sum_{j=n-m+1}^n \frac{1}{j^b} + \left(\frac{2}{n}\right)^b \sum_{m=2}^n \sum_{j=n-m+1}^n \frac{1}{m^a}. \tag{3.21}$$

If  $a, b > 1$ , then the first term is bounded by  $Cn^{1-a}$  and the second by  $Cn^{1-b}$ .  
 If  $a > 2, b > 0$ , then the first term is bounded by  $Cn^{2-a}$  and the second by  $Cn^{-b}$ .  
 If  $a > 2, b > 1$ , then the first term is bounded by  $Cn^{1-a}$  and the second by  $Cn^{-b}$ .  
 If  $a, b > 2$ , then the first term is bounded by  $Cn^{-a}$  and the second by  $Cn^{-b}$ .  $\square$

We also will make use of several estimates involving ratios. We begin with some preparation. Given a vector  $x = (x_l)$  with  $\sup_l |x_l| < 1$ , define  $\chi(x) = \sum_l \frac{|x_l|}{1-|x_l|}$ . The bound  $(1-t)^{-1} \leq \exp[t(1-t)^{-1}]$ , together with Taylor’s Theorem applied to  $f(t) = \prod_l \frac{1}{1-tx_l}$ , gives

$$\left| \prod_l \frac{1}{1-x_l} - 1 \right| \leq \chi(x)e^{\chi(x)}, \quad \left| \prod_l \frac{1}{1-x_l} - 1 - \sum_l x_l \right| \leq \chi(x)^2 e^{\chi(x)}. \tag{3.22}$$

We assume throughout the rest of this section that  $a(k) \leq \gamma(n+1)^{-1} \log(n+1)$  and  $2 \leq m \leq n+1$ , and define

$$\psi_{m,n} = \sum_{j=n+2-m}^n \frac{|r_j(0)|}{1-|r_j(0)|}, \quad \chi_{m,n}(k) = \sum_{j=n+2-m}^n \frac{v_j a(k) + |s_j(k)|}{1-v_j a(k) - |s_j(k)|}. \tag{3.23}$$

By (2.17) and (2.19),

$$\chi_{m,n}(k) \leq (m-1)a(k)Q(k) \quad \text{with} \quad Q(k) = [1 + C(K_2 + K_3)\beta][1 + Ca(k)]. \tag{3.24}$$

Since  $a(k) \leq \gamma(n+1)^{-1} \log(n+1)$ , we have  $Q(k) \leq [1 + C(K_2 + K_3)\beta][1 + C\gamma(n+1)^{-1} \log(n+1)]$ . Therefore

$$e^{\chi_{m,n}(k)} \leq e^{\gamma \log(n+1)Q(k)} \leq C(n+1)^{\gamma q}, \tag{3.25}$$

where  $q = 1 + C(K_2 + K_3)\beta$  may be taken to be as close to 1 as desired, by taking  $\beta$  to be small.

We now turn to the ratio bounds. It follows from (H3) and the first inequality of (3.22) that

$$\left| \frac{f_{n+1-m}(0)}{f_n(0)} - 1 \right| \leq \psi_{m,n} e^{\psi_{m,n}} \leq \sum_{j=n+2-m}^n \frac{CK_3\beta}{j^{(d-2)/2}} \leq \frac{CK_3\beta}{(n+2-m)^{(d-4)/2}}. \tag{3.26}$$

Therefore

$$\left| \frac{f_{n+1-m}(0)}{f_n(0)} \right| \leq 1 + CK_3\beta. \tag{3.27}$$

By (2.18),

$$\begin{aligned} \left| \frac{f_{n+1-m}(k)}{f_n(k)} - 1 \right| &\leq \left| \frac{f_{n+1-m}(0)}{f_n(0)} \right| \left| \prod_{j=n+2-m}^n [1 - v_j a(k) + s_j(k)]^{-1} - 1 \right| \\ &\quad + \left| \frac{f_{n+1-m}(0)}{f_n(0)} - 1 \right|. \end{aligned} \tag{3.28}$$

The first inequality of (3.22), together with (3.24)–(3.27), then gives

$$\left| \frac{f_{n+1-m}(k)}{f_n(k)} - 1 \right| \leq C(m-1)a(k)(n+1)^{\gamma q} + \frac{CK_3\beta}{(n+2-m)^{(d-4)/2}}. \tag{3.29}$$

Similarly,

$$\left| \frac{f_n(0)}{f_n(k)} - 1 \right| \leq \chi_{n+1,n}(k)e^{\chi_{n+1,n}(k)} \leq Ca(k)(n+1)^{1+\gamma q}. \tag{3.30}$$

Next, we estimate the quantity  $R_{m,n}(k)$ , which is defined by

$$R_{m,n}(k) = \prod_{j=n+2-m}^n [1 - v_j a(k) + s_j(k)]^{-1} - 1 - \sum_{j=n+2-m}^n [v_j a(k) - s_j(k)]. \tag{3.31}$$

By the second inequality of (3.22), together with (3.24) and (3.25), this obeys

$$|R_{m,n}(k)| \leq \chi_{m,n}(k)^2 e^{\chi_{m,n}(k)} \leq Cm^2 a(k)^2 (n+1)^{\gamma q}. \tag{3.32}$$

Finally, we apply (H3) to obtain

$$\begin{aligned} \left| \frac{f_{m-1}(k)}{f_m(k)} - 1 \right| &= \left| [1 - v_m a(k) + (r_m(k) - r_m(0)) + r_m(0)]^{-1} - 1 \right| \\ &\leq Ca(k) + \frac{CK_3\beta}{m^{(d-2)/2}}. \end{aligned} \tag{3.33}$$

### 3.3.3. The induction step

By definition,

$$r_{n+1}(0) = Y(0) + Z(0) + \zeta_{n+1} \tag{3.34}$$

and

$$r_{n+1}(k) - r_{n+1}(0) = X(k) + (Y(k) - Y(0)) + (Z(k) - Z(0)). \tag{3.35}$$

Since  $|\zeta_{n+1}| \leq CK_1\beta(n+1)^{-(d-2)/2}$  by Lemma 3.1, to prove (3.19) it suffices to show that

$$|Y(0)| \leq CK'_4\beta(n+1)^{-(d-2)/2}, \quad |Z(0)| \leq CK'_4\beta(n+1)^{-(d-2)/2} \quad (3.36)$$

and

$$\begin{aligned} |X(k)| &\leq CK'_4\beta a(k)(n+1)^{-\delta}, \quad |Y(k) - Y(0)| \leq CK'_4\beta a(k)(n+1)^{-\delta}, \\ |Z(k) - Z(0)| &\leq CK'_4\beta a(k)(n+1)^{-\delta}. \end{aligned} \quad (3.37)$$

The remainder of the proof is devoted to establishing (3.36) and (3.37).

*Bound on X.* We write  $X$  as  $X = X_1 + X_2$ , with

$$X_1 = \sum_{m=2}^{n+1} \left[ g_m(k) - g_m(0) - a(k)\sigma^{-2}\nabla^2 g_m(0) \right], \quad (3.38)$$

$$X_2 = \sum_{m=2}^{n+1} \left[ g_m(k) - g_m(0) \right] \left[ \frac{f_{n+1-m}(k)}{f_n(k)} - 1 \right]. \quad (3.39)$$

The term  $X_1$  is bounded using Lemma 2.5(iv) with  $\epsilon' \in (\delta, \epsilon)$ , and using the fact that  $a(k) \leq \gamma(n+1)^{-1} \log(n+1)$ , by

$$|X_1| \leq K'_4\beta a(k)^{1+\epsilon'} \sum_{m=2}^{n+1} \frac{1}{m^{(d-2-2\epsilon')/2}} \leq CK'_4\beta a(k)^{1+\epsilon'} \leq \frac{CK'_4\beta a(k)}{(n+1)^\delta}. \quad (3.40)$$

For  $X_2$ , we first apply Lemma 2.5(ii,iv), with  $\epsilon' = 0$ , to obtain

$$|g_m(k) - g_m(0)| \leq 2K'_4\beta a(k)m^{-(d-2)/2}. \quad (3.41)$$

Applying (3.29) then gives

$$|X_2| \leq CK'_4\beta a(k) \sum_{m=2}^{n+1} \frac{1}{m^{(d-2)/2}} \left( (m-1)a(k)(n+1)^{\gamma q} + \frac{K_3\beta}{(n+2-m)^{(d-4)/2}} \right). \quad (3.42)$$

By an elementary estimate, the contribution from the second term on the right side is bounded above by  $CK_3K'_4\beta^2 a(k)(n+1)^{-(d-4)/2}$ . The first term is bounded above by

$$CK'_4\beta a(k)(n+1)^{\gamma q-1} \log(n+1) \times \begin{cases} (n+1)^{0 \vee (6-d)/2} & (d \neq 6) \\ \log(n+1) & (d = 6). \end{cases} \quad (3.43)$$

Since we may choose  $q$  to be as close to 1 as desired, and since  $\delta + \gamma < 1 \wedge \frac{d-4}{2}$  by (2.2), this is bounded above by  $CK'_4\beta a(k)(n+1)^{-\delta}$ . With (3.40), this proves the bound on  $X$  in (3.37).

Bound on  $Y$ . By (2.18),

$$\frac{f_{n+1-m}(k)}{f_n(k)} = \frac{f_{n+1-m}(0)}{f_n(0)} \prod_{j=n+2-m}^n [1 - v_j a(k) + s_j(k)]^{-1}. \tag{3.44}$$

Recalling the definition of  $R_{m,n}(k)$  in (3.31), we can therefore decompose  $Y$  as  $Y = Y_1 + Y_2 + Y_3 + Y_4$  with

$$Y_1 = \sum_{m=2}^{n+1} g_m(0) \frac{f_{n+1-m}(0)}{f_n(0)} R_{m,n}(k), \tag{3.45}$$

$$Y_2 = \sum_{m=2}^{n+1} g_m(0) \frac{f_{n+1-m}(0)}{f_n(0)} \sum_{j=n+2-m}^n [(v_j - v_{n+1})a(k) - s_j(k)], \tag{3.46}$$

$$Y_3 = \sum_{m=2}^{n+1} g_m(0) \left[ \frac{f_{n+1-m}(0)}{f_n(0)} - 1 \right] (m - 1)v_{n+1}a(k), \tag{3.47}$$

$$Y_4 = \sum_{m=2}^{n+1} g_m(0) \left[ \frac{f_{n+1-m}(0)}{f_n(0)} - 1 \right]. \tag{3.48}$$

Then

$$Y(0) = Y_4 \quad \text{and} \quad Y(k) - Y(0) = Y_1 + Y_2 + Y_3. \tag{3.49}$$

For  $Y_1$ , we use Lemma 2.5, (3.27) and (3.32) to obtain

$$|Y_1| \leq CK'_4 \beta a(k)^2 (n + 1)^{\gamma q} \sum_{m=2}^n \frac{1}{m^{(d-4)/2}}. \tag{3.50}$$

As in the analysis of the first term of (3.42), we therefore have

$$|Y_1| \leq \frac{CK'_4 \beta a(k)}{(n + 1)^\delta}. \tag{3.51}$$

For  $Y_2$ , we use Lemma 2.5, (3.27), (H2) (now established up to  $n + 1$ ), (2.19) and Lemma 3.2 to obtain

$$|Y_2| \leq \sum_{m=2}^{n+1} \frac{K'_4 \beta}{m^{d/2}} C \sum_{j=n+2-m}^n \left[ \frac{K_2 \beta a(k)}{j^{(d-4)/2}} + \frac{K_3 \beta a(k)}{j^\delta} \right] \leq \frac{CK'_4 (K_2 + K_3) \beta^2 a(k)}{(n + 1)^\delta}. \tag{3.52}$$

The term  $Y_3$  obeys

$$|Y_3| \leq \sum_{m=2}^{n+1} \frac{K'_4 \beta}{m^{(d-2)/2}} \frac{CK_3 \beta}{(n + 2 - m)^{(d-4)/2}} a(k) \leq \frac{CK'_4 K_3 \beta^2 a(k)}{(n + 1)^{(d-4)/2}}, \tag{3.53}$$

where we used Lemma 2.5, (3.26), (2.17), and an elementary convolution bound. This proves the bound on  $|Y(k) - Y(0)|$  of (3.37), if  $\beta$  is sufficiently small.

We bound  $Y_4$  in a similar fashion, using Lemma 3.2 and the intermediate bound of (3.26) to obtain

$$|Y_4| \leq \sum_{m=2}^{n+1} \frac{K'_4 \beta}{m^{d/2}} \sum_{j=n+2-m}^n \frac{CK_3 \beta}{j^{(d-2)/2}} \leq \frac{CK'_4 K_3 \beta^2}{(n+1)^{(d-2)/2}}. \tag{3.54}$$

Taking  $\beta$  small then gives the bound on  $Y(0)$  of (3.36).

*Bound on  $Z$ .* We decompose  $Z$  as

$$\begin{aligned} Z &= \frac{e_{n+1}(0)}{f_n(0)} + \frac{1}{f_n(0)} [e_{n+1}(k) - e_{n+1}(0)] + \frac{e_{n+1}(k)}{f_n(0)} \left[ \frac{f_n(0)}{f_n(k)} - 1 \right] \\ &= Z_1 + Z_2 + Z_3. \end{aligned} \tag{3.55}$$

Then

$$Z(0) = Z_1 \quad \text{and} \quad Z(k) - Z(0) = Z_2 + Z_3. \tag{3.56}$$

Using Lemma 2.5(v,vi), and (3.27) with  $m = n + 1$ , we obtain

$$|Z_1| \leq CK'_4 \beta (n+1)^{-d/2} \quad \text{and} \quad |Z_2| \leq CK'_4 \beta a(k) (n+1)^{-(d-2)/2}. \tag{3.57}$$

Also, by (3.27) and (3.30), we have

$$|Z_3| \leq CK'_4 \beta (n+1)^{-d/2} a(k) (n+1)^{1+\gamma q} \leq CK'_4 \beta a(k) (n+1)^{-(1+\delta)}. \tag{3.58}$$

This completes the proof of (3.19), and hence completes the advancement of (H3) to  $n + 1$ .

### 3.4. Advancement of (H4)

In this section, we fix  $a(k) > \gamma(n+1)^{-1} \log(n+1)$ . To advance (H4) to  $j = n + 1$ , we first recall the definitions of  $b_{n+1}$ ,  $\zeta_{n+1}$  and  $X_1$  from (2.1), (3.3) and (3.38). After some algebra, (1.1) can be rewritten as

$$f_{n+1}(k) = f_n(k) \left( 1 - a(k)b_{n+1} + X_1 + \zeta_{n+1} \right) + W + e_{n+1}(k), \tag{3.59}$$

with

$$W = \sum_{m=2}^{n+1} g_m(k) [f_{n+1-m}(k) - f_n(k)]. \tag{3.60}$$

We already have estimates for most of the relevant terms. By Lemma 3.1, we have  $|\zeta_{n+1}| \leq CK_1 \beta (n+1)^{-(d-2)/2}$ . By (3.40),  $|X_1| \leq CK'_4 \beta a(k)^{1+\epsilon'}$ , for any  $\epsilon' \in (\delta, \epsilon)$ . By Lemma 2.5(v),  $|e_{n+1}(k)| \leq K'_4 \beta (n+1)^{-d/2}$ . It remains to estimate  $W$ . We will show below that  $W$  obeys the bound

$$|W| \leq \frac{CK'_4 \beta}{a(k)^{1+\rho} (n+1)^{d/2}} (1 + K_3 \beta + K_5). \tag{3.61}$$

Before proving (3.61), we will first show that it is sufficient for the advancement of (H4).

In preparation for this, we first note that it suffices to consider only large  $n$ . In fact, since  $|f_n(k; z)|$  is bounded uniformly in  $k$  and in  $z$  in a compact set by Assumption S, and since  $a(k) \leq 2$ , it is clear that both inequalities of (H4) hold for all  $n \leq N$ , if we choose  $K_4$  and  $K_5$  large enough (depending on  $N$ ). We therefore assume in the following that  $n \geq N$  with  $N$  large.

Also, care is required to invoke (H3) or (H4), as applicable, in estimating the factor  $f_n(k)$  of (3.59). Given  $k$ , (H3) should be used for the value  $n$  for which  $\gamma(n+1)^{-1} \log(n+1) < a(k) \leq \gamma n^{-1} \log n$  ((H4) should be used for larger  $n$ ). We will now show that, as anticipated in the discussion of (H4) in Section 2.3, the bound of (H3) actually implies the first bound of (H4) in this case. To see this, we use Lemma 2.2 to see that there are  $q, q'$  arbitrarily close to 1 such that

$$\begin{aligned} |f_n(k)| &\leq C e^{-qa(k)n} \leq \frac{C}{(n+1)^{q\gamma n/(n+1)}} \leq \frac{C}{n^{q'\gamma}} \\ &\leq \frac{C}{n^{d/2}} \frac{n^{2+\rho}}{n^{q'\gamma+2+\rho-d/2}} \leq \frac{C}{n^{d/2} a(k)^{2+\rho}}, \end{aligned} \tag{3.62}$$

where we used the fact that  $\gamma + 2 + \rho - d/2 > 0$  by (2.2). Thus, taking  $K_4 \gg 1$ , we may use the first bound of (H4) also for the value of  $n$  to which (H3) nominally applies. We will do so in what follows, without further comment.

*Advancement of the second bound of (H4) assuming (3.61).* To advance the second estimate in (H4), we use (3.59), (H4), and the bounds found above, to obtain

$$\begin{aligned} |f_{n+1}(k) - f_n(k)| &\leq |f_n(k)| \left| -a(k)b_{n+1} + X_1 + \zeta_{n+1} \right| + |W| + |e_{n+1}(k)| \\ &\leq \frac{K_4}{n^{d/2} a(k)^{2+\rho}} \left( a(k)b_{n+1} + CK'_4 \beta a(k)^{1+\epsilon'} + \frac{CK_1 \beta}{(n+1)^{(d-2)/2}} \right) \\ &\quad + \frac{CK'_4 \beta (1 + K_3 \beta + K_5)}{(n+1)^{d/2} a(k)^{1+\rho}} + \frac{K'_4 \beta}{(n+1)^{d/2}}. \end{aligned} \tag{3.63}$$

Since  $b_{n+1} = 1 + \mathcal{O}(\beta)$  by (3.9), and since  $(n+1)^{-(d-2)/2} < [a(k)/\gamma \log(n+1)]^{(d-2)/2} \leq Ca(k)$ , the second estimate in (H4) follows for  $n+1$  provided  $K_5 \gg K_4$  and  $\beta$  is sufficiently small.

*Advancement of the first bound of (H4) assuming (3.61).* To advance the first estimate of (H4), we argue as in (3.63) to obtain

$$\begin{aligned} |f_{n+1}(k)| &\leq |f_n(k)| \left| 1 - a(k)b_{n+1} + X_1 + \zeta_{n+1} \right| + |W| + |e_{n+1}(k)| \\ &\leq \frac{K_4}{n^{d/2} a(k)^{2+\rho}} \left( |1 - a(k)b_{n+1}| + CK'_4 \beta a(k)^{1+\epsilon'} + \frac{CK_1 \beta}{(n+1)^{(d-2)/2}} \right) \\ &\quad + \frac{CK'_4 \beta (1 + K_3 \beta + K_5)}{(n+1)^{d/2} a(k)^{1+\rho}} + \frac{K'_4 \beta}{(n+1)^{d/2}}. \end{aligned} \tag{3.64}$$

We need to argue that the right-hand side is no larger than  $K_4(n+1)^{-d/2}a(k)^{-2-\rho}$ . To achieve this, we will use separate arguments for  $a(k) \leq \frac{1}{2}$  and  $a(k) > \frac{1}{2}$ . These arguments will be valid only when  $n$  is large enough.

Suppose that  $a(k) \leq \frac{1}{2}$ . Since  $b_{n+1} = 1 + \mathcal{O}(\beta)$  by (3.9), for  $\beta$  sufficiently small we have

$$1 - b_{n+1}a(k) \geq 0. \tag{3.65}$$

Hence, the absolute value signs on the right side of (3.64) may be removed. Therefore, to obtain the first estimate of (H4) for  $n + 1$ , it now suffices to show that

$$1 - ca(k) + \frac{CK_1\beta}{(n+1)^{(d-2)/2}} \leq \frac{n^{d/2}}{(n+1)^{d/2}}, \tag{3.66}$$

for  $c$  within order  $\beta$  of 1. The term  $ca(k)$  has been introduced to absorb  $b_{n+1}a(k)$ , the order  $\beta$  term in (3.64) involving  $a(k)^{1+\epsilon'}$ , and the last two terms of (3.64). However,  $a(k) > \gamma(n+1)^{-1} \log(n+1)$ . From this, it can be seen that (3.66) holds for  $n$  sufficiently large and  $\beta$  sufficiently small.

Suppose, on the other hand, that  $a(k) > \frac{1}{2}$ . By (1.21), there is a positive  $\eta$ , which we may assume lies in  $(0, \frac{1}{2})$ , such that  $-1 + \eta < 1 - a(k) < \frac{1}{2}$ . Therefore  $|1 - a(k)| \leq 1 - \eta$  and

$$|1 - b_{n+1}a(k)| \leq |1 - a(k)| + |b_{n+1} - 1| |a(k)| \leq 1 - \eta + 2|b_{n+1} - 1|. \tag{3.67}$$

Hence

$$|1 - a(k)b_{n+1}| + CK'_4\beta a(k)^{1+\epsilon'} + \frac{CK_1\beta}{(n+1)^{(d-2)/2}} \leq 1 - \eta + C(K_1 + K'_4)\beta, \tag{3.68}$$

and the right side of (3.64) is at most

$$\begin{aligned} & \frac{K_4}{n^{d/2}a(k)^{2+\rho}} [1 - \eta + C(K_1 + K'_4)\beta] + \frac{CK'_4(1 + K_3\beta + K_5)\beta}{(n+1)^{d/2}a(k)^{2+\rho}} \\ & \leq \frac{K_4}{n^{d/2}a(k)^{2+\rho}} [1 - \eta + C(K_5K'_4 + K_1)\beta]. \end{aligned} \tag{3.69}$$

This is less than  $K_4(n+1)^{-d/2}a(k)^{-2-\rho}$  if  $n$  is large and  $\beta$  is sufficiently small.

This advances the first bound in (H4), assuming (3.61).

*Bound on  $W$ .* We now obtain the bound (3.61) on  $W$ . As a first step, we rewrite  $W$  as

$$W = \sum_{j=0}^{n-1} g_{n+1-j}(k) \sum_{l=j+1}^n [f_{l-1}(k) - f_l(k)]. \tag{3.70}$$

Let

$$m(k) = \begin{cases} 1 & (a(k) > \gamma 3^{-1} \log 3) \\ \max\{l \in \{3, \dots, n\} : a(k) \leq \gamma l^{-1} \log l\} & (a(k) \leq \gamma 3^{-1} \log 3). \end{cases} \tag{3.71}$$

For  $l \leq m(k)$ ,  $f_l$  is in the domain of (H3), while for  $l > m(k)$ ,  $f_l$  is in the domain of (H4). By hypothesis,  $a(k) > \gamma(n+1)^{-1} \log(n+1)$ . We divide the sum over  $l$  into two parts, corresponding respectively to  $l \leq m(k)$  and  $l > m(k)$ , yielding  $W = W_1 + W_2$ . By Lemma 2.5(i),

$$|W_1| \leq \sum_{j=0}^{m(k)} \frac{K'_4 \beta}{(n+1-j)^{d/2}} \sum_{l=j+1}^{m(k)} |f_{l-1}(k) - f_l(k)| \tag{3.72}$$

$$|W_2| \leq \sum_{j=0}^{n-1} \frac{K'_4 \beta}{(n+1-j)^{d/2}} \sum_{l=(m(k) \vee j)+1}^n |f_{l-1}(k) - f_l(k)|. \tag{3.73}$$

The term  $W_2$  is easy, since by (H4) and Lemma 3.2 we have

$$|W_2| \leq \sum_{j=0}^{n-1} \frac{K'_4 \beta}{(n+1-j)^{d/2}} \sum_{l=j+1}^n \frac{K_5}{a(k)^{1+\rho} l^{d/2}} \leq \frac{CK_5 K'_4 \beta}{a(k)^{1+\rho} (n+1)^{d/2}}. \tag{3.74}$$

For  $W_1$ , we have the estimate

$$|W_1| \leq \sum_{j=0}^{m(k)} \frac{K'_4 \beta}{(n+1-j)^{d/2}} \sum_{l=j+1}^{m(k)} |f_{l-1}(k) - f_l(k)|. \tag{3.75}$$

For  $1 \leq l \leq m(k)$ , it follows from Lemma 2.2 and (3.33) that

$$|f_{l-1}(k) - f_l(k)| \leq C e^{-qa(k)l} \left( a(k) + \frac{K_3 \beta}{l^{(d-2)/2}} \right), \tag{3.76}$$

with  $q = 1 - \mathcal{O}(\beta)$ . We fix a small  $r > 0$ , and bound the summation over  $j$  in (3.75) by summing separately over  $j$  in the ranges  $1 \leq j \leq (1-r)n$  and  $(1-r)n \leq j \leq m(k)$  (the latter range may be empty). We denote the contributions from these two sums by  $W_{1,1}$  and  $W_{1,2}$  respectively.

To estimate  $W_{1,1}$ , we will make use of the bound

$$\sum_{l=j+1}^{\infty} e^{-qa(k)l} l^{-p} \leq C e^{-qa(k)j} \quad (p > 1). \tag{3.77}$$

With (3.75) and (3.76), this gives

$$\begin{aligned} |W_{1,1}| &\leq \frac{CK'_4 \beta}{(n+1)^{d/2}} \sum_{j=0}^{(1-r)n} e^{-qa(k)j} (1 + K_3 \beta) \\ &\leq \frac{CK'_4 \beta}{(n+1)^{d/2}} \frac{1 + K_3 \beta}{a(k)} \leq \frac{CK'_4 \beta}{(n+1)^{d/2}} \frac{1 + K_3 \beta}{a(k)^{1+\rho}}. \end{aligned} \tag{3.78}$$

For  $W_{1,2}$ , we have

$$|W_{1,2}| \leq \sum_{j=(1-r)n}^{m(k)} \frac{CK'_4\beta}{(n+1-j)^{d/2}} \sum_{l=j+1}^{m(k)} e^{-qa(k)l} \left( a(k) + \frac{K_3\beta}{l^{(d-2)/2}} \right). \quad (3.79)$$

Since  $l$  and  $m(k)$  are comparable and large, it follows as in (3.62) that

$$e^{-qa(k)l} \left( a(k) + \frac{K_3\beta}{l^{(d-2)/2}} \right) \leq \frac{C}{a(k)^{2+\rho}l^{d/2}} \left( a(k) + \frac{K_3\beta}{l^{(d-2)/2}} \right) \leq \frac{C(1+K_3\beta)}{a(k)^{1+\rho}l^{d/2}}. \quad (3.80)$$

Hence, by Lemma 3.2,

$$\begin{aligned} |W_{1,2}| &\leq \frac{C(1+K_3\beta)K'_4\beta}{a(k)^{1+\rho}} \sum_{j=(1-r)n}^{m(k)} \frac{1}{(n+1-j)^{d/2}} \sum_{l=j+1}^{m(k)} \frac{1}{l^{d/2}} \\ &\leq \frac{C(1+K_3\beta)K'_4\beta}{a(k)^{1+\rho}(n+1)^{d/2}}. \end{aligned} \quad (3.81)$$

Summarising, by (3.78), (3.81), and (3.74), we have

$$|W| \leq |W_{1,1}| + |W_{1,2}| + |W_2| \leq \frac{CK'_4\beta}{a(k)^{1+\rho}(n+1)^{d/2}} (1 + K_3\beta + K_5), \quad (3.82)$$

which proves (3.61).

#### 4. Proof of the main results

As a consequence of the completed induction, it follows from Lemma 2.1 that  $I_1 \supset I_2 \supset I_3 \supset \dots$ , so  $\bigcap_{n=1}^{\infty} I_n$  consists of a single point  $z = z_c$ . Since  $z_0 = 1$ , it follows from (H1) that  $z_c = 1 + \mathcal{O}(\beta)$ . We fix  $z = z_c$  throughout this section. The constant  $A$  is defined by  $A = \prod_{i=1}^{\infty} [1 + r_i(0)] = 1 + \mathcal{O}(\beta)$ . By (H2), the sequence  $v_n(z_c)$  is a Cauchy sequence. The constant  $v$  is defined to be the limit of this Cauchy sequence. By (H2),  $v = 1 + \mathcal{O}(\beta)$  and

$$|v_n(z_c) - v| \leq \mathcal{O}(\beta n^{-(d-4)/2}). \quad (4.1)$$

##### 4.1. Proof of Theorem 1.1

*Proof of Theorem 1.1(a).* By (H3),

$$|f_n(0; z_c) - A| = \left| \prod_{i=1}^n [1 + r_i(0)] - \prod_{i=n+1}^{\infty} [1 + r_i(0)] \right| \leq \mathcal{O}(\beta n^{-(d-4)/2}). \quad (4.2)$$

Suppose  $k$  is such that  $a(k/\sqrt{\sigma^2 vn}) \leq \gamma n^{-1} \log n$ , so that (H3) applies. Here, we use the  $\gamma$  of (2.2). By (1.17),  $a(k) = \sigma^2 k^2 / 2d + \mathcal{O}(k^{2+2\epsilon})$  with  $\epsilon > \delta$ , where

we now allow constants in error terms to depend on  $L$ . Using this, together with (2.18)–(2.19), (4.1), and  $\delta < 1 \wedge \frac{d-4}{2} \wedge \epsilon$ , we obtain

$$\begin{aligned} \frac{f_n(k/\sqrt{v\sigma^2n}; z_c)}{f_n(0; z_c)} &= \prod_{i=1}^n \left[ 1 - v_i a\left(\frac{k}{\sqrt{v\sigma^2n}}\right) + \mathcal{O}\left(\beta a\left(\frac{k}{\sqrt{v\sigma^2n}}\right) i^{-\delta}\right) \right] \\ &= e^{-k^2/2d} [1 + \mathcal{O}(k^{2+2\epsilon} n^{-\epsilon}) + \mathcal{O}(k^2 n^{-\delta})]. \end{aligned} \tag{4.3}$$

With (4.2), this gives the desired result.

*Proof of Theorem 1.1(b).* Since  $\delta < 1 \wedge \frac{d-4}{4}$ , it follows from (2.34)–(2.35) and (4.1)–(4.2) that

$$\frac{\nabla^2 f_n(0; z_c)}{f_n(0; z_c)} = -v\sigma^2 n [1 + \mathcal{O}(\beta n^{-\delta})]. \tag{4.4}$$

*Proof of Theorem 1.1(c).* The claim is immediate from Lemma 2.3, which is now known to hold for all  $n$ .

*Proof of Theorem 1.1(d).* Throughout this proof, we fix  $z = z_c$  and drop  $z_c$  from the notation. The identity (1.31) follows after we let  $n \rightarrow \infty$  in (3.3), using Lemma 3.1.

To determine  $A$ , we use a summation argument. Let  $\chi_n = \sum_{k=0}^n f_k(0)$ . By (1.1),

$$\begin{aligned} \chi_n &= 1 + \sum_{j=1}^n f_j(0) = 1 + \sum_{j=1}^n \sum_{m=1}^j g_m(0) f_{j-m}(0) + \sum_{j=1}^n e_j(0) \\ &= 1 + z_c \chi_{n-1} + \sum_{m=2}^n g_m(0) \chi_{n-m} + \sum_{m=1}^n e_m(0). \end{aligned} \tag{4.5}$$

Using (3.3) to rewrite  $z_c$ , this gives

$$f_n(0) = \chi_n - \chi_{n-1} = 1 + \zeta_n \chi_{n-1} - \sum_{m=2}^n g_m(0) (\chi_{n-1} - \chi_{n-m}) + \sum_{m=1}^n e_m(0). \tag{4.6}$$

By Theorem 1.1(a),  $\chi_n \sim nA$  as  $n \rightarrow \infty$ . Therefore, using Lemma 3.1 to bound the  $\zeta_n$  term, taking the limit  $n \rightarrow \infty$  in the above equation gives

$$A = 1 - A \sum_{m=2}^{\infty} (m-1) g_m(0) + \sum_{m=1}^{\infty} e_m(0). \tag{4.7}$$

With (1.31), this gives (1.32).

To prove (1.33), we use (4.1), (2.1) and Lemma 2.5 to obtain

$$v = \lim_{n \rightarrow \infty} v_n = \frac{-\sigma^{-2} \sum_{m=2}^{\infty} \nabla^2 g_m(0)}{1 + \sum_{m=2}^{\infty} (m-1) g_m(0)}. \tag{4.8}$$

Equation (1.33) then follows, once we rewrite the denominator using (1.31).

4.2. Proof of Theorem 1.2

By Theorem 1.1(a),  $\chi(z_c) = \infty$ . Therefore  $z_c \geq p_c$ . We need to rule out the possibility that  $z_c > p_c$ . Theorem 1.1 also gives (1.27) at  $z = z_c$ . By assumption, the series

$$G(z) = \sum_{m=2}^{\infty} g_m(0; z), \quad E(z) = \sum_{m=2}^{\infty} e_m(0; z) \tag{4.9}$$

therefore both converge absolutely and are  $\mathcal{O}(\beta)$  uniformly in  $z \leq z_c$ . For  $z < p_c$ , since the series defining  $\chi(z)$  converges absolutely, the basic recursion relation (1.1) gives

$$\chi(z) = 1 + z\chi(z) + G(z)\chi(z) + E(z), \tag{4.10}$$

and hence

$$\chi(z) = \frac{1 + E(z)}{1 - z - G(z)} \quad (z < p_c). \tag{4.11}$$

It is implicit in the bound on  $\partial_z g_m(k; z)$  of Assumption G that  $g_m(k; \cdot)$  is continuous on  $[0, z_c]$ . By dominated convergence,  $G$  is also continuous on  $[0, z_c]$ . Since  $E(z) = \mathcal{O}(\beta)$  and  $\lim_{z \uparrow p_c} \chi(z) = \infty$ , it then follows from (4.11) that

$$1 - p_c - G(p_c) = 0. \tag{4.12}$$

By (1.31), (4.12) holds also when  $p_c$  is replaced by  $z_c$ . If  $p_c \neq z_c$ , then it follows from the mean-value theorem that

$$z_c - p_c = G(p_c) - G(z_c) = -(z_c - p_c) \sum_{m=2}^{\infty} \partial_z g_m(0; t) \tag{4.13}$$

for some  $t \in (p_c, z_c)$ . However, by a bound of Assumption G, the sum on the right side is  $\mathcal{O}(\beta)$  uniformly in  $t \leq z_c$ . This is a contradiction, so we conclude that  $z_c = p_c$ .

4.3. Proof of Theorem 1.3

We begin by noting that

$$\frac{1}{(2R + 1)^d} \sum_{y \in C_R(\lfloor x\sqrt{v\sigma^2 n} \rfloor)} p_n(y) = (Q_R * p_n)(\lfloor x\sqrt{n} \rfloor), \tag{4.14}$$

where  $Q_R(x) = (2R + 1)^{-d}$  for  $x \in C_R(0)$ , and otherwise equals zero. Therefore,

$$\frac{1}{(2R + 1)^d} \sum_{y \in C_R(\lfloor x\sqrt{v\sigma^2 n} \rfloor)} p_n(y) = \int_{[-\pi, \pi]^d} \frac{d^d k}{(2\pi)^d} e^{-ik \cdot \lfloor x\sqrt{v\sigma^2 n} \rfloor} \hat{Q}_R(k) f_n(k; z_c). \tag{4.15}$$

Before proceeding with the proof, we first derive some relevant properties of  $\hat{Q}_R$ .

By definition,  $|\hat{Q}_R(k)| \leq 1$ . The Fourier transform  $\hat{Q}_R(k)$  is given in terms of the Dirichlet kernel

$$M(t) = \sum_{j=-R}^R e^{ijt} = \frac{\sin[(2R + 1)t/2]}{\sin(t/2)} \tag{4.16}$$

by

$$\hat{Q}_R(k) = \prod_{i=1}^d \frac{M(k_i)}{2R + 1}. \tag{4.17}$$

Since  $|M(t)|$  is bounded above by both  $2R + 1$  and  $|\sin(t/2)|^{-1}$ , we have

$$|\hat{Q}_R(k)| \leq \frac{1}{(2R + 1) \sin(\|k\|_\infty/2)} \leq \frac{\pi}{2\|k\|_\infty R}. \tag{4.18}$$

Let  $\kappa \in [0, 1]$ . When the right side of (4.18) is less than 1, the bound is degraded by raising the right side to the power  $\kappa$ . On the other hand, when the right side is greater than 1, we may still raise it to the power  $\kappa$  and have a correct bound, since the left side is at most 1. Therefore, for every  $\kappa \in [0, 1]$ , we have

$$|\hat{Q}_R(k)| \leq \left[ \frac{\pi}{2\|k\|_\infty R} \right]^\kappa. \tag{4.19}$$

We divide the domain of integration  $[-\pi, \pi]^d$  of (4.15) into a small- $k$  region  $S_n = \{k \in [-\pi, \pi]^d : a(k) \leq \gamma n^{-1} \log n\}$  and a large- $k$  region  $L_n = \{k \in [-\pi, \pi]^d : a(k) > \gamma n^{-1} \log n\}$ . By Theorem 1.1(a), the integral over the small- $k$  region equals

$$A \int_{S_n} \frac{d^d k}{(2\pi)^d} e^{-ik \cdot [x\sqrt{v\sigma^2 n}]} e^{-nv\sigma^2 k^2/2d} [1 + \mathcal{O}(k^2 n^{1-\delta}) + \mathcal{O}(k^2 R^2)], \tag{4.20}$$

where we allow constants in error terms to depend on  $L$ . For the error terms, we bound the complex exponential by 1 and obtain a factor  $n^{-d/2}$  from the integration. For the leading term, we extend the integration domain to  $\mathbb{R}^d$  and perform the integral exactly. The error incurred by extending the integration domain is at most  $\mathcal{O}(n^{-d/2-\gamma^-})$ , where  $\gamma^-$  is any positive number less than  $\gamma$ . Therefore, the small- $k$  region gives

$$A \left( \frac{d}{2\pi v\sigma^2 n} \right)^{d/2} \left[ e^{-dx^2/2} + \mathcal{O}(n^{-\gamma^-}) + \mathcal{O}(n^{-\delta}) + \mathcal{O}(R^2 n^{-1}) \right]. \tag{4.21}$$

The integral over the large- $k$  region can be bounded using (H4), (4.19) and (1.19) by

$$K_4 n^{-d/2} \int_{L_n} \frac{d^d k}{(2\pi)^d} |\hat{Q}_R(k)| a(k)^{-2-\rho} \leq \mathcal{O}(n^{-d/2} R^{-\kappa}) \int_{L_n} d^d k k^{-4-2\rho-\kappa}. \tag{4.22}$$

The integral is bounded provided  $\kappa < d - 4 + 2\rho$ . By (2.2), we may choose a positive  $\kappa$  obeying this bound.

Therefore, the left side of (4.15) is given by

$$A \left( \frac{d}{2\pi v \sigma^2 n} \right)^{d/2} \left[ e^{-dx^2/2} + \mathcal{O}(R^{-\kappa}) + \mathcal{O}(n^{-\gamma^-}) + \mathcal{O}(n^{-\delta}) + \mathcal{O}(R^2 n^{-1}) \right]. \tag{4.23}$$

For  $x^2$  less than a sufficiently small multiple of  $\log R$ , this has the desired asymptotic form, provided  $R = R_n \rightarrow \infty$  as  $n \rightarrow \infty$ , with  $R_n = o(n^{1/2})$ .  $\square$

**A. Appendix: Example of  $D_L$**

In this Appendix, we verify that the function  $D_L$  defined in (1.22) obeys the requirements of Assumption D. The bounds of (1.17) and (1.18) follow easily, and clearly  $\hat{D}_L(0) = 1$ . We therefore concentrate on establishing (1.19) and (1.20)–(1.21). Before proceeding, we first note that it is sufficient to prove the lower bounds of (1.19) and (1.20) respectively for  $\|k\|_\infty \leq bL^{-1}$  and  $\|k\|_\infty \geq bL^{-1}$ , for any small positive  $b$ , since together these bounds imply the corresponding statements with  $b = 1$ . We will prove this modified form of the bounds.

For the upper bound of (1.19), we use only the symmetry of  $D_L$  and the inequality  $1 - \cos t \leq \frac{1}{2}t^2$  to obtain

$$\begin{aligned} 0 \leq a_L(k) &= \sum_{x \in \mathbb{Z}^d} [1 - \cos(k \cdot x)] D_L(x) \leq \frac{1}{2} \sum_{x \in \mathbb{Z}^d} (k \cdot x)^2 D_L(x) \\ &= \frac{1}{2} k^2 \sigma^2 \quad (k \in [-\pi, \pi]^d). \end{aligned} \tag{A.1}$$

For a lower bound, we use  $1 - \cos t \geq \frac{1}{2}t^2 - \text{const.}t^{2+2\epsilon}$ , (1.17) and (1.22) to conclude that there is a small positive  $b$  such that

$$a_L(k) \geq \frac{1}{2} k^2 \sigma^2 - \text{const.}k^{2+2\epsilon} L^{2+2\epsilon} \geq \text{const.}L^2 k^2 \quad (\|k\|_\infty \leq bL^{-1}). \tag{A.2}$$

For (1.20)–(1.21), it suffices to show that  $\hat{D}_L(k)$  is bounded away from 1 for  $\|k\|_\infty \geq bL^{-1}$  and bounded away from  $-1$  for  $k \in [-\pi, \pi]^d$ . Let  $\tilde{h}(k) = \int_{\mathbb{R}^d} h(x) e^{ik \cdot x} d^d x$  denote the Fourier integral transform of  $h$ . Then it suffices to show that

$$|\hat{D}_L(k/L) - \tilde{h}(k)| \rightarrow 0 \tag{A.3}$$

uniformly in  $k \in [-L\pi, L\pi]^d$ , since the characteristic function  $\tilde{h}$  of an almost everywhere continuous probability density is bounded away from  $-1$ , and is bounded away from 1 on the complement of any compact neighbourhood of 0. In the remainder of the proof, we prove (A.3).

Let  $X_L$  be a discrete random variable with probability mass function

$$\mathbb{P}(X_L = z) = D_L(Lz) = \frac{h(z)}{\sum_{w \in L^{-1}\mathbb{Z}^d} h(w)} \quad (z \in L^{-1}\mathbb{Z}^d). \tag{A.4}$$

Then  $X_L$  has characteristic function  $\hat{D}_L(k/L)$ . By definition,

$$\hat{D}_L(k/L) = \frac{\sum_{z \in L^{-1}\mathbb{Z}^d} h(z)e^{ik \cdot z}}{\sum_{z \in L^{-1}\mathbb{Z}^d} h(z)}. \tag{A.5}$$

The right side involves Riemann sums, and hence

$$\lim_{L \rightarrow \infty} \hat{D}_L(k/L) = \tilde{h}(k) \quad \text{for } k = \mathcal{O}(1). \tag{A.6}$$

We want to extend (A.6) to uniform convergence for  $k \in [-L\pi, L\pi]^d$ .

Given  $y \in \mathbb{R}^d$ , let  $z_y$  denote the closest point in  $L^{-1}\mathbb{Z}^d$  to  $y$ , with a fixed arbitrary rule used to choose from among the closest points when there is not a unique closest point. Let  $Y_L$  be a continuous random variable with density  $h_L$  given by

$$h_L(y) = \frac{L^d h(z_y)}{\sum_{z \in L^{-1}\mathbb{Z}^d} h(z)} \quad (y \in \mathbb{R}^d). \tag{A.7}$$

Thus  $Y_L$  has a density which is a piecewise constant analogue of the probability mass function of  $X_L$ . It follows that  $Y_L$  has the same distribution as  $X_L + U_L$ , where  $U_L$  is uniform on  $[-\frac{1}{2L}, \frac{1}{2L}]^d$  and independent of  $X_L$ . Let  $\tilde{h}_L$  and  $\tilde{u}_L$  respectively denote the characteristic functions of  $Y_L$  and  $U_L$ . Then

$$\hat{D}_L(k/L) = \frac{\tilde{h}_L(k)}{\tilde{u}_L(k)} \quad (k \in \mathbb{R}^d). \tag{A.8}$$

We claim that

$$|\tilde{h}_L(k) - \tilde{h}(k)| \rightarrow 0, \quad \text{uniformly in } k \in \mathbb{R}^d. \tag{A.9}$$

To see this, we fix  $\epsilon > 0$  and choose  $R$ , independent of  $L$ , such that we have  $\int_{\mathbb{R}^d \setminus [-R, R]^d} |h(y)| d^d y < \epsilon$  and  $\int_{\mathbb{R}^d \setminus [-R, R]^d} |h_L(y)| d^d y < \epsilon$ . This is possible since  $h$  is integrable, using the monotonicity assumption on  $H$ . Since  $h$  is almost everywhere continuous, for almost all  $y \in \mathbb{R}^d$  we have

$$\lim_{L \rightarrow \infty} h_L(y) = h(y). \tag{A.10}$$

Therefore, by dominated convergence,  $\int_{[-R, R]^d} |h(y) - h_L(y)| d^d y < \epsilon$  for  $L$  sufficiently large. The claim (A.9) then follows from the inequality  $|\tilde{h}_L(k) - \tilde{h}(k)| \leq \|h_L - h\|_1$ .

By definition,

$$\tilde{u}_L(k) = L^d \int_{[-\frac{1}{2L}, \frac{1}{2L}]^d} e^{ik \cdot x} d^d x = \prod_{i=1}^d \frac{2L \sin(\frac{k_i}{2L})}{k_i}. \tag{A.11}$$

Therefore, for any fixed  $\alpha > 0$ ,  $\tilde{u}_L(k) \rightarrow 1$  uniformly in  $k$  such that  $\|k\|_\infty \leq L^{1-\alpha}$ . With (A.8) and the uniform convergence of (A.9), this proves that the convergence in (A.3) holds uniformly in  $k$  such that  $\|k\|_\infty \leq L^{1-\alpha}$ .

It remains to consider the case  $\|k\|_\infty \geq L^{1-\alpha}$ . By (A.11),  $\tilde{u}_L(k) \geq (2\pi^{-1})^d$  uniformly in  $k \in [-L\pi, L\pi]^d$ . Also, by the Riemann–Lebesgue lemma,  $\lim_{L \rightarrow \infty} \sup_{\{k: \|k\|_\infty \geq L^{1-\alpha}\}} \tilde{h}(k) = 0$ . Therefore, by (A.8) and (A.9),

$$|\hat{D}(k/L) - \tilde{h}(k)| \leq \left(\frac{\pi}{2}\right)^d |\tilde{h}_L(k)| + |\tilde{h}(k)| \rightarrow 0 \quad (\text{A.12})$$

uniformly in  $k$  such that  $\|k\|_\infty \geq L^{1-\alpha}$ . This completes the proof of (1.20)–(1.21).

*Acknowledgements.* This work was supported in part by NSERC of Canada. The work of both authors was carried out in part at Microsoft Research and the Fields Institute. The work of G.S. was also carried out in part at McMaster University. We are grateful to an anonymous referee for numerous helpful suggestions.

## References

1. Aizenman, M., Barsky, D.J.: Sharpness of the phase transition in percolation models, *Commun. Math. Phys.*, **108**, 489–526 (1987)
2. Aizenman, M., Newman, C.M.: Tree graph inequalities and critical behavior in percolation models, *J. Stat. Phys.*, **36**, 107–143 (1984)
3. Barsky, D.J., Aizenman, M.: Percolation critical exponents under the triangle condition, *Ann. Probab.*, **19**, 1520–1536 (1991)
4. Bezuidenhout, C., Grimmett, G.: The critical contact process dies out, *Ann. Probab.*, **18**, 1462–1482 (1990)
5. Bolthausen, E., Ritzmann, C.: Strong pointwise estimates for the weakly self-avoiding walk, Preprint, (2001)
6. Brydges, D.C., Spencer, T.: Self-avoiding walk in 5 or more dimensions, *Commun. Math. Phys.*, **97**, 125–148 (1985)
7. Golowich, G.R., Imbrie, J.Z.: A new approach to the long-time behavior of self-avoiding random walks, *Ann. Phys.*, **217**, 142–169 (1992)
8. Grimmett, G.R. Hiemer, P.: Directed percolation and random walk, Preprint, (2001)
9. Hara, T., van der Hofstad, R., Slade, G.: Critical two-point functions and the lace expansion for spread-out high-dimensional percolation and related models, Preprint, (2001)
10. Hara, T., Slade, G.: Mean-field critical behaviour for percolation in high dimensions, *Commun. Math. Phys.*, **128**, 333–391 (1990)
11. Hara, T., Slade, G.: The lace expansion for self-avoiding walk in five or more dimensions, *Reviews in Math. Phys.*, **4**, 235–327 (1992)
12. van der Hofstad, R.: The lace expansion approach to ballistic behaviour for one-dimensional weakly self-avoiding walks, *Probab. Th. Rel. Fields* **119**, 311–349 (2001)
13. van der Hofstad, R., den Hollander, F., Slade, G.: A new inductive approach to the lace expansion for self-avoiding walks, *Probab. Th. Rel. Fields*, **111**, 253–286 (1998)
14. van der Hofstad, R., Slade, G.: Convergence of critical oriented percolation to super-Brownian motion above  $4 + 1$  dimensions, Preprint (2001)
15. van der Hofstad, R., Slade, G.: The lace expansion on a tree with application to networks of self-avoiding walks, Preprint, (2001)

16. Khanin, K.M., Lebowitz, J.L., Mazel, A.E., Sinai, Ya.G.: Self-avoiding walks in five or more dimensions: polymer expansion approach, *Russian Math. Surveys*, **50**, 403–434 (1995)
17. Madras, N., Slade, G.: *The Self-Avoiding Walk*, Birkhäuser, Boston, (1993)
18. Menshikov, M.V.: Coincidence of critical points in percolation problems, *Soviet Mathematics, Doklady*, **33**, 856–859 (1986)
19. Nguyen, B.G., Yang, W-S.: Triangle condition for oriented percolation in high dimensions, *Ann. Probab.*, **21**, 1809–1844 (1993)
20. Nguyen, B.G., Yang, W-S.: Gaussian limit for critical oriented percolation in high dimensions. *J. Stat. Phys.*, **78**, 841–876 (1995)
21. Penrose, M.D.: Self-avoiding walks and trees in spread-out lattices, *J. Stat. Phys.*, **77**, 3–15 (1994)
22. Sakai, A.: Mean-field critical behavior for the contact process, *J. Stat. Phys.* **104**, 111–143 (2001)
23. Slade, G.: The diffusion of self-avoiding random walk in high dimensions, *Commun. Math. Phys.*, **110**, 661–683 (1987)
24. Slade, G.: The lace expansion and the upper critical dimension for percolation, *Lectures in Applied Mathematics*, **27**:53–63, (1991). (Mathematics of Random Media, eds. W.E. Kohler and B.S. White, A.M.S., Providence).