

A construction of the Suzuki graph

A.E. Brouwer

2008-07-08

Abstract

We give an elementary construction of the rank 3 Suzuki graph, following Horiguchi, Kitazume & Nakasora.

1 Introduction

The Suzuki sporadic simple group Suz was discovered by M. Suzuki [3] as index 2 subgroup of the automorphism group of a rank 3 graph Σ , strongly regular with parameters $(v, k, \lambda, \mu) = (1782, 416, 100, 96)$, the largest graph in the Suzuki tower (of graphs for $\text{Suz}:2$, $G_2(4):2$, $\text{HJ}:2$, $U_3(3):2$, $L_3(2):2$). The spectrum of Σ is $416^1, 20^{780}, (-16)^{1001}$.

The Hoffman bound for the size of a coclique in Σ is 66, and in fact this graph has a single orbit of cocliques of this size, as was found by Kuzuta [2] and is easy to check.

If C is a 66-coclique in Σ , then each point outside C has 16 neighbours in C , so that the 1716 points outside C induce the structure of a block design with parameters 2 -(66,16,96) on C . It was observed by Horiguchi, Kitazume & Nakasora [1], that this block design happens to be a 3 -(66,16,21).

Something funny happens here: usually when one has a maximal coclique in a graph belonging to a nice group, then the group is transitive on the vertices of the coclique. But here the stabilizer of C in $\text{Suz}:2$ is $U_3(4):4$ acting with vertex orbit sizes $1 + 65$, 2-transitively on the orbit of length 65.

In [1] the graph Σ was constructed in this setting, starting from designs on the unital in $PG(2, 16)$. We repeat their construction here, perhaps in a more elegant way.

2 A 2-(65,15,21) design and the $G_2(4)$ graph

Consider the vector space V of dimension 3 over the field \mathbf{F}_{16} , provided with a nondegenerate Hermitean form $\langle \cdot, \cdot \rangle$. The projective plane PV has 65 isotropic and 208 nonisotropic points. There are $208 \cdot 12.1/6 = 416$ orthogonal bases (triples of orthogonal nonisotropic points).

We would like to construct the local graph Γ of Σ , a strongly regular graph with parameters $(v, k, \lambda, \mu) = (416, 100, 36, 20)$. The group $U_3(4):4$ of semilinear transformations preserving the form acts transitively on the 416 bases, with rank 5. The suborbit sizes are 1, 15, 100, 150, 150. The graph Γ is the graph on the 416 bases obtained by taking the suborbit of size 100 for adjacency.

These suborbits can be described geometrically as follows: Given one basis $\{a, b, c\}$, the suborbit of size 15 consists of the bases that have an element in common with $\{a, b, c\}$. The first suborbit of size 150 consists of the bases that are disjoint from $\{a, b, c\}$ but contain a point orthogonal to one of a, b, c . Associated with a basis $\{a, b, c\}$ is the triangle consisting of the 15 isotropic points on the three lines $ab, ac,$ and bc . The suborbits of sizes 1, 15, 100, 150, 150 correspond to the bases with triangles having 15, 5, 3, 2, 5 points in common, respectively. Thus, Γ can be described as the graph on the 416 triangles, adjacent when they have 3 points in common ([1]).

These 416 triangles form a 2-(65,15,21) design.

3 A 3-(66,16,21) design and the Suz graph

The set of 16 hyperbolic lines on any isotropic point y carries the structure of an affine plane of order 4. This affine plane has 20 lines of size 4, so we find 20 sets of four lines on y , and taking only the isotropic points distinct from y on these lines, we find for each y 20 sets of 16 isotropic points, and altogether 1300 such sets. Since $U_3(4):4$ acts 2-transitively these 1300 sets will be the blocks of a 2-design with parameters 2-(65,16,75).

If we add a point ∞ to the set of isotropic points, and to the blocks (triangles) of the 2-(65,15,21) design found earlier, then the union of both designs is clearly a 2-(66,16,96), and in fact even a 3-(66,16,21) design. Indeed, the group is transitive on collinear triples of isotropic points and on noncollinear such triples so only a single equality needs to be checked.

Let (X, \mathcal{B}) be the 3-(66,16,21) design just constructed, with 66 points and 1716 blocks. It has block intersection numbers 0, 3, 4, 6, 16, and each block meets n_i blocks in i points, where $n_0 = 75, n_3 = 800, n_4 = 400, n_6 = 440, n_{16} = 1$.

Now construct the Suz graph with $1782 = 66 + 1716 = 1 + 65 + 416 + 1300$ vertices, where the $1 + 65$ is a 66-coclique, the 416 are the neighbours of 1 and the $416 + 1300$ (viewed as blocks) are each adjacent to their 16 points. We already saw that "having 3 points in common" was the right adjacency for the 15-point triangles, and now that ∞ has been added the construction becomes: let two blocks be adjacent when they have 4 points in common. This defines the sporadic Suzuki graph.

References

- [1] N. Horiguchi, M. Kitazume & H. Nakasora, *A construction of the sporadic Suzuki graph from $U_3(4)$* , preprint, 2008.
- [2] K. Kuzuta, Master Thesis, Chiba Univ., 2007 (in Japanese).
- [3] M. Suzuki, *A simple group of order 448,345,497,600*, pp. 113–119 in: *Theory of Finite Groups (Symposium, Harvard Univ., Cambridge, Mass, 1968)*, Benjamin, New York, 1969.