

EXAMINATION MATHEMATICAL TECHNIQUES FOR IMAGE ANALYSIS

Course code: 8D020. Date: Thursday April 07, 2011. Time: 14h00–17h00. Place:

Read this first!

- Use a separate sheet of paper for each problem. Write your name and student ID on each paper.
- The exam consists of 4 problems. The maximum credit for each item is indicated in the margin.
- Motivate your answers. The use of course notes is allowed. The use of problem companion (“opgaven- en tentamenbundel”), calculator, laptop, or other equipment, is *not* allowed.
- You may provide your answers in Dutch or English.

GOOD LUCK!

(30) 1. CLIFFORD ALGEBRA

Attention: For each numeric symbol (matrix entry, scalar multiplier, et cetera) that you use in your arguments below please state explicitly whether it is real or complex.

Let σ be any 2×2 matrix with complex entries $\sigma_{ij} \in \mathbb{C}$ in i -th row and j -th column. The set of all complex 2×2 matrices constitutes a vector space over the *real* numbers (i.e. scalar multiplication pertains to *real* scalars), which we indicate here by $\mathbb{M}_{2 \times 2}$. For simplicity of notation we write the identity matrix as $1 \in \mathbb{M}_{2 \times 2}$. The complex conjugate of a complex number $z = a + bi$, $a, b \in \mathbb{R}$, is indicated by $z^* = a - bi$.

The *trace* operator $\text{tr} : \mathbb{M}_{2 \times 2} \rightarrow \mathbb{C}$ is defined by $\text{tr} \sigma = \sigma_{11} + \sigma_{22}$.

The *hermitian conjugate* operator $\dagger : \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ is defined by $\begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{pmatrix}^\dagger = \begin{pmatrix} \sigma_{11}^* & \sigma_{21}^* \\ \sigma_{12}^* & \sigma_{22}^* \end{pmatrix}$.

- (10) a. Show that $\text{tr} : \mathbb{M}_{2 \times 2} \rightarrow \mathbb{C} : \sigma \mapsto \text{tr} \sigma$ and $\dagger : \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2} : \sigma \mapsto \sigma^\dagger$ are linear operators.

The three so-called Pauli matrices are defined as follows:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Together they define a basis for a 3-dimensional vector space $V \subset \mathbb{M}_{2 \times 2}$ over the real numbers.

- (5) b. What is the general form of an element $\sigma \in V$?
- (5) c. Show that $\text{tr} \sigma = 0$ and $\sigma^\dagger = \sigma$ for all $\sigma \in V$.

We subsequently enrich the vector space V with a multiplication operator, viz. standard matrix multiplication.

- (5) **d.** Compute all nine products of the form $\sigma_k \sigma_\ell$ for $k, \ell = 1, 2, 3$.

We now consider the set A consisting of all real linear combinations of *all possible* products (i.e. with an arbitrary number of factors) of Pauli matrices. In this construct we allow for the “empty” product, and define it to produce the identity matrix $1 \in \mathbb{M}_{2 \times 2}$.

- (5) **e.** Interpreted as a vector space over the real numbers, show that A has dimension 8 by providing an explicit set of 8 basis vectors.
(*Hint:* $\dim \mathbb{M}_{2 \times 2} = 8$.)



(25) 2. STAIRCASE FUNCTION

We introduce the staircase function $f : \mathbb{R} \rightarrow \mathbb{R} : x \mapsto f(x)$ given by $f(x) = \lfloor x \rfloor$, i.e. the so-called *entier* of $x \in \mathbb{R}$, which is defined as the largest integer $k \in \mathbb{Z}$ such that $k \leq x$.

- (5) **a.** Sketch the graph of $y = f(x)$ on the interval $[-3, 3]$, clearly illustrating its discontinuities.
(5) **b.** Give the formula for the classical derivative $f'(x)$, together with its domain of definition.

The regular tempered distribution $T_g : \mathcal{S}(\mathbb{R}) \rightarrow \mathbb{R}$ associated with a function $g : \mathbb{R} \rightarrow \mathbb{R}$ is the tempered distribution given by

$$T_g(\phi) = \int_{-\infty}^{\infty} g(x) \phi(x) dx ,$$

for any $\phi \in \mathcal{S}(\mathbb{R})$.

- (5) **c.** Show that, in distributional sense, $T'_f \neq T_{f'}$ for the staircase function f defined above.

For any $a \in \mathbb{R}$ we furthermore define the (shifted) Dirac distribution $\delta_a : \mathcal{S}(\mathbb{R}) \rightarrow \mathbb{R}$ by

$$\delta_a(\phi) = \phi(a) ,$$

for any $\phi \in \mathcal{S}(\mathbb{R})$.

- (10) **d.** Prove that $T'_f = \sum_{k \in \mathbb{Z}} \delta_k$.



(20) **3. GROUP THEORY**

A discrete group G_n with n distinct elements $x_i \in G_n, i = 1, \dots, n$, can be represented by means of a multiplication table. The term multiplication refers to the group operation, which will be denoted by the infix operator $\circ : G \times G \rightarrow G : (x_i, x_j) \mapsto x_i \circ x_j$. The element on i -th row and j -th column in the table specifies the product $x_i \circ x_j \in G$:

\circ	x_1	\cdots	x_j	\cdots	x_n
x_1	$x_1 \circ x_1$	\cdots	$x_1 \circ x_j$	\cdots	$x_1 \circ x_n$
\vdots	\vdots		\vdots		\vdots
x_i	$x_i \circ x_1$	\cdots	$x_i \circ x_j$	\cdots	$x_i \circ x_n$
\vdots	\vdots		\vdots		\vdots
x_n	$x_n \circ x_1$	\cdots	$x_n \circ x_j$	\cdots	$x_n \circ x_n$

Of course, one obtains full knowledge about the group if all entries in this table are provided. However, due to the specific structure of a group it is not necessary to provide all entries in order to uniquely specify the group. An incomplete table may be completed or partially completed with the help of the defining group properties.

- (10) **a.** Prove that elements in any given row are all distinct. Likewise for elements in any given column.

We now consider the specific case of $G_4 = \{E, A, B, C\}$. This group has the following properties:

- E is the identity element,
- B equals its own inverse,
- $A \circ A = C \circ C \neq E$.

- (10) **b.** Complete the multiplication table for G_4 , and explain how you obtained your result:

\circ	E	A	B	C
E				
A				
B				
C				



(25) 4. FOURIER THEORY¹

Consider the following initial value problem for the function $u : \mathbb{R}^n \times \mathbb{R}^+ \rightarrow \mathbb{R}$:

$$\begin{cases} \frac{\partial u(x, t)}{\partial t} = (\Delta - m^2) u(x, t) & \text{for } (x, t) \in \mathbb{R}^n \times \mathbb{R}^+ \\ u(x, 0) = f(x) & \text{for } x \in \mathbb{R}^n \end{cases}$$

Here $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a real valued image and $m > 0$ a positive constant. Boundary and initial conditions are such that this initial value problem has a unique, sufficiently nice solution.

In this problem the following Fourier convention applies. For (sufficiently nice) $u : \mathbb{R}^n \rightarrow \mathbb{C}$ we define the function $\widehat{u} = \mathcal{F}(u) : \mathbb{R}^n \rightarrow \mathbb{C}$ as follows:

$$\widehat{u}(\omega) = \int_{\mathbb{R}^n} e^{-i\omega \cdot x} u(x) dx \quad \text{or, equivalently,} \quad u(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\omega \cdot x} \widehat{u}(\omega) d\omega,$$

in which $\omega \cdot x$ denotes $\omega_1 x_1 + \dots + \omega_n x_n$. For the sake of brevity we furthermore write $\|x\|^2 = x \cdot x$, respectively $\|\omega\|^2 = \omega \cdot \omega$.

In the problems below you may use the standard integral

$$\int_{-\infty}^{\infty} e^{-(x+iy)^2} dx = \sqrt{\pi} \quad \text{irrespective of the value of } y \in \mathbb{R}.$$

- (5) a. Show that the initial value problem above is equivalent to the following initial value problem in the Fourier domain:

$$\begin{cases} \frac{d\widehat{u}(\omega, t)}{dt} = -(\|\omega\|^2 + m^2) \widehat{u}(\omega, t) & \text{voor } (\omega, t) \in \mathbb{R}^n \times \mathbb{R}^+ \\ \widehat{u}(\omega, 0) = \widehat{f}(\omega) & \text{for } \omega \in \mathbb{R}^n \end{cases}$$

(Note that we may interpret this as an *ordinary* differential equation with initial condition, whence the alternative notation for the t -derivative on the left hand side.)

- (5) b. Find the solution $\widehat{u}(\omega, t)$.
(*Hint:* Stipulate a solution of type $\widehat{u}(\omega, t) = Ae^{Bt}$ and determine the (ω -dependent) parameters A, B .)

- (5) c. Show that for fixed $t \in \mathbb{R}^+$ the spatial solution is given by the convolution product

$$u(x, t) = (\phi_t * f)(x)$$

for some convolution filter $\phi_t : \mathbb{R}^n \rightarrow \mathbb{R}$.

- (5) d. Determine the form $\phi_t(x)$ of this convolution filter.

- (5) e. Prove: $\int_{\mathbb{R}^n} \phi_t(x) dx = e^{-m^2 t}$.
(*Hint:* Consider $\widehat{\phi}_t(\omega)$.)

THE END

¹Exam June 14, 2005, problem 3.