

An ω -Complete Equational Specification of Interleaving ^{*} Extended Abstract

W.J. Fokkink¹ and S.P. Luttik^{1,2,**}

¹ CWI, P.O. Box 94079, NL-1090 GB Amsterdam, The Netherlands

Wan.Fokkink@cwi.nl

² Department of Computer Science, University of Amsterdam

Kruislaan 403, NL-1098 SJ Amsterdam, The Netherlands

Bas.Luttik@cwi.nl

Abstract. We consider the process theory PA that includes an operation for parallel composition, based on the interleaving paradigm. We prove that the standard set of axioms of PA is not ω -complete by providing a set of axioms that are valid in PA, but not derivable from the standard ones. We prove that extending PA with this set yields an ω -complete specification, which is finite in a setting with finitely many actions.

1 Introduction

The interleaving paradigm consists of the assumption that two atomic actions cannot happen at the same time, so that concurrency reduces to nondeterminism. To express the concurrent execution of processes, many process theories have been accommodated with an operation for parallel composition that behaves according to the interleaving paradigm. For instance, CCS (see, e.g., Milner (1989)) has a binary operation for parallel composition —we shall denote it by $_||_-$ — that satisfies the so-called *Expansion Law*:

$$\text{if } p = \sum_{i=1}^m a_i.p_i \text{ and } q = \sum_{j=1}^n b_j.q_j, \text{ then } p || q \approx \sum_{i=1}^m a_i.(p_i || q) + \sum_{j=1}^n b_j.(q_j || p);$$

here the $a_i._-$ and the $b_j._-$ are unary operations that prefix a process with an atomic action, and summation denotes a nondeterministic choice between its arguments.

The Expansion Law generates an infinite set of equations, one for each pair of processes p and q . Bergstra and Klop (1984) enhanced the equational characterisation of interleaving. They replaced action prefixing with a binary operation

^{*} A full version that contains the omitted proofs is available as CWI Technical Report; see <http://www.cwi.nl/~luttik/>

^{**} Research supported by the Netherlands Organization for Scientific Research (NWO) under contract SION 612-33-008.

\cdot for sequential composition and added an auxiliary operation \parallel (the *left merge*; it is similar to \parallel , except that it must start execution with a step from its left argument). Their axiomatisation is finite for settings with finitely many atomic actions. Moller (1990) proved that interleaving is not finitely axiomatisable without an auxiliary operation such as the left merge.

The axioms of Bergstra and Klop (1984) form a ground-complete axiomatisation of bisimulation equivalence; ground terms p and q are provably equal if, and only if, they are bisimilar. Thus, it reflects for a large part our intuition about interleaving. On the other hand, it is not optimal. For instance, it can be shown by means of structural induction that every ground instance of the axiom $x \parallel (y \parallel z) \approx (x \parallel y) \parallel z$ is derivable (see Baeten and Weijland (1990)); however, the axiom itself is not derivable.

If an equational specification E has the property that $E \vdash t^\sigma \approx u^\sigma$ for all ground substitutions σ implies that $E \vdash t \approx u$, then E is called ω -complete (or: *inductively closed*). To derive any equation from such an equational specification it is never needed to use additional proof techniques such as structural induction. Therefore, in applications dealing with theorem proving, ω -completeness is a desirable property to have (see Lazrek *et al.* (1990)). In Heering (1986) it was argued that ω -completeness is desirable for the partial evaluation of programs.

Moller (1989) obtained an ω -complete axiomatisation for CCS without communication, by adding a law for *standard concurrency*:

$$(x \parallel y) \parallel z \approx x \parallel (y \parallel z).$$

In this paper we shall address the question whether PA, the subtheory of ACP without communication and encapsulation, is ω -complete. While the algebra studied by Moller (1989) has sequential composition in the form of prefix multiplication, PA incorporates the (more general) binary operation \cdot for sequential composition. Having this operation, it is no longer sufficient to add the law for standard concurrency to arrive at an ω -complete axiomatisation. However, surprisingly, it is sufficient to add this law and the set of axioms generated by a single scheme:

$$(x \cdot \alpha \parallel \alpha) \approx (x \parallel \alpha) \cdot \alpha,$$

where α ranges over alternative compositions of distinct atomic actions; if the set of atomic actions is finite, then this scheme generates finitely many axioms.

An important part of our proof has been inspired by the excellent work of Hirshfeld and Jerrum (1999) on the decidability of bisimulation equivalence for normed process algebra. In particular, they distinguish two kinds of *mixed equations*, in which a parallel composition is equated to a sequential composition. The first kind consists of equations

$$(t \cdot \alpha^k) \parallel \alpha^l \approx (t \parallel \alpha^l) \cdot \alpha^k$$

for positive natural numbers k and l , and for sums of atomic actions α . These equations can be derived using standard concurrency and our new axioms. The

second kind of mixed equations are the so-called *pumpable equations*, which are of a more complex nature (see p. 419 of Hirshfeld and Jerrum (1999)). Basically, we show that there cannot exist pumpable equations that contain variables by associating with every candidate $t \approx u$ a ground substitution σ such that $t^\sigma \not\approx u^\sigma$.

The notion of ω -completeness is related to *action refinement*, where each atomic action may be refined to an arbitrary process. That is, in a theory with action refinement, the actions take over the role played by variables in our theory; the actions, as they occur in our theory, are not present in theories for action refinement. Aceto and Hennessy (1993) presented a complete axiomatisation for PA (including a special constant `nil`, being a hybrid of deadlock and empty process) with action refinement, modulo timed observational equivalence from Hennessy (1988). In this setting, laws such as $a \parallel x \approx a \cdot x$, which hold in standard PA, are no longer valid, as the atomic action a can be refined into any other process.

This paper is set up as follows. In §2 we introduce the standard axioms of interleaving, and we prove that they do not form an ω -complete specification by proving that all ground substitution instances of our new axioms are derivable, while the axioms themselves are not. In §3 we state some basic facts about the theory of interleaving that we shall need in our proof of ω -completeness. In §4 we collect some results on certain mixed equations, and in §5 we investigate a particular kind of terms that consist of nestings of parallel and sequential compositions. In §6 we prove our main theorem, that the standard theory of interleaving enriched with the law for standard concurrency and our new axioms is ω -complete.

2 Interleaving

A *process algebra* is an algebra that satisfies the axioms A1–A5 of Table 1. Suppose that A is a set of constant symbols and suppose that \parallel and \ll are binary operation symbols; a process algebra with interpretations for the constant symbols in A and the operations \parallel and \ll satisfying M1, M4, M5, M2 _{a} and M3 _{a} for all $a \in A$, and M6 _{α} for all sums of distinct elements of A , we shall call an *A-merge algebra*; the variety of *A-merge algebras* we denote by PA _{A} .

Table 1. The axioms of PA _{A} , with $a \in A$ and α any sum of distinct elements of A .

(A1) $x + y$	$\approx y + x$	(M1) $x \parallel y$	$\approx x \ll y + y \ll x$
(A2) $x + (y + z)$	$\approx (x + y) + z$	(M2 _{a}) $a \ll x$	$\approx a \cdot x$
(A3) $x + x$	$\approx x$	(M3 _{a}) $a \cdot x \ll y$	$\approx a \cdot (x \parallel y)$
(A4) $(x + y) \cdot z$	$\approx x \cdot z + y \cdot z$	(M4) $(x + y) \ll z$	$\approx x \ll z + y \ll z$
(A5) $(x \cdot y) \cdot z$	$\approx x \cdot (y \cdot z)$	(M5) $(x \ll y) \ll z$	$\approx x \ll (y \parallel z)$
		(M6 _{α}) $x \cdot \alpha \ll \alpha$	$\approx (x \ll \alpha) \cdot \alpha$

The axioms A1–A5 together with the axioms M1–M4 form the standard axiomatisation of interleaving. Consider the single-sorted signature Σ with the elements of A as constants and the binary operations $+$, \cdot , \ll and \parallel . In writing terms we shall often omit the operation \cdot for sequential composition; we assume that sequential composition binds strongest and that the operation $+$ for alternative composition binds weakest.

Let \mathcal{R} consist of the axioms A3–A5 and M1–M4 of Table 1 interpreted as rewrite rules by orienting them from left to right. The term rewriting system $\langle \Sigma, \mathcal{R} \rangle$ is ground terminating and ground confluent modulo associativity and commutativity of $+$ (cf. the axioms A1 and A2). A ground term t is a *basic term* if there exist disjoint finite sets I and J , elements a_i and b_j of A and basic terms t_i , for $i \in I$ and $j \in J$ such that

$$t \approx \sum_{i \in I} a_i t_i + \sum_{j \in J} b_j \quad (\text{by A1 and A2}).$$

Every ground normal form of $\langle \Sigma, \mathcal{R} \rangle$ is a basic term.

It is well-known that the axioms A1–A5 together with M1–M4 do not constitute an ω -complete axiomatisation; all ground substitution instances of M5 are derivable, while the axiom itself is not. Moller (1989) has shown that, in a setting with prefix sequential composition instead of the binary operation \cdot , it suffices to add M5 to obtain an ω -complete axiomatisation (see Groote (1990) for an alternative proof). Clearly, neither $x\alpha \ll \alpha$ nor $(x \ll \alpha)\alpha$ is an instance of any of the axioms A1–A5 and M1–M5, so $M6_\alpha$ is not derivable. However, each ground substitution instance of $M6_\alpha$ is derivable.

Proposition 1. *If α is a finite sum of elements of A , then, for every ground term t ,*

$$A1, \dots, A5, M1, \dots, M4 \vdash t\alpha \ll \alpha \approx (t \ll \alpha)\alpha.$$

Consequently, in the case of binary sequential composition, the axioms A1–A5 together with M1–M5 do not constitute an ω -complete axiomatisation. In the sequel, we shall prove that PA_A is ω -complete.

3 Basic Facts

In every A -merge algebra, $+$ and \parallel are commutative and associative; we shall often implicitly make use of this. Also, we shall frequently abbreviate the statement $PA_A \vdash u \approx u + t$ by $t \preceq u$; if $t \preceq u$, then we call t a *summand* of u . Note that \preceq is a partial order on the set of terms modulo \approx ; in particular, if $t \preceq u$ and $u \preceq t$, then $t \approx u$.

Lemma 2. *Let a be an element of A and let t , u and v be ground terms. If $at \preceq u + v$, then $at \preceq u$ or $at \preceq v$.*

Suppose t is a ground normal form of the system $\langle \Sigma, \mathcal{R} \rangle$ and suppose that

$$t \approx \sum_{i \in I} a_i t_i + \sum_{j \in J} b_j;$$

then the *degree* $d(t)$ of t is defined by $d(t) = |I| + |J|$. We let the degree of an arbitrary ground term be the degree of its unique normal form in $\langle \Sigma, \mathcal{R} \rangle$. By $d_{\max}(t)$ we shall denote the maximal degree that occurs in t , i.e.,

$$d_{\max}(t) = \max(\{d(t)\} \cup \{d_{\max}(t') \mid \text{there exists an } a \in A \text{ such that } at' \approx t\}).$$

Lemma 3. *If α is a finite sum of elements of A , then*

$$\text{PA}_A \vdash x\alpha \parallel \alpha^n \approx (x \parallel \alpha^n)\alpha, \text{ and } \text{PA}_A \vdash x\alpha \parallel \alpha^n \approx (x \parallel \alpha^n)\alpha.$$

Proof. It is straightforward to show by induction on n that the identity (*) $\alpha^{n+1} \approx \alpha^n \parallel \alpha$ is derivable from PA_A ; we shall use it in the proof of the first set of equations (**) $x\alpha \parallel \alpha^n \approx (x \parallel \alpha^n)\alpha$, which is by induction on n . If $n = 1$, then (**) is an instance of M6_α , and for the induction step we have the following derivation:

$$\begin{aligned} x\alpha \parallel \alpha^{n+1} &\approx x\alpha \parallel (\alpha^n \parallel \alpha) && \text{(by *)} \\ &\approx (x\alpha \parallel \alpha^n) \parallel \alpha && \text{(by M5)} \\ &\approx (x \parallel \alpha^n)\alpha \parallel \alpha && \text{(by IH)} \\ &\approx ((x \parallel \alpha^n) \parallel \alpha)\alpha && \text{(by M6}_\alpha\text{)} \\ &\approx (x \parallel (\alpha^n \parallel \alpha))\alpha && \text{(by M5)} \\ &\approx (x \parallel \alpha^{n+1})\alpha && \text{(by *)}. \end{aligned}$$

The second set of equations is also derived by induction on n , using (**). □

Milner and Moller (1993) proved that if t, u and v are ground terms such that $t \parallel v$ and $u \parallel v$ are bisimilar, then t and u are bisimilar (a similar result was obtained earlier by Castellani and Hennessy (1989) in the context of distributed bisimulation). Also, they proved that every finite process has, up to bisimulation equivalence, a unique decomposition into prime components. Since PA_A is a sound and complete axiomatisation for bisimulation equivalence (Bergstra and Klop, 1984), the following two results are consequences of theirs.

Lemma 4. *If t, u and v are ground terms such that $\text{PA}_A \vdash t \parallel v \approx u \parallel v$, then $\text{PA}_A \vdash t \approx u$.*

Definition 5. *A ground term t we shall call parallel prime if there do not exist ground terms u and v such that $\text{PA}_A \vdash t \approx u \parallel v$.*

Theorem 6 (Unique factorisation). *Any ground term can be expressed uniquely as a parallel composition of parallel prime components.*

We associate to each term t a *norm* $\lfloor t \rfloor$ and a *depth* $\lceil t \rceil$ as follows:

$$\begin{aligned} \lfloor x \rfloor &= \lfloor a \rfloor = 1 & \lceil a \rceil &= \lceil x \rceil = 1 & (a \in A \text{ and } x \text{ a variable}); \\ \lfloor x * y \rfloor &= \lfloor x \rfloor + \lfloor y \rfloor & \lceil x * y \rceil &= \lceil x \rceil + \lceil y \rceil & \text{if } * \in \{\cdot, \parallel, \|\}; \text{ and} \\ \lfloor x + y \rfloor &= \min\{\lfloor x \rfloor, \lfloor y \rfloor\} & \lceil x + y \rceil &= \max\{\lceil x \rceil, \lceil y \rceil\}. \end{aligned}$$

Notice that if $t \approx u$, then t and u must have equal norm and depth.

Lemma 7. *If t, t', u and u' are ground terms such that $\lfloor t \rfloor = \lfloor t' \rfloor$, $\lfloor u \rfloor = \lfloor u' \rfloor$ and $\text{PA}_A \vdash tu \approx t'u'$, then $\text{PA}_A \vdash t \approx t'$ and $\text{PA}_A \vdash u \approx u'$.*

Definition 8. *Let t and t' be ground terms; we shall write $t \longrightarrow t'$ if there exists $a \in A$ such that $at' \preceq t$ and $\lfloor t' \rfloor < \lfloor t \rfloor$. We define the set $\text{red}(t)$ of reducts of t as the least set that contains t and is closed under \longrightarrow ; if $t \longrightarrow t'$, then we call t' an immediate reduct of t .*

Lemma 9. *Let t be a ground term. If $t \longrightarrow t'$ and $t \longrightarrow t''$ implies that $\text{PA}_A \vdash t' \approx t''$ for all ground terms t' and t'' , then there exists a parallel prime ground term t^* such that $\text{PA}_A \vdash t \approx t^* \parallel \dots \parallel t^*$.*

Proof. First, suppose that u and v are parallel prime, and let u' and v' be such that $u \longrightarrow u'$ and $v \longrightarrow v'$; then, $u \parallel v \longrightarrow u' \parallel v$ and $u \parallel v \longrightarrow u \parallel v'$. So, if $u' \parallel v \approx u \parallel v'$, then since $\lfloor u' \rfloor < \lfloor u \rfloor$, u cannot be a component of the prime decomposition of u' ; hence, by Theorem 6, $u \approx v$.

Suppose $t \approx t_1 \parallel \dots \parallel t_n$, with t_i parallel prime for all $1 \leq i \leq n$ and $\lfloor t_1 \rfloor \leq \dots \leq \lfloor t_n \rfloor$.

If $\lfloor t_1 \rfloor = 1$, then $\lfloor t_i \rfloor = 1$ for all $1 \leq i \leq n$; for suppose that t'_i is a ground term such that $t_i \longrightarrow t'_i$, then from $t_2 \parallel \dots \parallel t_n \approx t_1 \parallel \dots \parallel t_{i-1} \parallel t'_i \parallel t_{i+1} \parallel \dots \parallel t_n$, we get by Lemma 4 that $t_i \approx t_1 \parallel t'_i$, but t_i is parallel prime. From $t_1 \parallel \dots \parallel t_{i-1} \parallel t_{i+1} \parallel \dots \parallel t_n \approx t_1 \parallel \dots \parallel t_{j-1} \parallel t_{j+1} \parallel \dots \parallel t_n$, we conclude by Lemma 4 that $t_i \approx t_j$.

The remaining case is that $\lfloor t_i \rfloor > 1$ for all $1 \leq i \leq n$. Let t'_i and t'_j be ground terms such that $t_i \longrightarrow t'_i$ and $t_j \longrightarrow t'_j$ for some $1 \leq i < j \leq n$; then by Lemma 4 $t'_i \parallel t_j \approx t_i \parallel t'_j$. Since $\lfloor t'_i \rfloor < \lfloor t_i \rfloor$, t_i cannot be a component of the prime decomposition of t'_i , so by Theorem 6 $t_i \approx t_j$. □

4 Mixed Equations

We shall collect some results about *mixed equations*; these are equations of the form $tu \approx v \parallel w$.

Lemma 10. *If t, u and v are ground terms such that $\text{PA}_A \vdash tu \approx u \parallel v$, then there exists a finite sum α of elements of A such that $\text{PA}_A \vdash u \approx \alpha^k$ for some $k \geq 1$.*

Proof. Note that $[t] = [v]$; we shall first prove the following

Claim: if $[t], [v] = 1$, then there exists a $k \geq 1$ such that $u \approx t^k$ and $t \approx v$.

Let $t = a_1 + \dots + a_m$ with $a_1, \dots, a_m \in A$; we proceed by induction on $[u]$.

If $[u] = 1$, then there exists $a \in A$ such that $a \preceq u$, whence $at \preceq u \parallel v$. Since $a_1u + \dots + a_mu \approx u \parallel v$, there exists by Lemma 2 an i such that $a_iu \approx av$; hence by Lemma 7 $u \approx v$. Since $[v] = 1$ it follows that $tu \approx v \parallel v \approx vv \approx vu$, hence by Lemma 7 $t \approx v$.

If $[u] > 1$, then there exist $b_1, \dots, b_n \in A$ and ground terms u_1, \dots, u_n such that $u \approx b_1u_1 + \dots + b_nu_n$. Then $a_1u + \dots + a_mu \approx tu \approx u \parallel v \approx b_1(u_1 \parallel v) + \dots + b_n(u_n \parallel v) + vu$, so by Lemma 2 $u_i \parallel v \approx u$, for all $1 \leq i \leq n$. By Lemma 4 there exists u' such $u_i \approx u'$ for all $1 \leq i \leq n$, and, by A4, $(b_1 + \dots + b_n)u' \approx u \approx u' \parallel v$. Hence by the induction hypothesis $v \approx b_1 + \dots + b_n$ and $u' \approx v^k$ for some $k \geq 1$. So $u \approx v^{k+1}$, and from

$$\begin{aligned} tu &\approx vu + b_1(u' \parallel v) + \dots + b_n(u' \parallel v) \\ &\approx vu + vu && \text{(by A4)} \\ &\approx vu && \text{(by A3)} \end{aligned}$$

it follows, by Lemma 7, that $t \approx v$. This completes the proof of our claim.

The proof of the lemma is by induction on $[v]$. If $[t], [v] = 1$ then t is a finite sum of elements of A and by our claim $u \approx t^k$ for some $k \geq 1$. If $[t], [v] > 1$, then there exists $a \in A$ and ground terms t' and v' such that $av' \preceq v$ and $t'u \approx u \parallel v'$; hence, by the induction hypothesis, there exists a finite sum α of elements of A such that $u \approx \alpha^k$, for some $k \geq 1$. □

Lemma 10 has the following consequence.

Lemma 11. *If t, t', u and v are ground terms such that $\text{PA}_A \vdash tu \approx t'u \parallel v$, then there exists a finite sum α of elements of A such that $\text{PA}_A \vdash u \approx \alpha^k$ for some $k \geq 1$.*

Lemma 12. *Let α be a finite sum of elements of A ; if t, u and v are ground terms such that $\text{PA}_A \vdash t\alpha^k \approx u \parallel v$ for some $k \geq 1$, then $\text{PA}_A \vdash u \approx \alpha^l$ for some $l \leq k$, or there exists a ground term t' such that $\text{PA}_A \vdash u \approx t'\alpha^k$.*

Proof. The proof is by induction on the norm of v .

If $[v] = 1$, then there exists an $a \in A$ such that $a \preceq v$, whence $au \preceq t\alpha^k$. If $a \preceq t$, then $u \approx \alpha^k$, and if there exists a ground term t' such that $at' \preceq t$, then $u \approx t'\alpha^k$.

Suppose that $[v] > 1$ and let v' be a ground term such that $[v'] < [v]$ and $av' \preceq v$, whence $a(u \parallel v') \preceq t\alpha^k$. If $a \preceq t$, then $u \parallel v' \approx \alpha^k$, hence there exists an $l < k$ such that $u \approx \alpha^l$. Otherwise, suppose that t^* is a ground term such that $at^* \preceq t$ and $u \parallel v' \approx t^*\alpha^k$; by induction hypothesis $u \approx \alpha^l$ for some $l \leq k$, or there exists a ground term t' such that $u \approx t'\alpha^k$. □

Hirshfeld and Jerrum (1998) give a thorough investigation of a particular kind of mixed equations; we shall adapt some of their theory to our setting.

Let α be a finite sum of elements of A . A ground term t we shall call α -free if $t \not\approx \alpha$ and there exists no ground term t' such that $t \approx t' \parallel \alpha$. We shall call a ground term t an α -term if $t \approx \alpha^k$ for some $k \geq 1$. The α -norm $[t]_\alpha$ of a ground term t is the length of the shortest reduction of t to an α -term, or the norm of t if such a reduction does not exist. Note that if $t \approx u$, then $[t]_\alpha = [u]_\alpha$; the α -norm of an equation is the α -norm of both sides. We shall write $t \rightarrow_\alpha t'$ if $t \rightarrow t'$ and $[t']_\alpha < [t]_\alpha$; if $[u]_\alpha = 1$, then we say that u is an α -unit. In line with Definition 8, a ground term t' is an α -reduct of a ground term t if t' is reachable from t by an α -reduction.

It is easy to see that $[t \parallel u]_\alpha = [t]_\alpha + [u]_\alpha$, so we have the following lemma.

Lemma 13. *Let α be a finite sum of elements of A ; any α -free α -unit is parallel prime.*

Lemma 14. *If t is α -free, then $t\alpha$ is α -free.*

Hirshfeld and Jerrum (1998) proved a variant of Lemma 9.

Lemma 15. *Let α be a finite sum of elements of A , and let t be an α -free ground term. If $t \rightarrow_\alpha t'$ and $t \rightarrow_\alpha t''$ implies that $\text{PA}_A \vdash t' \approx t''$ for all ground terms t' and t'' , then there exists a parallel prime ground term t^* such that $\text{PA}_A \vdash t \approx t^* \parallel \dots \parallel t^*$.*

A pumpable equation is a mixed equation of the form

$$(t_1 \parallel \dots \parallel t_m)\alpha^k \approx u_1\alpha^k \parallel \dots \parallel u_n\alpha^k,$$

where α is a finite sum of elements of A , $k \geq 1$, $m, n \geq 2$ and t_i and u_j are α -free ground terms for $1 \leq i \leq m$ and $1 \leq j \leq n$. The following lemma occurs in Hirshfeld and Jerrum (1998) as Lemma 7.2.

Lemma 16. *There are no pumpable equations with α -norm less than three.*

Proposition 17. *Let t, u, u' and v be ground terms such that t and v are α -free and*

$$\text{PA}_A \vdash (t \parallel u)\alpha^k \approx v\alpha^k \parallel u'\alpha^k. \tag{1}$$

If u and u' are α -units, then $\text{PA}_A \vdash u \approx u'$.

Proof. If there exists a ground term u^* such that $u \approx u^* \parallel \alpha$, then by Lemma 3 $v\alpha^k \parallel u'\alpha^k \approx (t \parallel u^* \parallel \alpha)\alpha^k \approx (t \parallel u^*)\alpha^k \parallel \alpha$; by Lemma 14 $v\alpha^k$ is α -free, hence there exists a ground term u^{**} such that $u' \approx u^{**} \parallel \alpha$. Vice versa, from $u' \approx u^{**} \parallel \alpha$ we obtain the existence of a u^* such that $u \approx u^* \parallel \alpha$. In both cases $(t \parallel u^*)\alpha^k \parallel \alpha \approx v\alpha^k \parallel u^{**}\alpha^k \parallel \alpha$, whence

$$(t \parallel u^*)\alpha^k \approx v\alpha^k \parallel u^{**}\alpha^k.$$

Hence, we may assume without loss of generality that the α -units u and u' are α -free, so that (1) is a *pumpable equation*. By Lemma 16 there are no pumpable equations with α -norm less than three, so $[t]_\alpha, [v]_\alpha \geq 2$; we prove the lemma by induction on $[t]_\alpha$.

If there exist ground terms t' and v' such that $t \rightarrow_\alpha t', v \rightarrow_\alpha v'$ and $(t' \parallel u)\alpha^k \approx v'\alpha^k \parallel u'\alpha^k$, then we may conclude $u \approx u'$ from the induction hypothesis. Since the α -units $u'\alpha^k$ and u have unique immediate α -reducts, in the case that remains, t and v have unique immediate α -reducts t' and v' , respectively; hence, by Lemma 15 there exists a parallel prime ground term v^* such that $v \approx v^* \parallel \dots \parallel v^*$. By Lemma 3

$$(t' \parallel u)\alpha^k \approx v\alpha^k \parallel \alpha^{k+i} \approx (v \parallel \alpha^{k+i})\alpha^k, \text{ for some } i \geq 0,$$

so $t' \parallel u \approx v^* \parallel \dots \parallel v^* \parallel \alpha^{k+i}$. Since u is α -free, whence parallel prime by Lemma 14, it follows that $u \approx v^*$; hence

$$(t \parallel u)\alpha^k \approx (u \parallel \dots \parallel u)\alpha^k \parallel u'\alpha^k \text{ and } t' \approx u \parallel \dots \parallel u \parallel \alpha^{k+i}.$$

Clearly, there exists a $j \geq k$ such that $u'\alpha^k \parallel \alpha^j$ is an α -reduct of $v\alpha^k \parallel u'\alpha^k$ (α -reduce $v\alpha^k$ to α^j). Hence, by Lemma 3, $(u' \parallel \alpha^j)\alpha^k$ is an α -reduct of $(t \parallel u)\alpha^k$, so $u' \parallel \alpha^j$ is an α -reduct of $t \parallel u$. If $u' \parallel \alpha^j$ is obtained by reducing t to an α -term, then $u \approx u'$ follows, since u and u' are α -free. Otherwise, there exists $j' \leq j$ such that $u' \parallel \alpha^{j'}$ is an α -reduct of t , hence of the unique immediate α -reduct t' of t . Every α -reduct of t' with α -norm 1 is of the form $u \parallel \alpha^{j''}$. Since u and u' are parallel prime, $u \approx u'$ follows. \square

5 Mixed Terms

We shall now define the set of *head normal forms*, thus restricting the set of terms that we need to consider in our proof that PA_A is ω -complete. The syntactic form of head normal form motivate our investigation of a particular kind of terms that we shall call *mixed terms* (nestings of parallel and sequential compositions). We shall work towards a theorem that certain instantiations of mixed terms are either parallel prime or a parallel composition of a parallel prime and α^k for some finite sum α of elements of A .

Let x be a variable, suppose $t = x$ or $t = xt'$ for some term t' , and suppose $\bar{u} = u_1, \dots, u_j$ and $\bar{v} = v_1, \dots, v_j$ are sequences of terms; we define the set of *x-prefixes* $L_j[t, \bar{u}, \bar{v}]$ inductively as follows:

$$L_0[t] = t; \text{ and} \\ L_{j+1}[t, \bar{u}, u_{j+1}, \bar{v}, v_{j+1}] = (L_j[t, \bar{u}, \bar{v}] \parallel u_{j+1})v_{j+1}.$$

A term t is a *head normal form* if there exist finite sets I, J, K and L such that

$$t \approx \sum_{i \in I} a_i t_i + \sum_{j \in J} b_j + \sum_{k \in K} v_k \parallel u_k + \sum_{l \in L} w_l \quad (\text{by A1 and A2})$$

with the a_i and b_j elements of A , the t_i and u_k arbitrary terms and each v_k and w_l an x -prefix for some variable x .

Lemma 18. *For each term t there exists a head normal form t^* such that $\text{PA}_A \vdash t \approx t^*$.*

We shall associate with every equation $t \approx u$ a substitution σ such that $t^\sigma \approx u^\sigma$ implies that $t \approx u$. The main idea of our ω -completeness proof is to substitute for every variable in t or u a ground term that has a subterm φ_n of degree n , where, intuitively, n is large compared to the degrees already occurring in t and u . Let a be an element of A and let $n \geq 1$; we define

$$\varphi_n = a^n + a^{n-1} + \dots + a.$$

Lemma 19. *If $n \geq 2$ and t is a ground term, then $\varphi_n t$ is parallel prime.*

Suppose t is a term, and let $\bar{u} = u_1, \dots, u_j$ and $\bar{v} = v_1, \dots, v_j$ be sequences of terms; we define the set of *mixed terms* $M_j[t, \bar{u}, \bar{v}]$ inductively as follows:

$$M_0[t] = t; \text{ and} \\ M_{j+1}[t, \bar{u}, u_{j+1}, \bar{v}, v_{j+1}] = (M_j[t, \bar{u}, \bar{v}] \parallel u_{j+1})v_{j+1}.$$

Let t be a ground term; we denote by $d_{\max}^{\rightarrow}(t)$ the least upperbound for the degrees of all the reducts of t , i.e.,

$$d_{\max}^{\rightarrow}(t) = \max\{d(t') \mid t' \in \text{red}(t)\}.$$

Definition 20. *A mixed term $M_j[\varphi_n t, \bar{u}, \bar{v}]$ we shall call a generalised φ_n -term if*

$$d_{\max}^{\rightarrow}(M_j[t, \bar{u}, \bar{v}]) < n.$$

Note that there are no generalised φ_1 -terms.

Lemma 21. *Let $M_j[\varphi_n t, \bar{u}, \bar{v}]$ be a generalised φ_n -term and let u be a ground term such that*

$$\text{PA}_A \vdash M_j[\varphi_n t, \bar{u}, \bar{v}] \approx \varphi_n t v_1 \cdots v_j \parallel u.$$

Then there exists a finite sum α of elements of A such that $\text{PA}_A \vdash v_1 \cdots v_j \approx \alpha^k$ and $\text{PA}_A \vdash u \approx u_1 \parallel \cdots \parallel u_j \approx \alpha^l$ for some $k, l \geq 1$.

Proof. From $M_j[\varphi_n t, \bar{u}, \bar{v}] \longrightarrow M_j[t, \bar{u}, \bar{v}]$ and $d(M_j[t, \bar{u}, \bar{v}]) < n$ it follows that $M_j[t, \bar{u}, \bar{v}] \approx t v_1 \cdots v_j \parallel u$; hence

$$d_{\max}^{\rightarrow}(t v_1 \cdots v_j \parallel u) < n. \tag{2}$$

Note that $\lfloor u \rfloor = \lfloor u_1 \parallel \dots \parallel u_j \rfloor$; we shall prove the lemma by induction on $\lfloor u \rfloor$.

If $\lfloor u \rfloor = 1$, then $j = 1$ and $\lfloor u_1 \rfloor = 1$. By Lemma 11 there exists a finite sum α of elements of A such that $v_1 \approx \alpha^k$ for some $k \geq 1$, and hence by Lemma 12 $u \approx \alpha$. By Lemma 3 $(\varphi_n t \parallel u_1)\alpha^k \approx \varphi_n t \alpha^k \parallel \alpha \approx (\varphi_n t \parallel \alpha)\alpha^k$, so $u_1 \approx \alpha$.

If $\lfloor u \rfloor > 1$, then there are three cases: $\lfloor u_j \rfloor = 1$ and $j > 1$, $\lfloor u_j \rfloor > 1$ and $j = 1$, and $\lfloor u_j \rfloor > 1$ and $j > 1$. We shall only treat the last case; for the other two cases the proof is similar.

Let u'_j be an immediate reduct of u_j ; by (2) there exists an immediate reduct u' of u such that

$$M_j[\varphi_n t, u_1, \dots, u_{j-1}, u'_j, \bar{v}] \approx \varphi_n t v_1 \cdots v_j \parallel u'.$$

Since $M_j[\varphi_n t, u_1, \dots, u_{j-1}, u'_j, \bar{v}]$, there exists by the induction hypothesis a finite sum α of elements of A such that $v_1 \cdots v_j \approx \alpha^k$ and $u_1 \parallel \dots \parallel u_{j-1} \parallel u'_j \approx \alpha^l$ for some $k, l \geq 1$. By means of $j - 1$ applications of Lemma 3, we find that

$$M_j[\varphi_n t, \bar{u}, \bar{v}] \approx (\varphi_n t \alpha^{k-\lfloor v_j \rfloor} \parallel u_j) \alpha^{\lfloor v_j \rfloor} \parallel \alpha^l \approx \varphi_n t \alpha^k \parallel u.$$

Since $\varphi_n t \alpha^k$ is parallel prime by Lemma 19, there exists a ground term u^* with $\lfloor u^* \rfloor = \lfloor u_j \rfloor$ and $\lfloor u^* \rfloor_\alpha = \lfloor u_j \rfloor_\alpha$ such that $u \approx u^* \parallel \alpha^l$ and $(\varphi_n t \alpha^{k-\lfloor v_j \rfloor} \parallel u_j) \alpha^{\lfloor v_j \rfloor} \approx \varphi_n t \alpha^k \parallel u^*$. If $\lfloor u^* \rfloor \leq \lfloor v_j \rfloor$, then by Lemma 12 u^* is an α -term, which implies that u and u_j are also α -terms. So suppose that $\lfloor u^* \rfloor > \lfloor v_j \rfloor$, and let u^\dagger be a ground term such that $u^* \approx u^\dagger \alpha^{\lfloor v_j \rfloor}$. Since $\lfloor u_j \rfloor > \lfloor u^\dagger \rfloor$ and $\lfloor u_j \rfloor_\alpha = \lfloor u^\dagger \rfloor_\alpha$, by Proposition 17, u_j and u^\dagger are not α -units. Since u'_j is an α -term and $u_j \rightarrow u'_j$, u_j must be an α -term, so also u^\dagger is an α -term. Consequently, $u \approx u^* \parallel \alpha^l \approx u^\dagger \alpha^{\lfloor v_j \rfloor} \parallel \alpha^l$ is an α -term. □

Lemma 22. *Let $M_j[\varphi_n t, \bar{u}, \bar{v}]$ be a generalised φ_n -term. If $M_j[\varphi_n t, \bar{u}, \bar{v}]$ is not α -free, then there exists $i \leq j$ such that $\text{PA}_A \vdash v_i \cdots v_j \approx \alpha^k$ and u_i is not α -free.*

Proof. Let t^* be a ground term such that

$$M_j[\varphi_n t, \bar{u}, \bar{v}] \approx t^* \parallel \alpha. \tag{3}$$

Since $\varphi_n t v_1 \cdots v_j$ is a reduct of $M_j[\varphi_n t, \bar{u}, \bar{v}]$ and by Lemma 19 $\varphi_n t v_1 \cdots v_j$ is parallel prime, $\varphi_n t v_1 \cdots v_j$ must be a reduct of t^* . Then $\varphi_n t v_1 \cdots v_j \parallel \alpha$ is a reduct of $M_j[\varphi_n t, \bar{u}, \bar{v}]$, so there exist sequences of ground terms \bar{u}' and \bar{v}' such that for some $1 \leq j' \leq j$ and $1 \leq i \leq j$,

$$M_{j'}[\varphi_n t v_1 \cdots v_{i-1}, \bar{u}', \bar{v}'] \approx \varphi_n t v_1 \cdots v_j \parallel \alpha, \text{ where } v'_1 \cdots v'_{j'} \approx v_i \cdots v_j.$$

By Lemma 21 $v_i \cdots v_j \approx \alpha^k$, so in particular $v_j \approx \alpha^l$ for some $l \leq k$. Clearly, $\lfloor t^* \rfloor > l$, so by Lemma 12 there exists t^\dagger such that $t^* \approx t^\dagger \alpha^l$. We apply Lemma 3 to the right-hand side of (3) and cancel the α^l -tail on both sides to obtain

$$M_{j-1}[\varphi_n t, u_1, \dots, u_{j-1}, v_1, \dots, v_{j-1}] \parallel u_j \approx t^\dagger \parallel \alpha.$$

The remainder of the proof is by induction on j . If $j = 1$, then $\varphi_n t \parallel u_j \approx t^\dagger \parallel \alpha$ implies that u_j is not α -free and we are done. If $j > 1$ and u_j is α -free, then $M_{j-1}[\varphi_n t, u_1, \dots, u_{j-1}, v_1, \dots, v_{j-1}]$ is not α -free, so by the induction hypothesis there exists some $1 \leq i' \leq j - 1$ such that $u_{i'}$ is not α -free and $v_{i'} \cdots v_{j-1} \approx \alpha^{k'}$, whence $v_{i'} \cdots v_j \approx \alpha^{k'+1}$. \square

Proposition 23. *If a generalised φ_n -term t^* is not parallel prime, then there exists a finite sum α of elements of A and a parallel prime generalised φ_n -term t^\dagger such that $t^* \approx t^\dagger \parallel \alpha^k$ for some $k \geq 1$.*

Proof. Let $t^* \approx M_j[\varphi_n t, \bar{u}, \bar{v}]$ and let t_1, \dots, t_o be parallel prime ground terms such that

$$M_j[\varphi_n t, \bar{u}, \bar{v}] \approx t_1 \parallel \dots \parallel t_o.$$

Since $\varphi_n t v_1 \cdots v_j$ is a reduct of $M_j[\varphi_n t, \bar{u}, \bar{v}]$ and parallel prime, $\varphi_n t v_1 \cdots v_j$ must be a reduct of some t_i ($1 \leq i \leq o$); assume without loss of generality that it is a reduct of t_1 .

Suppose that $M_j[\varphi_n t, \bar{u}, \bar{v}]$ is not parallel prime and let $u \approx t_2 \parallel \dots \parallel t_o$. Since $\varphi_n t v_1 \cdots v_j \parallel u$ is a reduct of $M_j[\varphi_n t, \bar{u}, \bar{v}]$, there exist sequences of ground terms \bar{u}' and \bar{v}' such that, for some $1 \leq j' \leq j$ and $1 \leq i \leq j$,

$$M_{j'}[\varphi_n t v_1 \cdots v_{i-1}, \bar{u}', \bar{v}'] \approx \varphi_n t v_1 \cdots v_j \parallel u, \text{ where } v'_1 \cdots v'_{j'} \approx v_i \cdots v_j.$$

So by Lemma 21, there exists a finite sum α of elements of A such that $u \approx \alpha^k$, for some $k \geq 1$.

It remains to prove that if $M_j[\varphi_n t, \bar{u}, \bar{v}] \approx t^\dagger \parallel \alpha$, then t^\dagger is a generalised φ_n -term, for then it follows that t_1 is a generalised φ_n -term by induction on k . Since $M_j[\varphi_n t, \bar{u}, \bar{v}]$ is not α -free, there exists by Lemma 22 an $i \leq j$ such that u_i is not α -free and $v_i \cdots v_j \approx \alpha^l$ for some $l \geq 1$. So either $u_i \approx \alpha$ or there exists u'_i such that $u_i \approx u'_i \parallel \alpha$; we only consider the second possibility, as the other can be dealt with similarly. By Lemma 3 we obtain

$$M_j[\varphi_n t, \bar{u}, \bar{v}] \approx M_{j-1}[\varphi_n t, u_1, \dots, u_{i-1}, u'_i, u_{i+1}, \dots, u_j, \bar{v}] \parallel \alpha$$

Hence $t^\dagger \approx M_{j-1}[\varphi_n t, u_1, \dots, u_{i-1}, u'_i, u_{i+1}, \dots, u_j, \bar{v}]$ is a generalised φ_n -term. \square

6 ω -Completeness

Let A be a nonempty set; we shall now prove that PA_A is ω -complete. We shall assume that the variables used in an equation $t \approx u$ are enumerated by $x_1, x_2, \dots, x_k, \dots$. Let x_i be a variable and let m be a natural number; the particular kind of substitutions σ_m that we shall use in our proof satisfy

$$\sigma_m(x_i) = a(a\varphi_{i+m} + a)a.$$

We want to choose m large compared to the degrees already occurring in t and u ; with every term t we associate a natural number $d_{\max}^\sigma(t)$ that denotes the maximal degree that occurs in t after applying a substitution of the form described above, treating the terms φ_{i+m} as fresh constants.

Definition 24. *Suppose $\Xi = \{\xi_1, \xi_2, \dots, \xi_k, \dots\}$ is a countably infinite set of constant symbols such that $\Xi \cap A = \emptyset$. Let t be a term and let σ be a substitution such that*

$$\sigma(x_i) = a(a\xi_i + a)a.$$

We define $d_{\max}^\sigma(t)$ as the maximal degree that occurs in t^σ , i.e., $d_{\max}^\sigma(t) = d_{\max}^\sigma(t^\sigma)$.

Lemma 25. *If t is a term and let m be a natural number, then*

- i. $d_{\max}^{\rightarrow}(t^{\sigma^m}) \leq d_{\max}^\sigma(t)$; and
- ii. if $a \in A$ and t' is a ground term such that $at' \preceq t^{\sigma^m}$, then $d(t') \leq d_{\max}^\sigma(t)$.

If $L_j[t, \bar{u}, \bar{v}]$ is an x_i -prefix and m is a natural number, then

$$L_j[t, \bar{u}, \bar{v}]^{\sigma^m} \longrightarrow M_j[(a\varphi_{i+m} + a)t'', \bar{u}, \bar{v}]^{\sigma^m} \succcurlyeq aM_j[\varphi_{i+m}t'', \bar{u}, \bar{v}]^{\sigma^m}, \quad (4)$$

where $t'' = a$ if $t = x_i$ and $t'' = at'$ if $t = x_it'$ for some term t' . If $m \geq d_{\max}^\sigma(L_j[t, \bar{u}, \bar{v}])$, then $M_j[\varphi_{i+m}t'', \bar{u}, \bar{v}]^{\sigma^m}$ is a generalised φ_n -term; we shall call it the *generalised φ_{i+m} -term associated with $L_j[t, \bar{u}, \bar{v}]$ by σ_m* .

For ground terms t and t^* , let us write $t \mapsto t^*$ if there exists a ground term t' and an $a \in A$ such that $t \longrightarrow t' \succcurlyeq at^*$.

Lemma 26. *Let t be an x -prefix and suppose that $m \geq d_{\max}^\sigma(t)$. If $n > m$ and t^* and t^\dagger are generalised φ_n -terms such that $t^{\sigma^m} \mapsto t^*$ and $t^{\sigma^m} \mapsto t^\dagger$, then $\text{PA}_A \vdash t^* \approx t^\dagger$.*

Proof. Note that the unique immediate reduct of t^{σ^m} is of the form $M_j[(a\varphi_{i+m} + a)t', \bar{u}, \bar{v}]^{\sigma^m}$. Moreover, $m \geq d_{\max}^\sigma(t)$, so $M_j[\varphi_{i+m}t', \bar{u}, \bar{v}]^{\sigma^m}$ is the unique ground term t^* such that $at^* \preceq M_j[(a\varphi_{i+m} + a)t', \bar{u}, \bar{v}]^{\sigma^m}$ and $d(t^*) > m$. Hence, if t^* is any generalised φ_n -term with $n > d_{\max}^\sigma(t)$ such that $t^{\sigma^m} \mapsto t^*$, then $t^* \approx M_j[\varphi_{i+m}t', \bar{u}, \bar{v}]^{\sigma^m}$.

Note that if t is an x -prefix, $m \geq d_{\max}^\sigma(t)$ and t^* is the generalised φ_{i+m} -term associated with t by σ_m , then t^* has no reduct with a degree in $\{m + 1, m + 2, \dots, m + i - 1\}$. Lemma 26 has the following consequence.

Lemma 27. *Let t be an x -prefix, let u be a y -prefix and let $m \geq \max\{d_{\max}^\sigma(t), d_{\max}^\sigma(u)\}$. If $\text{PA}_A \vdash t^{\sigma^m} \approx u^{\sigma^m}$, then $x = y$.*

We generalise the definition of α -freeness to terms with variables: a term t is α -free if $t \not\approx \alpha$ and there exists no term t' such that $t \approx t' \parallel \alpha$.

Theorem 28 (ω -completeness). *Let A be a nonempty set; then PA_A is ω -complete, i.e., for all terms t and u ,*

if $\text{PA}_A \vdash t^\sigma \approx u^\sigma$ for all ground substitutions σ , then $\text{PA}_A \vdash t \approx u$.

Proof. Let $m \geq \max\{d_{\max}^\sigma(t), d_{\max}^\sigma(u)\}$. We shall prove by induction on the depth of t that if $t^{\sigma_m} \approx u^{\sigma_m}$, then $t \approx u$; clearly, this implies the theorem.

In the full version of this paper we simultaneously prove that one may assume without loss of generality that the generalised φ_n -term associated by σ_m with an x -prefix is parallel prime; we need this assumption in the remainder of the proof.

Suppose that $t^{\sigma_m} \approx u^{\sigma_m}$; it suffices to show that every summand of u is a summand of t , for then by a symmetric argument it follows that every summand of t is a summand of u , whence $t \approx u$. There are four cases:

1. If $b \in A$ such that $b \preceq u$, then $b \preceq t$ since σ_m -instances of summands of one of the three other types have a norm > 1 .
2. If $a \in A$ and u' is a term such that $au' \preceq u$, then by Lemma 25 $d(u') \leq d_{\max}^\sigma(u)$. Since $m \geq d_{\max}^\sigma(u)$, $(u')^{\sigma_m}$ cannot be an immediate reduct of a σ_m -instance of an x -prefix or of a term $v \parallel w$, with v an x -prefix. So there exists $i \in I$ such that $a(u')^{\sigma_m} \approx a_i t_i^{\sigma_m}$. By the induction hypothesis $u' \approx t_i$, hence $au' \preceq t$.
3. Let v be an x -prefix and let u' is a term such that $v \parallel u' \preceq u$. By our assumption that generalised φ_n -terms associated by σ_m with x -prefixes are parallel prime, there exists $k \in K$ such that $v^{\sigma_m} \approx v_k^{\sigma_m}$ and $(u')^{\sigma_m} \approx u_k^{\sigma_m}$; by Lemma 27 v_k is also an x -prefix. Hence, by the induction hypothesis $v \approx v_k$ and $u' \approx u_k$, so $v \parallel u \preceq t$.
4. If w is an x -prefix such that $w \preceq u$, then by our assumption that generalised φ_n -terms associated by σ_m with x -prefixes are parallel prime, there exists $l \in L$ such that $w^{\sigma_m} \approx w_l^{\sigma_m}$; by Lemma 27 w_l is also an x -prefix. If the generalised φ_n -term associated to w by σ_m is of the form $\varphi_n w'$, then clearly the generalised φ_n -term associated to w_l by σ_m must be of the form $\varphi_n w'_l$ and it is immediate by the induction hypothesis that $w' \approx w'_l$ and $w \approx w_l$. Let $w = (t' \parallel u')v'$ and let $w_l = (t'' \parallel u'')v''$, where t' and t'' are x -prefixes to which σ_m associates parallel prime generalised φ_n -terms t^\dagger and t^\ddagger . If $\lceil (v')^{\sigma_m} \rceil = \lceil (v'')^{\sigma_m} \rceil$, then by the induction hypothesis $(t' \parallel u') \approx (t'' \parallel u'')$ and $v' \approx v''$, whence $w \approx w_l$. So let us assume without loss of generality that $\lceil (v')^{\sigma_m} \rceil < \lceil (v'')^{\sigma_m} \rceil$; then there is a ground term v^* such that $(t' \parallel u')^{\sigma_m} \approx (t'' \parallel u'')^{\sigma_m} v^* \approx (v'')^{\sigma_m}$. Note that $(t^\ddagger \parallel u'')^{\sigma_m} v^* \approx (t^\dagger \parallel u')^{\sigma_m}$, which is not parallel prime. So there exists by Proposition 23 a finite sum α of elements of A and a parallel prime generalised φ_n -term t^* such that $(t^\ddagger \parallel (u'')^{\sigma_m})v^* \approx t^* \parallel \alpha^k$. Hence by Lemma 22 $v^* \approx \alpha^l$ for some $l \geq 1$. Consequently, $v'' \approx \alpha^l v'$ and by the induction hypothesis $t' \parallel u' \approx (t'' \parallel u'')\alpha^l$; hence, $w \preceq t$. □

References

- Aceto, L. and Hennessy, M. (1993). Towards action-refinement in process algebras. *Information and Computation*, **103**(2), 204–269.
- Baeten, J. C. M. and Weijland, W. P. (1990). *Process Algebra*. Number 18 in Cambridge Tracts in Theoretical Computer Science. Cambridge University Press.
- Bergstra, J. A. and Klop, J. W. (1984). Process algebra for synchronous communication. *Information and Control*, **60**(1–3), 109–137.
- Castellani, I. and Hennessy, M. (1989). Distributed bisimulations. *Journal of the ACM*, **36**(4), 887–911.
- Groote, J. F. (1990). A new strategy for proving ω -completeness applied to process algebra. In J. Baeten and J. Klop, editors, *Proceedings of CONCUR'90*, volume 458 of *LNCS*, pages 314–331, Amsterdam. Springer-Verlag.
- Heering, J. (1986). Partial evaluation and ω -completeness of algebraic specifications. *Theoretical Computer Science*, **43**, 149–167.
- Hennessy, M. (1988). Axiomatising finite concurrent processes. *SIAM Journal of Computing*, **17**(5), 997–1017.
- Hirshfeld, Y. and Jerrum, M. (1998). Bisimulation equivalence is decidable for normed process algebra. LFCS Report ECS-LFCS-98-386, University of Edinburgh.
- Hirshfeld, Y. and Jerrum, M. (1999). Bisimulation equivalence is decidable for normed process algebra. In J. Wiedermann, P. van Emde Boas, and M. Nielsen, editors, *Proceedings of ICALP'99*, volume 1644 of *LNCS*, pages 412–421, Prague. Springer.
- Lazrek, A., Lescanne, P., and Thiel, J.-J. (1990). Tools for proving inductive equalities, relative completeness, and ω -completeness. *Information and Computation*, **84**(1), 47–70.
- Milner, R. (1989). *Communication and Concurrency*. Prentice-Hall International, Englewood Cliffs.
- Milner, R. and Moller, F. (1993). Unique decomposition of processes. *Theoretical Computer Science*, **107**, 357–363.
- Moller, F. (1989). *Axioms for Concurrency*. Ph.D. thesis, University of Edinburgh.
- Moller, F. (1990). The importance of the left merge operator in process algebras. In M. Paterson, editor, *Proceedings of ICALP'90*, volume 443 of *LNCS*, pages 752–764, Warwick. Springer.